

# Taxonomic identification and distribution of *Lutzomyia* spp. in Cauca River Canyon municipalities of the Hidroituango project

## Identificación taxonómica y distribución de *Lutzomyia* spp., en el cañón del río Cauca, municipios del proyecto Hidroituango

Giovani Zapata-Úsuga<sup>1\*</sup>, Wilber Gómez-Vargas<sup>1,2</sup>, Paula Mejía-Salazar<sup>1</sup>, Boris Zuleta-Ruiz<sup>1</sup>, Walter Zuluaga-Ramírez<sup>1</sup>

### Abstract

In hydroelectric projects, few investigations have been carried out to determine the taxonomic composition of diptera of the family Psychodidae, subfamily Phlebotominae. In this work, a taxonomic inventory of phlebotomine sandflies was conducted in the area of influence of the Ituango hydroelectric power plant, in the department of Antioquia, Colombia. We highlight their medical importance as vectors of the protozoan of the genus *Leishmania* and update their distribution. Entomological monitoring was carried out in ten municipalities from the western subregion (Santa Fe de Antioquia, Peque, Olaya, Liborina and Sabanalarga) and the northern subregion (the municipalities of Ituango, Briceño, Valdivia, Toledo and San Andrés de Cuerquia). CDC traps and Shannon traps were used for monitoring. For taxonomic identification, the keys of Young and Duncan and Galati were used. A total of 7993 phlebotomine sandflies were collected, distributed in 39 species according to Galati, highlighting six species considered vectors of *Leishmania* spp. that causes cutaneous leishmaniasis, which are: *Lutzomyia* (*Trl.*) *gomezi*, *Lu.* (*Hel.*) *hartmanni*, *Psychodopygus panamensis*, *Pintomyia* (*Pif.*) *columbiana*, *Nyssomyia trapidoi* and *Ny. yuilli yuilli*. We conclude that the sampling area is endemic for leishmaniasis due to the presence of vectors transmitting this parasite. The inventory on the distribution of phlebotomine sandflies will serve as a basis for further studies that allow the implementation of surveillance and control strategies.

**Keywords:** hydroelectric, *Leishmania* spp., Phlebotomine, protozoan, vectors

### Resumen

En proyectos hidroeléctricos se han realizado pocas investigaciones para determinar la composición taxonómica de dípteros de la familia Psychodidae, subfamilia Phlebotominae. En este trabajo se realizó un inventario taxonómico de flebotomos en el área de influencia de la central hidroeléctrica de Ituango, departamento de Antioquia con el fin de actualizar su distribución y resaltar su importancia médica como vectores del protozoo del género *Leishmania*. Para esto, se realizaron monitoreos entomológicos con trampas CDC y trampas Shannon, en diez municipios de la subregión occidente (Santa Fe de Antioquia, Peque, Olaya, Liborina y Sabanalarga) y de la subregión norte (municipios de Ituango, Briceño, Valdivia, Toledo y San Andrés de Cuerquia). Para la identificación taxonómica se utilizaron las claves de Young y Duncan y Galati. En total se recolectaron 7993 flebotomos, distribuidos en 39 especies según Galati, destacándose seis especies consideradas vectores de especies de *Leishmania*, que causan la leishmaniasis cutánea. Estas especies fueron *Lutzomyia* (*Trl.*) *gomezi*, *Lu.* (*Hel.*) *hartmanni*, *Psychodopygus panamensis*,

<sup>1</sup> National School of Public Health, University of Antioquia, Medellín, Colombia.

<sup>2</sup> Colombian Institute of Tropical Medicine, CES University, Medellín, Colombia.

\* Corresponding author: [giovani.zapata@udea.edu.co](mailto:giovani.zapata@udea.edu.co)

Received: January 16, 2023; accepted: August 24, 2023; published: November 29, 2023.

*Pintomyia (Pif.) columbiana*, *Nyssomyia trapidoi* y *Ny. yuilli yuilli*. Los datos indican que la zona de muestreo es endémica para la leishmaniasis debido a la presencia de vectores transmisores de este parásito. El inventario sobre la distribución de flebotomos servirá de base para estudios posteriores que permitan la implementación de estrategias de vigilancia y control.

**Palabras clave:** hidroeléctrico, *Leishmania* spp., Phlebotominae, protozoo, vectores

## INTRODUCTION

Psychodidae are insects of the order Diptera (suborder Nematocera: Psychodidae: Phlebotominae), belonging to the family Psychodidae and the subfamily Phlebotominae (Young and Duncan, 1994). Phlebotomine sandflies are the main vectors of parasites of the genus *Leishmania* spp. that cause a zoonotic disease known as leishmaniasis that presents diverse clinical manifestations and its transmission occurs in tropical and subtropical areas, being endemic in 98 countries (Alvar et al., 2012; Ferro et al., 2015). There are 350 million people at risk of leishmaniasis and it is estimated that about 900,000 – 1.3 million new cases of leishmaniasis are reported each year (Pan American Health Organization [PAHO], 2023).

It is important to mention that, in Antioquia, Porter conducted the first inventory study of phlebotomine sandflies in the area of influence of the Providencia hydroelectric plant located in the municipality of Anorí in 1970. In that study, four species of the genus *Lutzomyia* (*Lu. hartmanni*, *Lu. gomezi*, *Lu. trapidoi* and *Lu. yuilli*) and one of the genus *Warileyia* (*Wa. rotundipennis*) were recorded (Porter and DeFoliart, 1981).

Another study was conducted between 1997 and 2000 by the Epidemiological Surveillance System Group (SVE) of the University of Antioquia at the Porce II hydroelectric plant, where eight *Lutzomyia* species (*Lu. barretoii*, *Lu. lichyi*, *Lu. gomezi*, *Lu. shannoni*, *Lu. panamensis*, *Lu. triramula*, *Lu. carrerai* and *Lu. walkeri*) were reported (Zuluaga et al., 2012). This group also conducted studies from 2007 to 2009 in the Porce III Hydroelectric Project, reporting 13 species of *Lutzomyia* (Zuluaga et al., 2012).

In 2003, at the ISAGEN dams in the municipality of San Carlos in Antioquia, the Colombian Institute of Tropical Medicine (ICMT) conducted a focus

study, recording the species *Lu. panamensis*, *Lu. gomezi*, *Lu. bifoliata* and *Wa. Rotundipennis*. Based on the abundance found in the sampling sites, anthropophilic behavior and vectorial antecedents, it was suggested that *Lu. panamensis* and *Lu. gomezi* are responsible for the transmission of the parasite that causes cutaneous leishmaniasis in this area (Parra-Henao and Echavarría, 2005).

In 2008 in the department of Caldas, a study of leishmaniasis outbreaks was reported in the hydroelectric plant La Miel I in the sectors of Guarinó and Manso. Five species of *Lutzomyia* were found (*Lu. longipalpis*, *Lu. gomezi*, *Lu. trapidoi*, *Lu. trinidadensis* and *Lu. cayennensis*) (Vergara et al., 2008). Another study conducted in this same plant in 2013 reported for the first time the maximum altitude of distribution of *Lu. longipalpis* at 1350 m.a.s.l. (Acosta et al., 2013).

In 2012 in the municipality of Chaparral (Tolima) in the area of influence of the Amoyá hydroelectric project (2008-2009), 13 species of phlebotomine sandflies were identified, of which the species *Lu. longiflocosa*, *Lu. columbiana*, *Lu. nuneztovari*, *Lu. suapiensis* and *Wa. rotundipennis* are epidemiologically important and *Lu. longiflocosa* has been recorded as a carrier of *Leishmania braziliensis* (Contreras et al., 2012).

In the department of Santander, during studies conducted in 2016 and 2017 in the Sogamoso hydroelectric power plant, 21 species of *Lutzomyia* were recorded, highlighting the presence of *Lu. gomezi*, *Lu. panamensis* and *Lu. ovallesi* as important transmitter species of parasites that cause cutaneous leishmaniasis. The species *Lu. longipalpis*, which transmits *Leishmania* species that cause visceral leishmaniasis, was predominant (Gómez-Vargas and Zapata-Úsuga, 2022).

Finally, in 2016 in the Urra hydroelectric plant, the presence of 12 species of *Lutzomyia* were reported, where it was also found that seven were naturally

infected with *Leishmania* spp. (*Lu. panamensis*, *Lu. gomezi*, *Lu. cayennensis*, *Lu. micropyga*, *Lu. shannoni*, *Lu. trinidadensis* and *Lu. yuilli yuilli*). Records for five new species in the department of Córdoba (*Lu. carpenteri*, *Lu. dysponeta*, *Lu. atroclavata* and *Lu. yuilli yuilli*) were also reported (Vivero et al., 2016).

In the New World, phlebotomine sandflies are involved in the propagation of protozoa of the genus *Leishmania* Ross, 1903, (Kinetoplastida: Trypanosomatidae) (Akhoundi et al., 2016; Lainson and Shaw, 1973; Vivero et al., 2015). Phlebotomine sandflies can also propagate arboviruses of the Bunyviridae family where the *Flebovirus* genus stands out, represented mainly by the phlebotomine fever (known as Panama fever in the New World and "pappataci fever" in the Old World) and arboviruses of the Rhabdoviridae family represented by the *Vesiculovirus* genus (which produces Chandipura virus encephalitis), summer meningitis (Toscana virus), vesicular stomatitis (called New Jersey stomatitis or Indian stomatitis), which affect cattle, horses, swine and humans. Additionally, phlebotomine sandflies transmit other pathogens such as the bacterium *Bartonella bacilliformis* that produces Carrion's disease (Acevedo and Arrivillaga, 2008; Maroli, et al., 2013; Vivero et al., 2015; Young, 1977). Disease transmission is due to the hematophagous behavior of females which need the blood of their endothermic vertebrate hosts (cattle, dogs, rodents and man) (Sales et al., 2015; Sant'Anna et al., 2008; Tanure et al., 2015) and exothermic vertebrates, such as reptiles and amphibians (Abbate et al., 2020; Young and Duncan, 1994), to carry out their gonotrophic cycle. Males feed preferentially on sugars produced by plants or by aphids and/or coccidia and females supplement their diet with these sugars (Bastidas et al., 2004; Cameron et al., 1994). Phlebotomine sandflies are considered telmophages, because they lacerate the skin of the animal using their mandibles, causing a small accumulation of blood from which they feed (Gonzalez, 2019).

There are 1000 species of phlebotomine sandflies, 535 which have been recorded in the Americas (Galati, 2018) and approximately 45 being vectors of different species of *Leishmania* (Cecílio et al., 2022; Gradoni, 2018). In Colombia, 167 species have been described (Ferro et al., 2015), 153 which are part of the genus *Lutzomyia* (Bejarano

and Estrada, 2016), 14 of which are confirmed as vectors, and an additional eight as potential vectors (Gómez-Vargas and Zapata-Úsuga, 2022; Méndez-Cardona, 2021).

The species confirmed as vectors in Colombia are the following: *Lu. columbiana* (Bejarano et al., 2003; Grimaldi et al., 1989; Montoya-Lerma et al., 1999; Pan American Health Organization [PAHO], 2012), *Lu. evansi* (Bejarano et al., 2002; Bejarano et al., 2003; PAHO, 2012; Travi et al., 2001; Urango and Hoyos-López, 2022), *Lu. gomezi* (Alexander et al., 2001; Bejarano et al., 2002; Montoya-Lerma et al., 1999), *Lu. hartmanni* (Alexander et al., 2001; Grimaldi et al., 1989), *Lu. lichyi* (Alexander et al., 1995; Warburg et al., 1991), *Lu. longiflocosa* (Bejarano et al., 2003; Cárdenas et al., 1999; Méndez-Cardona, 2021; Pardo et al., 1999), *Lu. longipalpis* (Corredor et al., 1989, 1990; Goenaga-Mafud et al., 2020), *Lu. ovallesi* (Alexander et al., 2001; Bejarano et al., 2003), *Lu. panamensis* (Alexander et al., 1995; Bejarano et al., 2003), *Lu. scorzai* (Alexander et al., 1995), *Lu. spinicrassa* (Morales et al., 2005), *Lu. trapidoi* (Alexander et al., 2001; Corredor et al., 1990; Martínez et al., 2018; Santamaría et al., 2006, Travi et al., 1988), *Lu. umbratilis* (Grimaldi et al., 1989), and *Lu. yuilli yuilli* (Martínez et al., 2018; Santamaría et al., 2006).

The potential vectors are the following: *Lu. antunesi* (Niño and Pérez-Español, 2021; Vásquez-Trujillo et al., 2008), *Lu. davisii* (Bejarano et al., 2006), *Lu. flaviscutellata* (Cabrera et al., 2009; PAHO, 2012), *Lu. hirsuta hirsuta* (Cabrera et al., 2009; Ministerio de Protección Social [Minsalud], 2010), *Lu. torvida*, *Lu. townsendi* (Bejarano et al., 2003), *Lu. trinidadensis* (Vivero et al., 2017) and *Lu. quasitownsendi* (Bejarano et al., 2003; Garcia-Leal et al., 2020).

For the municipalities in the area of influence of the Ituango hydroelectric power plant, the first records of leishmaniasis began in 1995 in municipalities in the northern subregion such as Briceño (eight cases), Ituango (10 cases), and Valdivia (38 cases). In the western subregion, only one case has been recorded in Sabanalarga. For the department, 895 cases were recorded that same year (Dirección Seccional de Salud de Antioquia [DSSA], 1998). In 1997, the department registered 951 cases of leishmaniasis, including municipalities in the area of influence of the Ituango hydroelectric power

plant such as Briceño (2 cases), Ituango (44 cases), Valdivia (70 cases), and Santa Fe de Antioquia (one case) (DSSA and Laboratorio Departamental de Salud [LDS], 1998). Between 1998 and 2001, there are no records for these areas.

In 2002, some municipalities in the study area were considered endemic for leishmaniasis, such as the municipality of Valdivia in the northern subregion, which registered 1825 cases of leishmaniasis between 2002 and 2011, and 1198 cases between 2012-2020. This is followed by Ituango (785 cases between 2002-2011 and 464 cases from 2012 to 2020), Briceño (204 cases from 2002 to 2011 and 121 cases from 2012 to 2020). For the municipalities of the western sub-region, Santa Fe de Antioquia recorded the most cases with a total of 458 distributed as follows: 52 cases from 2002 to 2011 and 406 cases from 2012 to 2020. Other municipalities in this subregion that reported cases were Sabanalarga (8 cases 2002 to 2011 and 284 cases from 2012 to 2020) and Buriticá (4 cases from 2002 to 2011 and 131 cases from 2012 to 2020) (DSSA, 2011, 2020). Because of this, the northern subregion of the department of Antioquia is considered to be high incidence and endemic for leishmaniasis, especially the municipalities of Briceño, Ituango and Valdivia, the latter, where focus studies have been conducted (Posada-López et al., 2014).

## MATERIALS AND METHODS

### Study area

The study was conducted in the rural area of the western subregion of the municipalities of Buriticá (6°43'12"N, 75°54'27"O), Liborina (6°40'41"N 75°48'44"O), Olaya (6°37'40"N 75°48'43"O), Sabanalarga (6°50'54"N 75°49'01"O), Peque and Santafé de Antioquia (6°33'23"N 75°49'39"O) and in the northern subregion in the municipalities of Briceño (7°6'38"N, 75°33'4"O), Valdivia (7°09'49"N 75°26'21"O), Ituango (7°10'16"N 75°45'49"O), San Andrés de Cuerquia (6°54'52"N 75°40'33"O) and Toledo (7°00'37"N 75°42'06"O), municipalities where the Hidroituango project is being executed (Figure 1). These municipalities are located in the Cauca River canyon between the central and western mountain ranges, and have two ecotopes, a dry forest type fragmented mainly with coffee and cocoa crops in the western subregion, while the northern subregion is

dominated by tropical rainforest, predominantly gallery forest, coffee, cocoa and illicit crops.

### Entomological sampling

Sampling was carried out during two nights at each site, with a CDC type light trap between 18:00 and 06:00 hours located in the intra, peri and extradomicile, and with a Shannon trap between 18:00 and 21:00 using a mouth aspirator. Monitoring was conducted every three months for eight consecutive years (years 2012-2020), including dry and rainy seasons. All specimens collected were stored in 1.5 ml tubes, with 70% alcohol, subsequently rinsed with 10% KOH. Hoyer's liquid was used for semi-permanent mounts. For taxonomic identification, the dichotomous keys of Young (1977, 1979), Young and Duncan (1994) and Galati (Galati, 2018; Galati et al., 2017) were used.


An Excel database was used for the calculations of entomological indicators for the species of medical importance. Past 3 software was used for the statistical analysis of diversity and Arcgis 10.1 and Maxent 3.4.4 software was used for the projection of species and maps.

## RESULTS

A total of 7993 phlebotomine sandfly specimens were identified (5469 specimens for the municipalities of the northern subregion and 2524 for the municipalities of the western subregion), distributed in 39 species belonging to 13 genera according to Galati's classification (Galati, 2018; Galati et al., 2017). The abundance and total number of specimens collected per municipality according to the subregion to which that municipality belongs is shown in Table 1.

Of the species collected in the sampling area (Table 1), six are implicated in the transmission of parasites of the genus *Leishmania* in Colombia: *Lu. (Trl.) gomezi*, *Lu. (Hel.). hartmanni*, *Ps. panamensis*, *Pi. (Pif.) columbiana*, *Ny. trapidoi* and *Ny. yuilli yuilli*. Figure 2 shows the spermathecae of these species as a morphological character of importance.

Given the epidemiological importance of these vector species, potential distribution maps were constructed in MaxEnt (Figure 3). For the



**Figure 1.** Map of phlebotomine species of public health concern recorded in the Cauca river canyon area, an area of influence of the Ituango Hydroelectric Project, Antioquia.

**Table 1.** Abundance of phlebotomine sandfly species collected in the ten municipalities located in the area of influence of the Ituango Hydroelectric Project, Antioquia between 2012-2020.

Specie	Northern subregion							Western subregion							
	Briceño	Ituango	San Andres de Cuerquia	Toledo	Valdivia	Total N	Pi	Buritica	Liborina	Olaya	Peque	Sabanalarga	Santa Fe de Antioquia	Total N	Pi
<i>Bi. olmeca bicolor</i>	0.00%	0.00%	0.00%	0.00%	0.04%	2	0.04%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Bi. olmeca sp</i>	0.00%	0.00%	0.00%	0.02%	0.00%	1	0.02%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Da. vespertilionis</i>	0.00%	0.02%	0.00%	0.00%	0.00%	1	0.02%	0.00%	0.00%	0.00%	0.04%	0.00%	0.00%	1	0.04%
<i>Da. rosabali</i>	0.00%	0.00%	0.00%	0.00%	0.02%	1	0.02%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Ev. dubitans</i>	0.37%	0.84%	0.00%	0.38%	0.11%	93	1.70%	0.51%	0.20%	0.28%	0.87%	1.93%	0.00%	96	3.79%
<i>Ev. walkeri</i>	0.00%	0.00%	0.00%	0.02%	0.00%	1	0.02%	0.00%	0.00%	0.00%	0.00%	0.20%	0.00%	5	0.20%
<i>Ev. saulensis</i>	0.00%	0.00%	0.00%	0.00%	0.02%	1	0.02%	0.00%	0.00%	0.12%	0.00%	0.00%	0.00%	3	0.12%
<i>Lu. cirrita</i>	0.00%	0.02%	0.00%	0.11%	0.16%	16	0.29%	2.92%	0.00%	0.00%	0.04%	0.28%	0.00%	82	3.24%
<i>Lu. hartmanni</i>	1.53%	3.63%	0.00%	4.69%	1.73%	635	11.59%	1.26%	0.24%	0.00%	2.09%	2.92%	0.08%	167	6.59%
<i>Lu. sanguinaria</i>	0.00%	0.09%	0.00%	0.00%	0.05%	8	0.15%	0.00%	0.00%	0.00%	0.04%	0.00%	0.00%	1	0.04%
<i>Lu. scorzai</i>	0.04%	0.29%	0.04%	0.07%	0.18%	34	0.62%	0.00%	0.00%	0.04%	0.28%	0.04%	0.00%	9	0.36%
<i>Lu. strictivilla</i>	0.07%	0.62%	0.02%	0.29%	0.16%	64	1.17%	0.36%	0.00%	0.00%	0.79%	0.20%	0.00%	34	1.34%
<i>Lu. bifoliata</i>	0.00%	0.00%	0.00%	0.04%	0.00%	2	0.04%	0.04%	0.00%	0.00%	0.00%	0.00%	0.00%	1	0.04%
<i>Lu. lichyi</i>	0.04%	0.00%	0.00%	0.04%	0.05%	7	0.13%	0.00%	0.12%	0.08%	0.28%	0.04%	0.00%	13	0.51%
<i>Lu. gomezi</i>	3.74%	12.80%	0.27%	14.80%	18.73%	2758	50.35%	11.40%	6.27%	10.46%	2.60%	17.32%	1.97%	1268	50.04%
<i>Lu. spp</i>	0.49%	0.88%	0.00%	0.47%	0.29%	117	2.14%	0.67%	0.32%	0.36%	0.39%	1.03%	0.00%	70	2.76%
<i>Mg. migonei</i>	0.00%	0.00%	0.00%	0.02%	0.00%	1	0.02%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Mi. cayennensis</i>	0.00%	0.05%	0.00%	0.07%	0.02%	8	0.15%	1.93%	1.18%	0.83%	0.08%	0.75%	0.12%	124	4.89%
<i>Mi. micropyga</i>	0.00%	0.02%	0.00%	0.00%	0.00%	1	0.02%	0.59%	0.20%	0.16%	0.00%	0.16%	0.47%	40	1.58%
<i>Mi. pilosa</i>	0.00%	0.02%	0.00%	0.02%	0.00%	2	0.04%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Mi. trinidadensis</i>	0.38%	1.59%	0.00%	1.72%	0.16%	211	3.85%	0.95%	1.34%	0.59%	0.28%	2.09%	0.47%	145	5.72%
<i>Mi. venezuelensis</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%	0.00%	0.00%	0.00%	0.04%	0.00%	0.00%	1	0.04%
<i>Ny. trapidoi</i>	0.11%	0.16%	0.00%	0.02%	7.14%	407	7.43%	0.28%	0.00%	0.20%	0.00%	0.04%	0.43%	24	0.95%
<i>Ny. yuilli yuilli</i>	0.00%	0.02%	0.00%	0.00%	1.20%	67	1.22%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Ny. ylephiletor</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%	0.00%	0.08%	0.00%	0.00%	0.00%	0.04%	3	0.12%
<i>Pa. carpenteri</i>	0.00%	0.00%	0.00%	0.00%	0.02%	1	0.02%	0.16%	0.00%	0.00%	0.00%	0.04%	0.00%	5	0.20%
<i>Pa. shannoni</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%	0.12%	0.00%	0.00%	0.00%	0.00%	0.00%	3	0.12%
<i>Pi. colombiana</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%	2.17%	0.08%	0.04%	7.10%	0.36%	0.59%	262	10.34%
<i>Pi. nunetovari</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%	0.16%	0.00%	0.00%	4.10%	0.00%	0.00%	108	4.26%
<i>Pi. ovallesi</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%	0.08%	0.00%	0.00%	0.24%	0.00%	0.00%	8	0.32%
<i>Pi. pia</i>	0.04%	0.00%	0.00%	0.00%	0.02%	3	0.05%	0.28%	0.00%	0.00%	0.00%	0.00%	0.00%	7	0.28%
<i>Pi. spinicrassa</i>	0.00%	0.00%	0.00%	0.02%	0.00%	1	0.02%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Pr. camposi</i>	0.05%	0.78%	0.00%	0.44%	0.15%	78	1.42%	0.12%	0.00%	0.00%	0.04%	0.28%	0.00%	11	0.43%
<i>Ps. carrerai carrerai</i>	0.00%	0.02%	0.00%	0.00%	0.07%	5	0.09%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Ps. carrerai thula</i>	0.02%	0.02%	0.00%	0.00%	0.00%	2	0.04%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Ps. hirsutus hirsutus</i>	0.40%	0.05%	0.00%	0.09%	0.04%	32	0.58%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Ps. nocticolus</i>	0.00%	0.00%	0.00%	0.02%	0.02%	2	0.04%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0	0.00%
<i>Ps. paramensis</i>	1.84%	4.98%	0.00%	6.90%	2.72%	901	16.45%	0.39%	0.28%	0.00%	0.12%	0.00%	0.00%	20	0.79%
<i>Ty. triramula</i>	0.00%	0.00%	0.00%	0.11%	0.11%	6	0.11%	0.04%	0.04%	0.00%	0.39%	0.04%	0.00%	13	0.51%
<b>Total</b>	<b>9.13%</b>	<b>26.91%</b>	<b>0.33%</b>	<b>30.25%</b>	<b>33.22%</b>	<b>5469</b>	<b>100%</b>	<b>24.43%</b>	<b>10.34%</b>	<b>13.14%</b>	<b>19.81%</b>	<b>27.70%</b>	<b>4.18%</b>	<b>2524</b>	<b>100%</b>

analysis of the spatial distribution of the six species, 90 georeferenced points were used. For the construction of the predictive models with MaxEnt, 21 bioclimatic variables which correspond to environmental layers in raster format for the area were used with a resolution of 30 seconds (1 km). Monthly mean precipitation and temperature data were taken from the Worldclim database (worldclim, 2020). The variables used were the following: BIO1, annual average temperature (°C); BIO2, diurnal range of temperature (°C); BIO3, isothermality (°C); BIO4, seasonality of temperature (°C); BIO5, maximum temperature of the hottest period (°C); BIO6, minimum temperature of the coldest period (°C); BIO7, annual range of temperature (°C); BIO8, average temperature in the rainiest quarter (°C); BIO9, average temperature in the driest quarter (°C); BIO10, average temperature in the hottest quarter (°C); BIO11, average temperature in the coldest


quarter (°C); BIO12, annual precipitation (mm); BIO13, precipitation in the rainiest period (mm); BIO14, precipitation in the driest period (mm); BIO15, seasonality of precipitation (%); BIO16, precipitation in the rainiest quarter (mm); BIO17, precipitation in the driest quarter (mm); BIO18, precipitation in the hottest quarter (mm); and BIO19, precipitation in the coldest quarter (mm). Topographic and ecological variables could not be included in this study.

The collected species *Lu. (Hel.) sanguinaria* (Fairchild and Hertig, 1957), *Mg. (Mig.) migonei* (França, 1920), *Pi. (Pif.) spinicrassa* (Morales et al., 1969) and *Pi. (Pif.) moralesi* (Young, 1979) correspond to new records for the department of Antioquia, thus expanding the geographical expansion of these species in the country.

In addition, five species were collected using human



*Lu. (Trl.) gomezi*




*Lu. (Hel.) hartmanni*




*Ps. panamensis*



*Ny. trapidoi*




*Ny. yuilli yuilli*



*Pi. (Pif.) columbiana*

**Figure 2.** Spermathecae of phlebotomine sandfly species of epidemiological importance collected in the area of influence of the Ituango Hydroelectric Project, Antioquia.



**Figure 3.** Potential distribution of phlebotomine sandflies of epidemiological importance in the area of influence of the Ituango Hydroelectric Project, Antioquia.

bait, suggesting that they have anthropophilic habits. These were: *Lu. (Lut) lichyi*; for which this behavior has already been recorded in the Cauca valley (Alexander et al., 2001) and in Venezuela (Cazorla-Perfetti, 2015). *Lu. (Hel.) strictivilla*; a species that according to Young, feeds on humans (Young, 1979). *Mi. (Sau.) trinidadensis*; reported to have a feeding preference for geckos and lizards (Young, 1979), although it has been reported to bite humans in the Colombian Orinoquia (Vivero et al., 2010). *Ev. (Ald.) dubitans* is considered saurophilous (Young and Duncan, 1994), but in Colombia there are reports of it biting humans in the Montes de María (Cortés et al., 2009). Finally, *Lu. (Hel.) cirrita*, recorded in southwestern Colombia in Alto Anchicayá (Valle del Cauca) and reported in human baiting (Barreto et al., 1997).

## DISCUSSION

This research determined the diversity and abundance of phlebotomine sandfly species present in the rural areas of the Cauca River canyon, which are part of the Ituango hydroelectric project. It was also possible to update the inventory of epidemiologically important species as vectors of

leishmaniasis in these municipalities.

In addition, the presence and wide dispersion of species reported in the literature as vectors of *Leishmania* species that cause cutaneous leishmaniasis was verified: *Lu. (Trl.) gomezi*, *Lu. (Hel.) hartmanni*, *Ps. panamensis*, *Ny. trapidoi*, *Ny. yuilli yuilli* and *Pi. (Pif.) columbiana*. It is important to note that this research presents new records of *Lu. (Hel.) sanguinaria*, *Mg. (Mig.) migonei*, *Pi. (Pif.) spinicrassa* and *Pi. (Pif.) moralesi* for the department of Antioquia. However, these species have not been incriminated in the transmission of the parasite to date.

The results show that in both subregions the dominant species present in all sampling points was *Lu. (Trl.) gomezi*, which presented a  $Pi = 50.04$  for the western subregion and a  $Pi = 50.35$  for the northern subregion. This species was collected in extra, peri and intradomiciliary, in protected human bait and in Shannon traps.

The behavior of *Lu. (Trl.) gomezi* is consistent with that reported by other authors, which shows that this species could easily adapt to different environments due to a high degree of adaptability



(Niño and Pérez-Español, 2021). At the same time it is the most widely distributed species in Colombia, being registered in 24 departments (Bejarano, 2006; Bejarano and Estrada, 2016; Ferro et al., 2015). In addition, this species has been recorded in different hydroelectric power plants in the country, since the work carried out by Porter, in 1970 in the Providencia hydroelectric power plant (Porter and DeFoliart, 1981) to later works such as in the Porce II-III hydroelectric power plant (Zuluaga et al., 2012) at Isagen-San Carlos (Parra-Henao and Echavarría, 2005) at the Sogamoso hydroelectric power plant (Gómez-Vargas and Zapata-Úsuga, 2022) and in the La Miel I hydroelectric power plant (Veléz-Bernal et al., 1999). In relation to works carried out in other countries, where there are records of important species, Brazil reports the presence of *Lu. (Trl.) gomezi* in the Usina Luís Eduardo Magalhães hydroelectric power plants (Vilela et al., 2011) and in the Santo Antônio do Jari hydroelectric system (Furtado et al., 2016). It is also important to highlight that in Colombia, *Lu. (Trl.) gomezi* is the most important vector in the transmission of *L. panamensis* and *L. braziliensis* (World Health Organization, 2012) which cause cutaneous leishmaniasis (INS, 2019). Regarding the other species of importance, there are reports of *Lu. (Hel.) hartmanni* and *Ny. trapidoi* in the Toaschi- Piltáon hydroelectric plant in Ecuador, where their abundance implies that *Ny. trapidoi* is responsible for the transmission of leishmaniasis cases in the area (León et al., 2014).

Since 1997, the Cauca river canyon, especially the northern sub-region, has been an area where, before the construction of the Hidroituango project began, some of its municipalities have registered cases of leishmaniasis and is considered endemic by the VTE control and prevention program of the Health Department of Antioquia, which is why it is justified to conduct other studies on this entomofauna in the area.

Some species were collected only in a specific subregion, such as *Pi. (Pif.) columbiana*, which was collected only in the western subregion where tropical dry forests predominate. This species has been associated with *L. mexicana* transmission in an outbreak of cutaneous leishmaniasis in the southwestern part of the country (Cárdenas et al., 1999; Montoya et al., 1999) and is distributed from the Western and Central Cordilleras to the steppe zones of La Guajira, in a wide altitudinal range

from 100 to 2700 m.a.s.l. (Bejarano et al., 2003). *Ny. yuilli yuilli* was only collected in the northern subregion where tropical rainforest predominates. As for its bionomics, it has been recorded in eight departments, namely: Amazonas, Antioquia, Caquetá, Chocó, Guaviare, Meta, Putumayo and Santander (Bejarano, 2006).

When analyzing the bioclimatic variables with the species of epidemiological importance, it can be observed that they are associated with the biological corridor of the Cauca River, where there are points of higher concentration, indicating a greater probability of collecting them. This is related to the areas of the Cauca River canyon that are critical for leishmaniasis, as can be seen in Figure 3, where the area in red represents the places with the greatest presence of these phlebotomine sandflies and which are important in the transmission casuistry.

## CONCLUSIONS

In general, we can conclude that the area of the Cauca River canyon where the Ituango hydroelectric project is being executed is an area where phlebotomine sandflies are present, where there are species that are involved in the transmission of *Leishmania* species that cause cutaneous and mucocutaneous leishmaniasis, which is confirmed by the presence of patients who have registered this disease.

Of the 55 species reported for the department (Bejarano and Estrada, 2016), 39 species of phlebotomine sandflies were found in the study area and at least six species are confirmed vectors of parasite transmission: *Lu. (Trl.) gomezi*, *Lu. (Hel.) hartmanni*, *Ps. panamensis*, *Ny. trapidoi* and *Ny. yuilli yuilli* and *Pi. (Pif.) columbiana*.

Based on the abundance of phlebotomine sandflies of public health interest found in the sampling sites, and the endemic nature of the disease in some of the municipalities, it remains an area of great epidemiological importance for the transmission of *Leishmania* species that cause cutaneous leishmaniasis, with *Lu. (Trl.) gomezi* standing out as a possible main vector, due to its great abundance in the northern and western subregion. However, the vectorial role of this species as well as that of other important species is not clear, nor are the transmission dynamics of each one;

future systematic studies are needed to resolve these concerns and to design appropriate control strategies.

## FINANCING

This study was carried out thanks to Contract CT- I 2012-000030 established between EPM S.A. E.S.P. with the National School of Public Health of the University of Antioquia, between 2012-2020.

## ACKNOWLEDGMENTS

The authors express their sincere thanks to EPM S.A.E.S.P. for funding the study, the National School of Public Health of the University of Antioquia, the Department of Biosciences of the Faculty of Sciences of the National University of Colombia, Medellín, and Dr. Sandra Inés Uribe Soto. Special thanks to the people of the study areas that provided us with accompaniment, security and access to their homes to carry out the entomological samplings. The English version was translated by *Actualidades Biológicas* Journal.

## AUTHORS CONTRIBUTION

Giovani Zapata-Úsuga participated as principal investigator of the project, in the preparation of the manuscript and in the analysis of results. Wilber Gómez-Vargas participated in the collection of field material and in the preparation of the manuscript. Paula Mejía-Salazar Vargas participated in the collection of field material. Boris Zuleta-Ruiz Vargas participated in the collection of field material. Walter Zuluaga-Ramírez Vargas participated in the coordination of the project. All authors approved and reviewed the final version of the manuscript.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

## REFERENCES

- Abbate, J. M., Maia, C., Pereira, A., Arfuso, F., Gaglio, G., Rizzo, M., Caracappa, G., Marino, G., Pollmeier, M., Giannetto, S., & Brianti, E. (2020). Identification of trypanosomatids and blood feeding preferences of phlebotomine sand fly species common in Sicily, Southern Italy. *PLoS ONE*, 15(3), Article e0229536. <https://doi.org/10.1371/journal.pone.0229536>
- Acevedo, M. Á. & Arrivillaga, J. (2008). Eco-Epidemiología de flebovirus (Bunyaviridae, Phlebovirus) transmitidos por flebotomos (Psychodidae, Phlebotominae). *Boletín de Malaria y Salud Ambiental*, 48(1), 3-16. [http://ve.scielo.org/scielo.php?script=sci\\_arttext&pid=S1690-46482008000100001&lng=en&tlng=es](http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S1690-46482008000100001&lng=en&tlng=es).
- Acosta, L. A., Mondragón-Shem, K., Vergara, D., Vélez-Mira, A., Cadena, H., & Carrillo-Bonilla, L. (2013). Ampliación de la distribución de *Lutzomyia longipalpis* (Lutz & Neiva, 1912) (Diptera: Psychodidae) en el departamento de Caldas: potencial aumento del riesgo de leishmaniasis visceral. *Biomédica*, 33(2), 319-325. <https://doi.org/10.7705/biomedica.v33i2.762>
- Akhoundi, M., Kuhls, K., Cannet, A., Votýpka, J., Marty, P., Delaunay, P., & Sereno, D. (2016). A historical overview of the classification, evolution, and dispersion of *Leishmania* parasites and sandflies. *PLoS Neglected Tropical Diseases* 10(3), Article e0004770 <https://doi.org/10.1371/journal.pntd.0004349>
- Alexander, B., Usma, M. C., Cadena, H., Quesada, B.L., Solarte, Y., Roa, W., Montoya, J., Jaramillo, C., & Travi, B. L. (1995). Phlebotomine sandflies associated with a focus of cutaneous leishmaniasis in Valle del Cauca, Colombia. *Medical and Veterinary Entomology*, 9(3), 273-278. <https://doi.org/10.1111/j.1365-2915.1995.tb00133.x>
- Alexander, B., Agudelo, L., Navarro, F., Ruiz, F., Molina, J., Aguilera, G., & Quiñónez, M. (2001). Phlebotomine sandflies and leishmaniasis risks in Colombian coffee plantations under two systems of cultivation. *Medical and Veterinary Entomology*, 15(4), 364-373. <https://doi.org/10.1046/j.0269-283x.2001.00322.x>
- Alvar, J., Vélez, I. D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jamin, J., & den Boer, M. (2012). Leishmaniasis worldwide and global estimates of its incidence. *PLoS ONE*, 7(5), Article e35671. <https://doi.org/10.1371/journal.pone.0035671>
- Barreto, M., Burbano, M. E., & Barreto, P. (1997). Nuevos registros de flebotominos (Diptera: Psychodidae) y triatominos (Hemiptera: Reduviidae) para Risaralda, Cauca y Valle del Cauca, Colombia. *Colombia Médica*, 28(3), 116-122. [https://bibliotecadigital.univalle.edu.co/bitstream/handle/10893/6998/Nuevos\\_registros\\_de.pdf?sequence=1](https://bibliotecadigital.univalle.edu.co/bitstream/handle/10893/6998/Nuevos_registros_de.pdf?sequence=1)
- Bastidas, R. G., Oviedo, M., Vivenes, J. A., & González, A. (2004). Determinación del azúcar preferencial en la dieta de *Lu. evansi* (Nuñeztovar) (Diptera: Psychodidae). *Revista Colombiana de Entomología*, 30(2), 193-196. [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S0120-04882004000200011&lng=en&tlng=es](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0120-04882004000200011&lng=en&tlng=es).
- Bejarano, E. E., Patricia, D., & Vélez, I. D. (2006).

- Estudio de los flebotómicos (Diptera: Psychodidae) antropofílicos de la Serranía de La Macarena, Colombia. *Revista colombiana de entomología*, 32(2), 176–178. [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S0120-04882006000200012&lng=en&tlng=es](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0120-04882006000200012&lng=en&tlng=es).
- Bejarano, E. E., Uribe, S., Rojas, W., & Vélez, I. D. (2002). Phlebotomine sand flies (Diptera: Psychodidae) associated with the appearance of urban leishmaniasis in the City of Sincelejo, Colombia. *Memorias do Instituto Oswaldo Cruz*, 97(5), 645–647. <https://doi.org/10.1590/S0074-02762002000500010>
- Bejarano, E. E., Sierra, D., & Vélez, I. D. (2003). Novedades en la distribución geográfica del grupo verrucarum (Diptera: Psychodidae) en Colombia. *Biomédica*, 23(3), 341–350. <https://doi.org/10.7705/biomedica.v23i3.1228>
- Bejarano, E. E. (2006). Lista actualizada de los psicódidos (Diptera: Psychodidae) de Colombia. *Folia Entomológica Mexicana*, 45(1), 47–56. <https://www.redalyc.org/pdf/424/42445106.pdf>
- Bejarano, E. E., & Estrada, L. G. (2016). Family psychodidae. *Zootaxa*, 4122(1), 187–238. <https://doi.org/10.11646/zootaxa.4122.1.20>
- Cabrera, O. L., Mosquera, L., Santamaría, E., & Ferro, C. (2009). Flebotomos (Diptera: psychodidae) del departamento de Guaviare, Colombia, con nuevos registros para el país. *Biomedica*, 29(1), 73. <https://doi.org/10.7705/biomedica.v29i1.43>
- Cameron, M. M., Davies, C. R., Monje, J., Villaseca, P., Ogasaku, E., & Llanos-Cuentas, A. (1994). Comparative activity of phlebotomine sandflies (Diptera: Psychodidae) in different crops in the Peruvian Andes. *Bulletin of Entomological Research*, 84(4), 461–467. <https://doi.org/10.1017/S0007485300032685>
- Cárdenas, R., Romo, G. M., Santamaría, E., Bello, F., & Ferro, C. (1999). *Lutzomyia longiflora* (Diptera: Psychodidae) posible vector en el foco de leishmaniasis cutánea del municipio de Planadas, zona cafetera del Tolima. *Biomedica*, 19(3), 239–244. <https://doi.org/10.7705/biomedica.v19i3.1028>
- Cazorla-Perfetti, D. (2015). Lista comentada de los flebotominos (Diptera: Psychodidae, Phlebotominae) citados para Venezuela. *Saber*, 27(2), 178–231. [http://ve.scielo.org/scielo.php?script=sci\\_arttext&pid=S1315-01622015000200003&lng=en&tlng=es](http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S1315-01622015000200003&lng=en&tlng=es).
- Cecílio, P., Cordeiro-da-Silva, A., & Oliveira, F. (2022). Sand flies: Basic information on the vectors of leishmaniasis and their interactions with *Leishmania* parasites. *Communications Biology*, 5(1), 305–316. <https://doi.org/10.1038/s42003-022-03240-z>
- Contreras, M. A., Vivero, R. J., Bejarano, E. E., Carrillo, L. M., & Vélez, I. D. (2012). Nuevos registros de flebotómicos (Diptera : Psychodidae) en el área de influencia del río Amoyá en Chaparral, Tolima. *Biomédica*, 32(2), 263–268. [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S0120-41572012000200014](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0120-41572012000200014)
- Corredor, A., Gallego, J. F., Tesh, R. B., Morales, A., De Carrasquilla, C. F., Young, D. G., Kreutzer, R. D., Boshell, J., Palau, M. T., Cáceres, E., & Pelaez, D. (1989). Epidermiology of visceral leishmaniasis in Colombia. *American Journal of Tropical Medicine and Hygiene*, 40(5), 480–486. <https://doi.org/10.4269/ajtmh.1989.40.480>
- Corredor, A., Kreutzer, R., Tesh, R., Boshell, J., Palau, M., Cáceres, E., Duque, S., Pelaez, D., Rodríguez, G., Nichols, S., Hernández, C., Morales, A., Young, D., & Ferro, C. (1990). Distribution and etiology of leishmaniasis in Colombia. *The American journal of tropical medicine and hygiene*, 42(3), 206–214. <http://dx.doi.org/10.4269/ajtmh.1990.42.206>
- Cortés, A. L., Pérez, D. A., & Bejarano, E. E. (2009). Flebotómicos (Diptera: Psychodidae) antropofílicos de importancia en salud pública en Los Montes de María, Colombia. *Revista Cubana de Medicina Tropical*, 61(3), 220–225. [http://scielo.sld.cu/scielo.php?script=sci\\_arttext&pid=S0375-07602009000300003&lng=es&tlng=es](http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0375-07602009000300003&lng=es&tlng=es)
- Dirección Seccional de Salud de Antioquia. (1998). *Leishmaniasis incidencias por grupos de edad 1995-1997: Servicio Seccional de Salud de Antioquia*, Boletín Informativo de Antioquia.
- Dirección Seccional de Salud de Antioquia & Laboratorio Departamental de Salud Pública. (1998). *Distribución de Leishmaniasis para el departamento de Antioquia* [Data set]. SIVIGILA.
- Dirección Seccional de Salud de Antioquia. (2011). *2002-2010. Información de interés- enfermedades transmitidas por vectores* [Data set]. SIVIGILA.
- Dirección Seccional de Salud de Antioquia. (2022). *Eventos de Salud Pública por municipios. 2007-2022*. Recuperado de: <https://dssa.gov.co/index.php/vigilancia-en-salud-publica>
- Fairchild, G. B., & Hertig, M. (1957). Notes on the *Phlebotomus* of Panama XIII. The Vexator Group, with Descriptions of New Species from Panama and California. *Annals of the Entomological Society of America*, 50(4), 325–334. <https://doi.org/10.1093/aesa/50.4.325>
- Ferro, C., López, M., Fuya, P., Lugo, L., Cordovez, J. M., & González, C. (2015). Spatial distribution of sand fly vectors and eco-epidemiology of cutaneous leishmaniasis transmission in Colombia. *PLoS ONE*, 10(10), Article e0139391. <https://doi.org/10.1371/journal.pone.0139391>
- França, C. (1920) Observations sur le genre *Phlebotomus*. II. -*Phlebotomus* du Nouveau Monde (Phlébotomes du Brasil et du Paraguay). *Bulletin de la Société Portugaise*

- des Sciences Naturelles*, 8, 215–236. <https://biostor.org/reference/275790>
- Furtado, N. V. R., Galardo, A. K. R., Galardo, C. D., Firmino, V. C., & Dos Santos, T. V. (2016). Phlebotomines (Diptera: Psychodidae) in a Hydroelectric System Affected Area from Northern Amazonian Brazil: further insights into the effects of environmental changes on vector ecology. *Journal of Tropical Medicine*, 1–12. <https://doi.org/10.1155/2016/9819723>
- Galati, E. A. B., Galvis-Ovallos, F., Lawyer, P., Léger, N., & Depaquit, J. (2017). An illustrated guide for characters and terminology used in descriptions of Phlebotominae (Diptera, Psychodidae). Guide illustré des caractères et de la terminologie utilisés dans les descriptions de Phlebotominae (Diptera, Psychodidae). *Parasite (Paris, France)*, 24, 26. <https://doi.org/10.1051/parasite/2017027>
- Galati, E. A. B. (2018). *Morfología e terminología de Phlebotominae (Diptera: Psychodidae). Classificação e identificação de táxons das Américas (Vol. 1)* [archivo PDF]. Recuperado de: <http://www.fsp.usp.br/~egalati/>
- García-Leal, J., Carrero-Sarmiento, D., & Hoyos- López, R. (2022). Diversidad del género *Lutzomyia* (Diptera: Psychodidae) en municipios del departamento de Córdoba – Colombia. *Acta Biológica Colombiana*, 27(3), 394 – 402. <https://doi.org/10.15446/abc.v27n3.90684>
- Goenaga-Mafud, L. C., Eyes-Escalante, M., & Florez-Arrieta, F. (2020). Primera evaluación sobre la fauna de flebotomos (Psychodidae: Phlebotominae) del departamento del Atlántico de Colombia. *Acta Biológica Colombiana*, 25(3), 284–292. <https://doi.org/10.15446/abc.v25n3.78771>
- Gómez-Vargas, W., & Zapata-Úsuga, G. (2022). Especies de *Lutzomyia* (Diptera: Psychodidae, Phlebotominae) en el área de influencia de la Central Hidroeléctrica Sogamoso (Santander, Colombia). *Boletín Científico Centro de Museos Museo de Historia Natural*, 26(1), 121–133. <https://doi.org/10.17151/bccm.2022.26.1.9>
- González, E. (2019, Junio 14). *Estudios entomológicos en el foco de leishmaniosis del suroeste de la Comunidad de Madrid* [Tesis Doctoral]. Universidad Complutense de Madrid, Madrid. <https://hdl.handle.net/20.500.14352/17155>
- Gradoni, L. (2018). A brief introduction to leishmaniasis epidemiology. *The Leishmaniasis: Old Neglected Tropical Diseases* (pp. 245). Springer International Publishing AG 2018. [https://doi.org/10.1007/978-3-319-72386-0\\_1](https://doi.org/10.1007/978-3-319-72386-0_1)
- Grimaldi, G., Tesh, R. B., & McMahon-Pratt, D. (1989). A review of the geographic distribution and epidemiology of leishmaniasis in the New world. *American Journal of Tropical Medicine and Hygiene*, 41(6), 687–725. <https://doi.org/10.4269/ajtmh.1989.41.687>
- Instituto Nacional de Salud. (2019). *Informe de vigilancia entomológica de Leishmaniasis. Colombia 2018*. [archivo PDF]. Recuperado de <https://www.ins.gov.co/buscador-eventos/Informacin%20de%20laboratorio/Informe-vigilancia-entomologica-Leishmaniasis-Colombia-2018.pdf>
- Lainson, R., & Shaw, J. J. (1973). Las Leishmanias y la leishmaniasis del Nuevo Mundo con particular referencia al Brasil. *Boletín de la Oficina Sanitaria Panamericana*, VII N° 4, 1–19. <https://iris.paho.org/bitstream/handle/10665.2/10793/v76n2p93.pdf?sequence=1>
- León, R., Ortega, L., Gualapuro, M., Morales, F., Rojas, M. I., & Espinel, M. (2014). Identificación de Dípteros Nematóceros de interés médico en zonas aledañas a la construcción de la Represa Hidroeléctrica Toachi-Pilatón. *ACI Avances en Ciencias e Ingenierías*, 6(2), 25–31. <https://doi.org/10.18272/aci.v6i2.176>
- Maroli, M., Feliciangeli, M. D., Bichaud, L., Charrel, R. N., & Gradoni, L. (2013). Phlebotomine sandflies and the spreading of leishmaniasis and other diseases of public health concern. *Medical and Veterinary Entomology*, 27(2), 123–147. <https://doi.org/10.1111/j.1365-2915.2012.01034.x>
- Martínez, D. C., Ávila, J. L., & Molano, F. (2018). Actividad nocturna de *Nyssomyia yuilli* (Young & Porter, 1972) y *Nyssomyia trapidoi* (Farchild & Hertig, 1952) (Diptera: Psychodidae: Phlebotominae) en el municipio de Otanche (Boyacá, Colombia), zona endémica de leishmaniasis cutánea. *Salud UIS*, 50(2), 106–114. <https://doi.org/10.18273/revsal.v50n2-2018002>
- Méndez-Cardona, S. A. (2021, Agosto). *Descripción del cuarto estadio larval de *Pintomyia longiflocosa* e implicaciones taxonómicas en el subgénero *Pifanomyia** [Tesis Doctoral]. Universidad de los Andes, Bogotá. <https://repositorio.uniandes.edu.co/bitstream/handle/1992/53899/24942.pdf?sequence=1>
- Ministerio de Protección Social [Minsalud]. (2010). *Gestión para la vigilancia entomológica y control de la transmisión de Leishmaniasis: Guía de Vigilancia Entomológica y control de Leishmaniasis*. [archivo PDF]. Recuperado de [http://simudatsalud-risaralda.co/normatividad\\_inv7/Entomologica%20Leishmaniasis.pdf](http://simudatsalud-risaralda.co/normatividad_inv7/Entomologica%20Leishmaniasis.pdf)
- Montoya-Lerma, J., Cadena, H., Segura, I., & Travi, B. L. (1999). Association of *Lutzomyia columbiana* (Diptera: Psychodidae) with a leishmaniasis focus in Colombia due to species of the *Leishmania mexicana* complex. *Memórias do Instituto Oswaldo Cruz*, 94(3), 277–283. <https://doi.org/10.1590/S0074-02761999000300001>
- Morales, A., E. Osorno-Mesa, F. Osorno, and P. Muñoz Hoyos. (1969). Phlebotominae de Colombia (Diptera, Psychodidae). V. Descripción de una nueva especie de *Lutzomyia*. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 13(51), 383–390. <https://raccefyn.co/index.php/raccefyn/issue/>

- view/93/237
- Morales, A., Bello, F., & Cárdenas, E. (2005). Establecimiento, mantenimiento y productividad de una colonia de laboratorio de *Lutzomyia spinicrassa* Morales, Osorno-Mesa, Osorno y Hoyos (Diptera: Psychodidae) en Colombia. *Revista Ciencias de la Salud*, 3(2), 129–135. <http://www.scielo.org.co/pdf/recis/v3n2/v3n2a3.pdf>
- Niño, L., & Pérez-Español, S. (2021). Distribución geográfica y factores ambientales asociados a flebotomíneos del género *Lutzomyia* França, 1924 (Diptera: Psychodidae) en Cundinamarca, Colombia. *Revista Chilena De Entomología*, 47(2), 275–303. <https://doi.org/10.35249/rche.47.2.21.15>
- Organización Mundial de la Salud. (2012). *Control de las Leishmaniasis: informe de una reunión del Comité de Expertos de la OMS sobre el Control de las Leishmaniasis, Ginebra, 22 a 26 de marzo de 2010*. Recuperado de [https://apps.who.int/iris/bitstream/handle/10665/82766/WHO\\_TRS\\_949\\_spa.pdf;sequence=1](https://apps.who.int/iris/bitstream/handle/10665/82766/WHO_TRS_949_spa.pdf;sequence=1)
- Organización Panamericana de la Salud. (2023). *Leishmaniasis*. Paho.org. Recuperado de; <https://www.paho.org/es/temas/leishmaniasis>
- Organización Panamericana de la Salud. (2019). *Manual de procedimientos para vigilancia y control de las leishmaniasis en las Américas*. Washington, D.C. Recuperado de <https://doi.org/https://doi.org/10.37774/9789275320631>
- Pardo, R., Ferro, C., Lozano, G., Lozano, C., Cabrera, O., & Davies, C. (1999). Flebotomos vectores de leishmaniasis cutánea y sus determinantes ecológicos en la zona cafetera del Departamento del Huila. [Memorias XXVI Congreso de la Sociedad Colombiana de Entomología]. *Socolen*, 147–163. <http://hdl.handle.net/20.500.12324/34421>
- Parra-Henao, G. J., & Echavarría, J. C. (2005). Experiencias en el control de un foco de leishmaniasis cutánea en San Carlos, Antioquia. *CES Medicina*, 19(1), 31–36. <https://revistas.ces.edu.co/index.php/medicina/article/view/193>
- Porter, C. H., & DeFoliart, G. R. (1981). The man biting activity of phlebotomine sand flies in a tropical wet forest environment in Anori central providencia Colombia. *Archivos De Zoología*, 30(2), 81–158. <https://doi.org/10.11606/issn.2176-7793.v30i2p81-158>
- Posada-López, L., Vélez-Mira, A., Acosta, L., Cadena, H., Agudelo, D., & Vélez, I. D. (2014). Descripción de un foco endémico de leishmaniasis cutánea en Puerto Valdivia, Antioquia, Colombia. *Revista CES Salud Pública*, 5(1), 3–10. [https://revistas.ces.edu.co/index.php/ces\\_salud\\_publica/article/view/3029](https://revistas.ces.edu.co/index.php/ces_salud_publica/article/view/3029)
- Sales, K. G. D. S., Costa, P. L., De Morais, R. C. S., Otranto, D., Brandão-Filho, S. P., Cavalcanti, M. D. P., & Dantas-Torres, F. (2015). Identification of phlebotomine sand fly blood meals by real-time PCR. *Parasites and Vectors*, 8(1), 1–6. <https://doi.org/10.1186/s13071-015-0840-3>
- Sant’Anna, M. R. V., Jones, N. G., Hindley, J. A., Mendes-Sousa, A. F., Dillon, R. J., Cavalcante, R. R., Alexander, B., & Bates, P. A. (2008). Blood meal identification and parasite detection in laboratory-fed and field-captured *Lutzomyia longipalpis* by PCR using FTA databasing paper. *Acta Tropica*, 107(3), 230–237. <https://doi.org/10.1016/j.actatropica.2008.06.003>
- Santamaría, E., Ponce, N., Zipa, Y., & Ferro, C. (2006). Presencia en el peridomicilio de vectores infectados con *Leishmania (Viannia) panamensis* en dos focos endémicos en el occidente de Boyacá, piedemonte del valle del Magdalena medio, Colombia. *Biomédica* 26 (sup1), 82–94. [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S0120-41572006000500011](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0120-41572006000500011)
- Tanure, A., Peixoto, J. C., Afonso, M. M. D. S., Duarte, R., Pinheiro, A. D. C., Coelho, S. V. B., & Barata, R. A. (2015). Identification of Sand flies (Diptera: Psychodidae: Phlebotominae) blood meals in an endemic leishmaniasis area in Brazil. *Revista do Instituto de Medicina Tropical de São Paulo*, 57(4), 321–324. <https://doi.org/10.1590/s0036-46652015000400008>
- Travi, B. L., Montoya, J., Solarte, Y., Lozano, L., & Jaramillo, C. (1988). Leishmaniasis in Colombia. I. Studies on the phlebotomine fauna associated with endemic foci in the Pacific Coast region. *American Journal of Tropical Medicine and Hygiene*, 39(3), 261–266. <https://doi.org/10.4269/ajtmh.1988.39.261>
- Travi, B. L., Tabares, C. J., Cadena, H., Ferro, C., & Osorio, Y. (2001). Canine visceral leishmaniasis in Colombia: Relationship between clinical and parasitologic status and infectivity for sand flies. *American Journal of Tropical Medicine and Hygiene*, 64(3), 119–124. <https://doi.org/10.4269/ajtmh.2001.64.119>
- Urango, H. J., & Hoyos-López, R. (2022). Flebotomíneos (Psychodidae: Phlebotominae) en la zona periurbana de Montería (Córdoba-Colombia). *Acta Biológica Colombiana*, 27(3), 377–385- <https://doi.org/10.15446/abc.v27n3.92751>
- Vásquez-Trujillo, Adolfo, Santamaría-Herreño, Erika, González-Reina, Angélica E, Buitrago-Álvarez, Luz S, Góngora-Orjuela, Agustín, & Cabrera-Quintero, Olga L. (2008). *Lutzomyia antunesi*, Probable Vector de Leishmaniasis Cutánea en el Área Rural de Villavicencio. *Revista de Salud Pública*, 10(4), 625–632., [http://www.scielo.org.co/scielo.php?script=sci\\_arttext&pid=S0124-00642008000400012&lng=en&tlng=es](http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0124-00642008000400012&lng=en&tlng=es)
- Vélaz-Bernal, I. D., Agudelo-Urbe, L. A., Cuartas-Hernández, S. E., Arango, M., & Díaz, M. R. (1999). Programa de estudio y control de enfermedades tropicales-pecet

- determinación del riesgo de infección por *Leishmania* spp. en la zona de influencia del proyecto hidroeléctrico miel I en el departamento de Caldas. *Rev. ECM*, 4, 45–59. <https://pesquisa.bvsalud.org/portal/resource/pt/lil-385731>
- Vergara, D., Carrillo, L., Bejarano, E., & Vélez, I. (2008). Primer informe de *Lutzomyia yuilli* (Young & Porter, 1972) y *Lutzomyia triramula* (Fairchild & Hertig 1952) (Diptera: Psychodidae) en el departamento de Caldas, Colombia. *Biota Neotropica*, 8(3), 251-253 <https://www.scielo.br/j/bn/a/MDLrNkcZBwZN3c8T-BfGZkXM/?lang=es&format=pdf>
- Vilela, M. L., Azevedo, C. G., Carvalho, B. M., & Rangel, E. F. (2011). Phlebotomine fauna (diptera: Psychodidae) and putative vectors of leishmaniasis in impacted area by Hydroelectric Plant, State of Tocantins, Brazil. *PLoS ONE*, 6(12), Article e27721. <https://doi.org/10.1371/journal.pone.0027721>
- Vivero, R. J., Bejarano, E. E., Castro, M., Vélez, A., Pérez, J. E., Pérez-Doria, A., Vélez, I. D., & Carrillo, L. M. (2010). Trece registros nuevos de *Lutzomyia* (Diptera: Psychodidae) para el departamento de Vichada, Orinoquia Colombiana. *Biota Neotropica*, 10(2), 401–404. <https://doi.org/10.1590/s1676-06032010000200037>
- Vivero, R. J., Torres-Gutiérrez, C., Bejarano, E. E., Cadena-Peña, H., Estrada, L. G., Flórez, F., Ortega, E., Aparicio, Y., & Muskuz, C. E. (2015). Study on natural breeding sites of sand flies (Diptera: Phlebotominae) in areas of *Leishmania* transmission in Colombia. *Parasites and Vectors*, 8(1), 1-14- <https://doi.org/10.1186/s13071-015-0711-y>
- Vivero, R. J., Quintero, L. S., Peña, H. C., Alvar, J., Mira, A. V., Atencia, M., Tovar, C., Vélez, I. D (2016, Abril ). *Listado taxonómico y distribución espacial de especies de importancia médica del género Lutzomyia presentes en los municipios de Tierra Alta.*, [Postér presentado en la Primera Reunión Colombiana de *Leishmaniasis* y enfermedad de *Chagas*]. XV Simposio PECET: Avances en la investigación de enfermedades tropicales, Medellín, Colombia. <https://issuu.com/comunicaciones.pecet/docs/memoriasreunionsimposiopecet>
- Vivero, R. J., Contreras, M. A., & Suaza, J. D. (2017). Especies de flebotomíneos (Diptera : Psychodidae ) recolectados en reservas naturales de las regiones del Darién y del Pacífico en Colombia. *Biomédica*, 37(Supl.2), 215–223. <https://doi.org/10.7705/biomedica.v37i0.3382>
- Warburg, A., Montoya-Lerma, J., Jaramillo, C., Cruz-Ruiz, A., & Ostrovska, K. (1991). Leishmaniasis vector potential of *Lutzomyia* spp. in Colombian coffee plantations. *Medical and Veterinary Entomology*, 5(1), 9-16. <https://doi.org/10.1111/j.1365-2915.1991.tb00514.x>
- WorldClim.org. (2020). <https://www.worldclim.org/>
- Young, D. G. (1977). *A biosystematic review of the bloodsucking psychodid flies of Colombia (diptera: Phlebotominae and Sycoracinae)* [Tesis Doctoral]. University of Florida, Florida. <https://doi.org/10.5962/bhl.title.42420>.
- Young, D. G. (1979). A review of the bloodsucking psychodid flies of Colombia (diptera: Phlebotominae and Sycoracinae. *Bulletin 806 (Technical) of the Agricultural Experiment Stations, University of Florida*, pp. 1–266. <https://apps.dtic.mil/sti/trecms/pdf/ADA080842.pdf>
- Young, D. G., & Duncan, M. A. (1994). Guide to the identification and geographic distribution of *Lutzomyia* sand flies in Mexico, the West Indies, Central and South America (Diptera: Psychodidae). *Memoirs of the American Entomologic Institute* (pp. 881). Gainesville, Florida, USA: Associated Publishers. <https://apps.dtic.mil/sti/tr/pdf/ADA285737.pdf>
- Zuluaga, W. A., López, Y., Ríos-Osorio, L. A., Salazar, L., González, M. C., Ríos, C. M., Wolff, M., & Escobar, J. P. (2012). Vigilancia entomológica de insectos de importancia en salud pública durante la construcción de los proyectos hidroeléctricos Porce II y Porce III, Antioquia, Colombia, 1990-2009. *Biomédica*, 32, 321-332. <https://doi.org/10.7705/biomedica.v32i3.668>