

Food and nutrition surveillance system in municipalities of paraíba: data reliability, coverage and management's perception

Sistema de Vigilancia Alimentaria y Nutricional en municipios de Paraíba: confiabilidad, cobertura y percepción de gestores

Sistema de Vigilância Alimentar e Nutricional em municípios da Paraíba: confiabilidade, cobertura e percepção de gestores

Dixis Figueroa Pedraza¹

¹ PhD in Nutrition. Department of Nursing and Postgraduate. Program in Public Health, State University of Paraíba, Campina Grande, Paraíba, Brazil. Correo: dixisfigueroa@gmail.com

Recibido: 31/05/2017. Aprobado: 05/09/2017. Publicado: 10/09/2017

Figueroa-Pedraza D. Food and nutrition surveillance system in municipalities of paraíba: data reliability, coverage and management's perception. Rev. Fac. Nac. Salud Pública, 2017; 35(3): 313-321. DOI: 10.17533/udea.rfnsp.v35n3a02

Abstract

Objectives: To evaluate the reliability of anthropometric data of under-five children based on the SISVAN-Web, describing the coverage of the System and aspects of its implementation.

Methods: Cross-sectional study developed in two cities of Paraíba. The sample consisted of 18 family health teams and 324 children, randomly selected. Anthropometric measures (weight and height) were collected (1) by the researcher using standard equipment (R); (2) by the researcher using service equipment (RS) and; (3) during routine service (S). Municipal managers answered a questionnaire about the System.

Results: Among the 319 children observed, 10.28% were overweight/age, 14.02% were overweight and 7.16% were stunted. Coverage was 2.8% in one municipality and 3.3% in

the other, being higher for children and smaller for the elderly. The agreement between S and R were weaker than those with RS, especially for height/age (Kappa=0.38) and BMI/age (Kappa=0.39). The measurements obtained in S overestimated overweight and stunting. Managers mentioned that the information of the System is restricted to the health sector.

Conclusions: The study pointed to major shortcomings related to SISVAN-Web. The system's coverage and reliability of anthropometric measurements were both low. The System is not used to its full potential.

-----**Keywords:** Nutritional Surveillance. Information Systems. Nutritional status. Anthropometry. Nutrition Programs.

Resumen

Objetivos: Con foco en el SISVAN-Web, evaluar la confiabilidad de datos antropométricos de niños menores de cinco años, describir la cobertura del sistema y conocer aspectos de la implantación. **Metodología:** Estudio transversal realizado en dos municipios de Paraíba. La muestra se constituyó por 18 equipos de salud de la familia y 324 niños. Las medidas antropométricas (peso y estatura) fueron tomadas (1) por el investigador que utilizando equipamiento patrón (P); (2) por el investigador usando equipamientos del servicio (PS) y; (3) en la rutina del servicio (S). Los gestores municipales responden a un cuestionario sobre el Sistema. **Resultados:** De los 319 niños observados, 10,28% estaban con exceso de peso/edad, 14,02% con exceso de peso y 7,16% con déficit de estatura. La cobertura fue de 2,8% en un municipio y 3,3%

en el otro, siendo mayor para niños y menor en ancianos. Las concordancias de S con P fueron más débiles que las de PS, principalmente para estatura/edad ($Kappa = 0,38$) y IMC/edad ($Kappa = 0,39$). Las medidas obtenidas en S superestimaron el exceso de peso y el déficit de estatura. Los gestores refirieron que las informaciones del Sistema son restrictas al sector salud.

Conclusiones: Se apuntan importantes deficiencias relacionadas con el SISVAN-Web que incluyen baja confiabilidad de las medidas antropométricas y de la cobertura. El Sistema no es utilizado en su potencial.

-----**Palabras-clave:** Vigilancia nutricional. Sistemas de información. Estado nutricional. Antropometría. Atención Primaria de Salud. Programas de Nutrición.

Resumo

Objetivos: Com foco no SISVAN-Web, avaliar a confiabilidade de dados antropométricos de crianças menores de cinco anos, descrever a cobertura do Sistema e conhecer aspectos da implantação. **Métodos:** Estudo transversal realizado em dois municípios da Paraíba. A amostra foi constituída por 18 equipes de saúde da família e 324 crianças. As medidas antropométricas (peso e estatura) foram aferidas (1) pelo pesquisador utilizando equipamento padrão (P); (2) pesquisador usando equipamentos do serviço (PS) e; (3) na rotina do serviço (S). Os gestores municipais responderam um questionário sobre o Sistema. **Resultados:** Das 319 crianças observadas, 10,28% estavam com excesso de peso/idade, 14,02% com excesso de peso e 7,16% com déficit de estatura. A cobertura foi de

2,8% em um município e 3,3% no outro, sendo maior para as crianças e menor em idosos. As concordâncias de S com P foram mais fracas do que as de PS, sobretudo para estatura/idade ($Kappa=0,38$) e IMC/idade ($Kappa=0,39$). As medidas obtidas em S superestimaram o excesso de peso e o déficit de estatura. Os gestores referiram que as informações do Sistema ficam restritas ao setor saúde. **Conclusões:** Apontaram-se importantes deficiências relacionadas ao SISVAN-Web que incluem a baixa confiabilidade das medidas antropométricas e pouca cobertura. O Sistema não é utilizado no seu potencial.

-----**Palavras-chave:** Vigilância nutricional. Sistemas de informação. Estado nutricional. Antropometria. Atenção Primária à Saúde. Programas de Nutrição.

Introduction

The Brazilian population has experienced the process of nutritional transition in the last decades, implying the need for transforming the organization of health practices [1, 2]. In this context, two extremes share the same scenario: the vertiginous increase of overweight and the persistence of high frequencies of specific nutritional deficiencies [2].

To address this nutritional paradox, the National Food and Nutrition Policy (NFNP) proposes Food and Nutrition Surveillance (FNS) among its guidelines as a central element in the monitoring of the food and nutritional situation and as a support to political decisions aiming to improve health [3]. The FNS consists in the “continuous description and prediction of trends in food and nutrition conditions of the population and its determinants” [3]. It includes different strategies such as population surveys, nutrition calls, research and scientific production funding [1, 2, 4, 5].

In order to overcome limitations related to the high cost and poor representativeness of population surveys and nutritional calls, which limit health and nutrition policy subsidies, the Food and Nutrition Surveillance System (SISVAN) was established as the basic instrument of the FNS in the decade 1990s. The SISVAN represents a system of data collection, processing and continuous analysis that allows to know the nutritional situation of the population served by the Unified Health System (SUS) [6, 7]. With the creation of the Family Grant Program in 2004, the SISVAN became the main provider of information on the health of beneficiary families [2].

In an attempt to overcome some limitations of the System and integrated with the new technological times, the implementation of SISVAN-Web was suggested in 2008 as a computerized system of Primary Health Care (PHC) replacing the SISVAN Municipal Module [8, 9, 10]. This new system allows the registration of food consumption and anthropometric markers (based on

current references) through an interface that facilitates its use by users [11].

Recent literature review pointed out the lack of studies on SISVAN in the scope of the Family Health Strategy (FHS) [12]. Furthermore, few studies have paid attention to the quality of the information collected to feed the system [13]. However, assessing the SISVAN is important for guiding the NFNP and the subjects involved with it (public managers, health professionals and target population) [9].

Focusing on the SISVAN-Web, this study aimed to evaluate the reliability of anthropometric data of under-five children, describe the coverage of the System and know aspects of the implementation from the perception of managers.

Methodology

Research scenario

The study was conducted in the context of FHS in two municipalities in the state of Paraíba, Brazil, selected for convenience. The choice of municipalities considered similarities of geographic position (location in the metropolitan area of the state capital with access to the available network of services), degree of urbanization (almost 100%), demographic-social indicators (average human development index - 0.748 in a municipality 1 and 0.649 in the other one) and traditional organization of PHC (almost 100% of the population covered by the FHS). The municipalities have a population of 4,596 and 7,862 under-five children, with 19 FHS teams and the other with 28 teams, respectively.

Sampling plan

The sampling plan aimed at the inclusion of 314 randomly selected under-five children (157 from each municipality) from 18 FHS teams (nine from each municipality that corresponded to about half of the teams of a municipality and 1/3 of those of the other municipality). This calculation was established in the larger research from which the results of the present study are part, which evaluated the implementation of feeding and nutrition actions in the FHS and their relation with the quality of care. The SISVAN-Web analysis was one of the objectives. A study developed in the municipality of Montes Claros (MG) [14] indicated a higher proportion (58.1%) of negative evaluation of health services provided by FHS teams formed by physicians and nurses without residence in Family Health compared to the proportion (41.9%) of the teams composed by professionals with this training. Based on this, it was necessary to observe 157 children in each of

the municipalities of Paraíba, considering the expected proportion of the negative evaluation of the health services by the users of 41.9% in the municipality with a better human resource situation of health professionals and 58.1% in the other, with 95% of confidence level, 80% of power, 10% of increase to compensate for possible losses and design effect of 1.2.

Data collection

Data collection was carried out in health facilities between July and December 2014. The field team was formed by professionals and students of the health area. The quality control of the study included: training and standardization of interviewers, use of the Instruction Manual, pilot study and supervision. In order to train the anthropometrists, the recommendations for PHC of the Ministry of Health were adopted [15]. Data collection includes the measurement of the children's weight and height, consultation with the SISVAN-Web to verify the individuals monitored by the System, the quantification of beneficiaries of private health plans and of those who use the public health service, and the application of a questionnaire to the managers responsible for the System.

Anthropometric data were obtained from children scheduled for childcare consultations on the day of data collection, on a typical work day. It was established that each child included in the study would be measured in three situations:

- by health professionals with the use of the available equipment in the health units, respecting their work routine;
- by the anthropometrists of the research using their own equipment, considered of adequate reliability (standard gold measure);
- by the anthropometrists of the research using the equipment of the health units.

The measures obtained by health professionals were collected directly from the medical record or from the Child Health Handbook, after the consultation. At the time, the child's birth date was also noted, and later used to calculate age in the date of the interview.

The anthropometric evaluation by the anthropometrists of the research was performed after the consultation, in order to avoid the observation of the procedures used and interfere with the work of health professionals. All measures were obtained by a single pair of anthropometrists, one of which would be in the auxiliary function, following training standards. Children were measured and weighed twice, with the researchers' own equipment and with those available at health facilities. The average of the two measurements was used in the registration in a specific form for this purpose. Children under 24 months of age had length measured in dorsal decubitus. Children aged 25-60

months had the height measured in orthostatic position. Height corresponds to the set of measures of length and height. All children were weighed in light clothing. In children younger than 24 months of age, weight was obtained by the difference between the weight of the mother with the child in her lap and the value of the mother's weight.

Research equipment were: child anthropometer made of wood with amplitude of 130 cm and subdivisions of 0.1 cm, stadiometer (WCS®) with amplitude of 200 cm and subdivisions of 0.1 cm and electronic platform-type scale with capacity for 150 kg and graduation in 100g (Tanita UM-080®). The equipment of the health units were used in the same conditions by assistance professionals and research anthropometrists.

For the three study situations, the Z-scores of Weight/Age (W/A), Body Mass Index/Age (BMI/A) and Height/Age (H/A) were calculated using the *Software* WHO Anthro 2009, considering the population of the *Multicentre Growth Reference Study* as reference [16]. The categories of nutritional diagnosis of overweight for age ($W/A \geq 2$ Z-score), overweight ($BMI/A \geq 2$ Z-score) and stunting ($H/A < -2$ Z-score) were considered [15].

The municipal coverage of SISVAN-Web for the total population and per life stage (children aged 0 to 4 years, children aged 5 to 10 years, adolescents, adults and the elderly) was calculated based on the totals of individuals monitored, obtained directly in the System, and the number of users using the public health service, estimated by the difference between the number of inhabitants of the municipality and the number of beneficiaries of private health plans. The 2010 demographic census [17] was used as data source of the number of inhabitants in the municipality. Also related to 2010, information provided by the National Agency for Supplementary Health Services made it possible to identify the total number of individuals by age group and sex linked to private health plans [18].

The total coverage was defined as the percentage of individuals, from all life stages, monitored by SISVAN-Web in the health units in 2010. Total coverage was estimated by dividing the number of records of nutritional status of individuals followed in 2010 by the approximated population that used public health services. The same calculation was performed to obtain SISVAN-Web coverage for the different life phases considered in the study.

In order to analyze aspects of the implementation of SISVAN-Web, a questionnaire was applied to the managers responsible for the System in both municipalities. The questionnaire addressed data related to the profile of these managers responsible for feeding the system and transferring information (education and training for implementation of the System), ease of use (user interface, information structure, navigation, search

system efficiency, loading time of data files, feeding and flow) and the use of information (reliability: use of protocols, coverage: focused on vulnerable segments or in all life cycles; origin: data exclusive of the health sector or population-based data; nature: anthropometric and/or food consumption data; and intersectorial character: restriction to demands of the health sector or coverage to other sectors such as food security, Family Grant Program and school diet).

Data analysis

Data were organized in spreadsheets and typed in double entry. The Validate application of the Epi Info *Software* version 3.3.2 was used to analyze the consistency of the data.

Three types of comparisons were made in order to test the reliability of anthropometric measurements:

- inter-equipment, considering the measurements obtained by the research team using their own equipment vs. the measurements obtained by the research team using the service equipment;
- inter-examiners (anthropometric techniques), considering the measurements obtained by the research team using the service equipment vs. the measurements obtained in the routine of the service;
- inter-examiners and inter-equipment, considering the measurements obtained by the research team using their own equipment vs. the measurements obtained in the routine of the service.

A t-test to paired samples was applied to verify the difference in means of weight (kg) and length/height (cm) in the three situations. In order to know the degree of agreement between nutritional classifications, the classification obtained by the research team using their own equipment was used as the "gold standard" to comparisons with the other two classifications. The degree of agreement was verified through the Cohen's Kappa Coefficient. Landis & Koch' criterion was used for interpretation of the agreement [19].

The level of significance adopted was 5%. The *Software* Stata version 12.0 was used.

Ethical aspects

This study was approved by the Research Ethics Committee of the State University of Paraíba under protocol number 19689613.3.0000.5187. All participants signed an Informed Consent form.

Results

A total of 319 under-five children were examined for anthropometric evaluation, 152 from one municipality and 167 from the other. However, the pairing of anthropometric data obtained by the research

anthropometrists with their own equipment was possible for 266 cases on weight and 286 on length/height, when they used the equipment of the health services. For data obtained by health professionals, pairing was possible for 132 cases on weight, and 137 cases on length/height. Losses in the first situation were due to the lack or poor quality of anthropometric equipment in some health units. Losses of data obtained in the service were due to lack of measurement and/or lack of registration of measures (our study did not intervene in the work routine). It is important to highlight that the number of children analyzed vs. losses per municipality did not differ with respect to nutritional status.

Table 1 presents the description of the anthropometric status of the study sample. The prevalence of overweight for age, overweight and stunting were 10.28%, 14.02% and 7.16%, respectively. None of the children were diagnosed with weight-for-age deficit (Z-score W/A < -2) nor with weight deficit (Z-score BMI/A < -2).

When analyzing the measurements obtained with the equipment of the health service, it was observed

that health professionals obtained smaller values of weight and height in 540 g and 3.17 cm, respectively, being these differences statistically significant. Similar results were found in the measurements of the research anthropometrists with their own equipment in comparison to the ones that they obtained using the equipment of the service (Table 2).

Table 3 shows the agreement between classifications of the nutritional status of the children according to anthropometric evaluation of the research anthropometrists with their own equipment (gold standard) with the other two classifications. It was noticed that lower agreement values were found for classifications related to the measurements obtained by the health professionals, being worse for H/A (Kappa Coefficient = 0.38) and BMI/A (Kappa Coefficient = 0.39). Health professionals overestimated overweight and stunting. When the parameter was the equipment, substantial agreement values were obtained for the three anthropometric indices, being higher for H/A (Kappa Coefficient = 0.74).

Table 1. Description of anthropometric status of under-five children living in two municipalities in the state of Paraíba, 2014

Variables	n	%
<i>Weight/Age</i>		
Adequate (Z-scores ≥ -2 and ≤ 2)	286	89.72
Excessive (Z-scores > 2)	33	10.28
Mean \pm Standard deviation (Z-score)	0.431 \pm 0.266	
<i>BMI/Age</i>		
Adequate (Z-scores ≥ -2 and ≤ 2)	274	85.98
Excessive (Z-scores > 2)	45	14.02
Mean \pm Standard deviation (Z-score)	0.968 \pm 0.302	
<i>Height/Age</i>		
Adequate (Z-scores ≥ -2)	296	92.84
Deficit (Z-scores < -2)	23	7.16
Mean \pm Standard deviation (Z-score)	-0.444 \pm 0.256	

Table 2. Differences in the measurement of weight and length/height of under-five children assisted by Family Health Strategy health teams of two municipalities of Paraíba, according to the origin of the measures, 2014.

Statistical Parameters	Anthropometric parameters					
	Weight (kg)			Length/Height (cm)		
	P	PS	S	P	PS	S
Number of observations	266	266	132	286	286	137
Mean	10.65	10.61	10.07	75.27	75.50	72.33
Standard error	0.23	0.23	0.33	0.79	0.78	1.06
Standard deviation	3.74	3.71	3.75	13.46	13.30	12.41
<i>Difference of means</i>						
- P vs. PS (inter-equipment analysis)	0.04 p=0.87			-0.23 p=0.48		
- PS vs. S (inter-examiner/technique analysis)	0.54 p=0.03			3.17 p=0.01		
- P vs. S (inter-equipment and inter-examiners analysis)	0.58 p=0.02			2.94 p=0.03		

P = Measurement obtained by the researcher using high reliability equipment (gold standard measure).

PS = Measurement obtained by the researcher, using the equipment of the service (the same used in “S” by the professional of the health unit).

S = Measurement obtained in the routine of the service.

Table 3. Agreement between anthropometric classifications of under-five children assisted by Family Health Strategy teams in two municipalities of Paraíba, according to the origin of the measurements, 2014.

P	Nutritional diagnosis								Kappa	p-value		
	PS			Kappa	p-value	S					Kappa	p-value
	Eutrophy	Malnutrition	Excessive			Eutrophy	Malnutrition	Excessive				
Weight/Age				0.66	<0.001				0.56	<0.001		
Eutrophy	227	3	8			109	3	4				
Excessive	7	0	21			6	0	10				
BMI/Age				0.64	<0.001				0.39	<0.001		
Eutrophy	193	4	13			83	3	11				
Excessive	8	0	29			9	0	12				
Height/Age				0.74	<0.001				0.38	<0.001		
Eutrophy	260	5	-			121	7	-				
Malnutrition	5	16	-			5	4	-				

P = Measurement obtained by the researcher using high reliability equipment (gold standard measure).

PS = Measurement obtained by the researcher, using the equipment of the service (the same used in “S” by the professional of the health unit).

S = Measurement obtained in the routine of the service.

The coverages of SISVAN-Web were 2.8% and 3.3%. In both municipalities, higher coverage values occurred among children, being 6.5% and 10.1% in the 0-4 year age range and 8.4% and 7.1% for 5-10 years. The elderly presented the lowest coverage, of 0.3% and 0.7%. Among adolescents, coverage was 2.1% and 1.8%; and among adults, 2.3% and 2.2%.

Regarding the answers of SISVAN-Web managers, it was verified that both municipalities had one single person responsible for feeding and transferring information, and this person had complete secondary education and was not trained for the implementation of the System. Regarding ease of use, the efficiency of the searching system was the only aspect perceived as bad by one of the managers. Regarding the use of information, the two managers reported not using protocols, approaching all life cycles, being based exclusively on health care services and restricted to the demands of the health sector. The use of anthropometric data and food consumption was pointed out by one of the managers, while the other pointed exclusive use of anthropometry.

Discussion

Generally, this study indicate several limitations in relation to the SISVAN-Web, such as poor structure, low coverage, lack of integration with other systems and low reliability of anthropometric measurements. This indicates little progress since these factors are highlighted in the literature prior to the implantation of the computerizations of the System [4, 5, 8].

According to Cohen's Kappa coefficients, the results obtained indicate errors in the nutritional classification, due both to the techniques used by health professionals and to problems of the equipment available in the services. This reality compromises the SISVAN-Web information. The diagnosis of insufficient linear growth (smaller values of stature) becomes overestimated, which is reflected in higher values of BMI/A (greater chance of being overweight). Low agreement between anthropometric measures obtained in the basic health network have also been reported by other researchers [13, 20, 21]. In this study, failure to observe the procedures adopted by health professionals makes it impossible to specify deficiencies in measurement techniques. However, the diversity, poor positioning and lack of calibration of instruments available in the health services were problems detected that may be related to the observed variations, which coincides with previous reports [20].

The results of the present study revealed very low coverage, similar to those found in other localities [1, 2, 9], and against the increase in the number of follow-

ups suggested after computerization of the SISVAN [4]. These findings are reason for concern, especially considering the importance of the System for decisions, programs and policies related to nutrition actions, including those related to the promotion of healthy food practices and the management of the Family Grant Program [3, 22].

Regarding the population coverage in the different phases of life, the highest coverage recorded among children and the lower among the elderly coincides with the results of a national survey⁴ and of more recent studies [1, 2]. This situation may be related to the historical priority of child-centered surveillance [5, 22]. Despite the positive consequences derived from this finding, i.e. favoring the care of children who are biologically vulnerable, ignoring the elderly population limits the care to a group that has increased proportionally in the population and who presents significant risks of infectious, deficiency and chronic diseases [1, 23].

The lack of training to implement the SISVAN-Web of those responsible for feeding the system and transferring information in the municipalities of the current study has been previously reported by professionals who carry out similar actions [9]. The need to promote professional training and retraining is, therefore, encouraged so as to improve the reliability of the information generated by the System [1]. The non-use of protocols, indicated by the managers interviewed, can also be detrimental to the credibility of data [4, 22]. It is necessary that the guiding documents of the SISVAN actions [10, 11, 15] be available in the health services and be known and used, which does not seem to be the reality in the country [22, 24]. More favourable situation was found in Minas Gerais, where 68.6% of those responsible for the System indicated to use technical documents of the Ministry of Health [25].

In general, the managers' perception of ease of use is indicative of good handling of SISVAN-Web. This result is consistent with the proposal of computerization of the System and the need for a friendly user interface [2].

As verified in Belo Horizonte [9], the non-use of food consumption data was indicated in one of the municipalities of the current study. In Minas Gerais, the registry of this information was informed by only 1/3 of those responsible for the system [25]. This situation is worrisome due to the importance of knowing food consumption data to subsidize the implantation and development of nutritional interventions [26], especially in the current situation marked by major changes in the food pattern associated with social transformations of modern life [3]. These changes include increased consumption of processed foods in detriment of fruits and vegetables, with possible negative effects on

children's health, which demands the implementation of actions to promote healthy eating [3, 27].

The NFNP suggests the issue of food and nutrition be addressed through the articulation of health with other sectors, such as education and social assistance, and programs, such as the Family Grant Program, in a way that will break the history of overlapping and disarticulated actions [3]. However, the report of information generated by SISVAN-Web restricted to the health sector was rather observed in the present study, likewise the observed in Minas Gerais [26]. It is necessary to recognize that the concretization of intersectoriality is related to the historical process of strengthening and expanding the concept of food and nutritional security, in a conjuncture of governmental integration with civil society and technical levels [3, 27].

The prevalence of overweight and stunting found in this study is in line with the findings reported in a meta-analysis of samples from public health services [28]. Another study that also focused on SISVAN presented similar results [13]. Despite expressive prevalences that reveal the importance of monitoring the growth of children assisted in public health services, many of them did not have their weight and/or height measured in the current study, which converges with results of previous studies [20, 29]. Possible characteristics of professionals associated with this situation such as training, skills and motivation [13, 29], were not analyzed in this study.

The limitations of this study are related to the caution to extrapolate the results, as they represent the reality of two municipalities only. Furthermore, information related to managers may involve information bias. However the results are important considering that research about the validity and accuracy of measured anthropometric data is lacking. The accuracy of anthropometric data is critical to produce robust and interpretable findings. Future researches are important to identify the provision of appropriate training and assessment as essential for health professionals responsible for collecting routine data.

Conclusion

This study allowed us to illustrate problems arising from the reliability of anthropometric data of under-five children obtained in public health services, low SISVAN-Web coverage, and restrictions on the use of information from the System in two municipalities in the state of Paraíba. It is necessary to invest in structure and human resources with a view to the adequate the operationalization of SISVAN-Web and monitoring of nutritional status. These arrangements are fundamental for the planning, management and evaluation of food and nutrition actions under the PHC in the municipalities studied. Further evaluative studies of SISVAN-Web are

necessary for its improvement and for the implementation of NFNP guidelines.

Acknowledges

Financial support by National Council for Scientific and Technological Development and Foundation for Research Support of the State of Paraíba.

References

- Enes CC, Loiola H, Oliveira MRM. Cobertura populacional do Sistema de Vigilância Alimentar e Nutricional no Estado de São Paulo, Brasil. *Cien Saude Colet.* 2014;19(5):1543-51.
- Jung NM, Bairros FS, Neutzling MB. Utilização e cobertura do Sistema de Vigilância Alimentar e Nutricional no Estado do Rio Grande do Sul, Brasil. *Cien Saude Colet.* 2014;19(5):1379-88.
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Política Nacional de Alimentação e Nutrição. Brasília (DF): Editora do Ministério da Saúde; 2012. (Série B. Textos Básicos de Saúde).
- Coutinho JG, Cardoso AJC, Toral N, Silva ACF, Ubarana JA, Aquino KKNC, et al. A organização da Vigilância Alimentar e Nutricional no Sistema Único de Saúde: histórico e desafios atuais. *Rev Bras Epidemiol.* 2009;12(4):688-99.
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Indicadores de Vigilância Alimentar e Nutricional: Brasil 2006. Brasília (DF): Editora do Ministério da Saúde; 2009. (Série B. Textos Básicos de Saúde).
- Bagni UV, Barros DC. Capacitação em antropometria como base para o fortalecimento do Sistema de Vigilância Alimentar e Nutricional no Brasil. *Rev Nutr.* 2012;25(3):393-402.
- Camilo SMB, Camilo GBC, Toledo GC, Camilo Júnior RD, Toledo CC. Vigilância nutricional no Brasil: criação e implementação do SISVAN. *Rev APS.* 2011;14(2):224-8.
- Pereira SMP, Brito LAMH, Palácio MAV, Monteiro MPA. Operacionalização do sistema de vigilância alimentar e nutricional em Juazeiro do Norte, Ceará. *Rev Baiana Saude Publica.* 2012;36(2):577-86.
- Ferreira CS, Cherchiglia ML, Cesar CC. O Sistema de Vigilância Alimentar e Nutricional como instrumento de monitoramento da Estratégia Nacional para Alimentação Complementar Saudável. *Rev Bras Saude Mater Infan.* 2013;13(2):167-77.
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Orientação para avaliação de marcadores de consumo alimentar na atenção básica. Brasília (DF): Editora do Ministério da Saúde; 2015.
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Marco de referência da vigilância alimentar e nutricional na atenção básica. Brasília (DF): Editora do Ministério da Saúde; 2015.
- Vasconcelos IAL, Santos LMP. Research potential of food and nutrition in the Family Health Strategy: A structured review. *Rev Nutr.* 2014; 27(1):107-26.
- Damé PKV, Pedrosa MRO, Marinho CL, Gonçalves VM, Duncan BB, Fisher PD, et al. Sistema de Vigilância Alimentar e Nutricional (SISVAN) em crianças do Rio Grande do Sul, Brasil: cobertura, estado nutricional e confiabilidade dos dados. *Cad Saude Pública.* 2011;27(11):2155-65.

- 14 Leão CDA, Caldeira AP. Avaliação da associação entre qualificação de médicos e enfermeiros em atenção primária em saúde e qualidade da atenção. *Cienc Saude Colet*. 2011;16(11):4415-23.
- 15 Ministério de Saúde do Brasil. Protocolos do Sistema de Vigilância Alimentar e Nutricional – SISVAN na assistência à saúde. Brasília: Ministério da Saúde; 2008. (Série B. Textos Básicos de Saúde).
- 16 World Health Organization. WHO Child Growth Standards. Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age. Methods and development. Geneva: World Health Organization; 2006.
- 17 Instituto Brasileiro de Geografia e Estatística. Censo demográfico 2010. Dados populacionais dos municípios [documento en Internet] 2010. [Acceso 26 de enero de 2017]. Disponible en: <http://www.ibge.gov.br/cidadesat/>.
- 18 Ministério de Saúde do Brasil. Informações de Saúde Suplementar-Beneficiários [documento en Internet] 2010. [Acceso 26 de enero de 2017]. Disponible en: <http://www2.datasus.gov.br/DATASUS>.
- 19 Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977; 33: 15974.
- 20 Lima MAA, Oliveira MAA, Ferreira HS. Confiabilidade dos dados antropométricos obtidos em crianças atendidas na Rede Básica de Saúde de Alagoas. *Rev Bras Epidemiol*. 2010; 13(1): 69-82.
- 21 Bryant M, Santorelli G, Fairley L, Petherick ES, Bhopal R, Lawlor DA, et al. Agreement between routine and research measurement of infant height and weight. *Arch Dis Child*. 2015;100:24-9.
- 22 Jaime PC, Silva ACF, Lima AMC, Bortolini GA. Ações de alimentação e nutrição na atenção básica: a experiência de organização no Governo Brasileiro. *Rev Nutr*. 2011;24(6):809-24.
- 23 Mayer BLD, Stumm EMF, Barbosa DA, Guido LA, Kirchner RM. Reflexões acerca do envelhecimento, das doenças crônicas e da qualidade de vida em renais crônicos. *Rev Contexto Saúde*. 2011;10(20):1315-8.
- 24 Pinheiro ARO, Recine E, Alencar B, Fagundes AA, Sousa JS, Monteiro RA, et al. Percepção de professores e estudantes em relação ao perfil de formação do nutricionista em saúde pública. *Rev Nutr*. 2012;25(5):631-43.
- 25 Rolim MD, Lima SML, Barros DC, Andrade CLT. Avaliação do SISVAN na gestão de ações de alimentação e nutrição em Minas Gerais, Brasil. *Cien Saude Colet*. 2015;20(8):2359-69.
- 26 Figueroa Pedraza D, Queiroz D, Gama JSFA. Consumo alimentar de crianças brasileiras assistidas em creches: uma revisão sistemática. *Rev Bras Saude Matern Infant*. 2015; 15 (1):17-31.
- 27 Rigon AS, Schmidt ST, Bógus CM. Desafios da nutrição no Sistema Único de Saúde para construção da interface entre a saúde e a segurança alimentar e nutricional. *Cad Saude Publica*. 2016; 32 (3):e00164514.
- 28 Sousa CPC, Olinda RA, Figueroa Pedraza D. Prevalence of stunting and overweight/obesity among Brazilian children according to different epidemiological scenarios: systematic review and meta-analysis. *Sao Paulo Med J*. 2016; 134 (3):251-62.
- 29 Figueroa Pedraza D. Growth surveillance in the context of the Primary Public Healthcare Service Network in Brazil: literature review. *Rev Bras Saude Matern Infant*. 2016; 16 (1):7-19.