

Clinical and Microbiological Characterization of Osteoarticular Infections in Children from a Clinic in Rionegro, Antioquia

Álvaro Hoyos-Orrego¹ , Natalia Echeverri-Ramírez² , Santiago Moreno-Ramírez² 
Jhojan Herrera-Vargas³ 

¹Pediatrician, specialist in infectious diseases, Clínica Somer, Rionegro, Colombia.

²Physician, Clínica Somer, Rionegro, Colombia.

³Respiratory therapist and epidemiologist, Clínica Somer, Rionegro, Colombia.

ARTICLE INFORMATION

KEYWORDS

Anti-Bacterial Agents;
Osteomyelitis;
Staphylococcus aureus

Received: July 29, 2024

Accepted: September 10, 2024

Correspondence:

Álvaro Hoyos-Orrego;
alvaromicro@hotmail.com

How to cite: Hoyos-Orrego A, Echeverri-Ramírez N, Moreno-Ramírez S, Herrera-Vargas J. Clinical and Microbiological Characterization of Osteoarticular Infections in Children from a clinic in Rionegro, Antioquia. *Iatreia* [Internet]. 2025 Jul-Sept;38(3):424-437. <https://doi.org/10.17533/udea.iatreia.328>



Copyright: © 2025
Universidad de Antioquia.

ABSTRACT

Introduction: Osteomyelitis and septic arthritis are common invasive bacterial infections in children.

Objective: To describe the microorganisms, antimicrobial susceptibility, and microbiological diagnostic procedures.

Methods: A case series of children with a confirmed diagnosis, from 2017 to 2022, was characterized.

Results: Forty-eight out of 84 patients were included in the study. The diagnoses were osteomyelitis in 20.8%, septic arthritis in 37.5% and osteoarthritis in 41.6% of cases. Isolated microorganisms included methicillin-susceptible *Staphylococcus aureus* (MSSA) in 30.7% of cases, methicillin-resistant *S. aureus* (MRSA) in 28.8%, *Pseudomonas aeruginosa* in 5.7%, *S. epidermidis* in 3.8%, *Escherichia coli* in 3.8%, *Enterococcus faecalis* in 3.8%, *Aeromonas hydrophila* in 3.8%, *Mycobacterium tuberculosis* in 1.9%, and *Salmonella* spp. in 1.9%, among others. In 40 infections originating from skin and soft tissues, orthopedic hardware, limb trauma, and without an identified source, MSSA/MRSA predominated (75%). MSSA/MRSA showed susceptibility to vancomycin at 100%, trimethoprim-sulfamethoxazole at 100% and 93%, and clindamycin at 100% and 86.6%, respectively. Blood cultures (n=22), synovial tissue cultures (n=7), bone cultures (n=28), and bone pus or synovial fluid in blood culture bottles (n=3) were 100% positive.

Conclusions: *S. aureus* was the most frequent microorganism with a high proportion of MRSA. MRSA exhibited high susceptibility to trimethoprim-sulfamethoxazole and vancomycin. The high positivity of primary cultures suggests the importance of always obtaining them.

Caracterización clínica y microbiológica de las infecciones osteoarticulares en niños de una clínica en Rionegro, Antioquia

Álvaro Hoyos-Orrego¹ , Natalia Echeverri-Ramírez² , Santiago Moreno-Ramírez² ,
Jhojan Herrera-Vargas³ ,

¹Pediatra especialista en enfermedades infecciosas, Clínica Somer, Rionegro, Colombia.

²Médico(a) general, Clínica Somer, Rionegro, Colombia.

³Terapeuta respiratorio y epidemiólogo, Clínica Somer, Rionegro, Colombia.

INFORMACIÓN ARTÍCULO

PALABRAS CLAVE

Antibacterianos;
Osteomielitis;
Staphylococcus aureus

Recibido: julio 29 de 2024

Aceptado: septiembre 10 de 2024

Correspondencia:

Álvaro Hoyos-Orrego;
alvaromicro@hotmail.com

Cómo citar: Hoyos-Orrego A, Echeverri-Ramírez N, Moreno-Ramírez S, Herrera-Vargas J. Caracterización clínica y microbiológica de las infecciones osteoarticulares en niños de una clínica en Rionegro, Antioquia. *Iatreia* [Internet]. 2025 Jul-Sept;38(3):424-437.

<https://doi.org/10.17533/udea.iatreia.328>



Copyright: © 2025
Universidad de Antioquia.

RESUMEN

Introducción: la osteomielitis y la artritis séptica son infecciones bacterianas invasoras comunes en niños.

Objetivo: describir los microorganismos, su susceptibilidad a los antibióticos y los procedimientos diagnósticos microbiológicos.

Métodos: se caracterizó una serie de casos en niños con diagnóstico confirmado entre 2017 y 2022.

Resultados: se estudiaron 48 de 84 pacientes. La osteomielitis, la artritis séptica y la osteoartritis representaron el 20,8 %, 37,5 % y 41,6 % de los casos, respectivamente. Los microorganismos aislados incluyeron *Staphylococcus aureus* susceptible a meticilina (SAMS) en el 30,7 % de los casos, *S. aureus* resistente a meticilina (SAMR) en el 28,8 %, *Enterococcus faecalis* en el 3,8 %, *Aeromonas hydrophila* en el 3,8 %, *Mycobacterium tuberculosis* en el 1,9 % y *Salmonella* spp. en el 1,9 %, entre otros. En 40 casos de infecciones originadas en piel y tejidos blandos, material de osteosíntesis, trauma sobre la extremidad y sin una fuente identificada, predominó SAMS/SAMR (75 %). La susceptibilidad de SAMS/SAMR a vancomicina fue del 100 %, para trimetoprim-sulfametoxazole del 100 % y 93 %, y clindamicina 100 % y 86,6 %, respectivamente. Los hemocultivos (n = 22), cultivos de tejido sinovial (n = 7), hueso (n = 28) y pus de hueso o líquido sinovial en botella de hemocultivo (n = 3) fueron positivos en el 100 %.

Conclusiones: *S. aureus* fue el microorganismo más frecuente con una elevada proporción de SAMR, y presentó una alta susceptibilidad a trimetoprim-sulfametoxazol y vancomicina. La alta positividad de los cultivos primarios sugiere la importancia de siempre obtenerlos.

INTRODUCTION

Osteomyelitis, septic arthritis, and osteoarthritis (OA) are common invasive bacterial infections in children, primarily affecting those under 5 years old. The incidence of osteoarticular infections (OAI) is estimated at 10 to 80 cases per 100,000 children per year, with a lower rate in developed countries at 4 to 10 cases per 100,000 children per year (1-3). While *Staphylococcus aureus* has been the most common pathogen causing musculoskeletal infections, community-acquired methicillin-resistant *S. aureus* (CA-MRSA) has emerged as a significant microorganism involved in invasive infections, including OAI (1,4-6). In South America, various studies have reported CA-MRSA as a major etiological agent in cases of OAI, like other countries worldwide (7-12). In Colombia, the first report of multisystem invasive infection and osteomyelitis caused by CA-MRSA was documented in 2009 (13). Since then, additional cases have been described (14-16).

Other microorganisms causing OAI in children include *Streptococcus pyogenes*, *S. agalactiae*, *Neisseria gonorrhoeae*, *Escherichia coli*, *Salmonella* spp., *Kingella kingae*, and *Candida* spp. (5,11,17,18). With the introduction of conjugated vaccines into immunization schedules, microorganisms such as *S. pneumoniae* and *Haemophilus influenzae* serotype B have decreased as causative agents of invasive infections, including OAI (4,18).

Locally, the etiology of OAI in children is not well-characterized, and published data are scarce (19-23). Recognizing the microbiology involved in OAI is crucial, allowing for the selection of empirical antibiotic therapy based on epidemiology (7,9-11,24). The objective of this study was to describe the microbiology, antibiotic susceptibility, and microbiological procedures in children with OAI treated at a high-complexity clinic.

METHODOLOGY

The study was an observational and retrospective case series. Children with a confirmed diagnosis of OAI who were hospitalized at Clínica Somer (CS) in the city of Rionegro, Antioquia, Colombia, between January 1, 2017, and July 30, 2022, were included. CS is a high-complexity (fourth-level) institution that provides care for both adults and children, with a capacity of 110 pediatric beds distributed across emergency services, hospitalization, and intensive care units. CS serves as a referral center for complex patients from urban and rural areas inside and outside the state of Antioquia.

Medical records of patients diagnosed with septic arthritis and osteomyelitis, according to the International Classification of Diseases (ICD) valid at that time (25), were reviewed. A total of 84 medical records were evaluated. The study received approval from the Ethics and Research Committee of CS (Minutes number 54 dated July 15, 2022). Due to the nature of the study, informed consent was not required.

Inclusion and Exclusion Criteria

All patients aged over 30 days to 15 years, hospitalized with a diagnosis of osteomyelitis, septic arthritis, or acute, subacute, or chronic OA confirmed by a primary culture (bone, pus, tissue, or synovial fluid) or a positive blood culture were included.

Patients without microbiological confirmation, non-infectious arthritis, those under 1 month or over 15 years old, and those with incomplete medical information in the medical records were excluded. In patients with multiple positive cultures, only the first isolation was recorded.

Operational Definitions

Osteomyelitis, septic arthritis, and acute OA were defined when the duration of symptoms was ≤ 14 days. Subacute and chronic infections were considered if the symptom duration was less than 3

months or more than 3 months, respectively (25).

Microbiology

The cultured samples included blood, bone biopsy, bone pus, tissue, and synovial fluid. The samples collected and the types of cultures ordered depended on the orthopedic surgeon's criteria. The cytological study of synovial fluid was not included in the analysis. Samples for microbiological study were processed according to the laboratory's microbiology protocols. The laboratory does not have any molecular techniques for the microbiological diagnosis of OAI.

Conventional Cultures

Samples were inoculated on MacConkey agar, chocolate agar, and blood agar. Synovial or bone tissue was also inoculated in brain heart infusion (BHI) enriched medium. For mycobacteria, Ogawa-Kudoh solid medium was used, following protocols from the National Institute of Health of Colombia and recommendations by the Pan American Health Organization (26).

Blood Cultures

Bottles inoculated with blood, bone pus, or synovial fluid were incubated for 7 days in the automated Bactec™ FX system (Becton Dickinson, USA). Identification and antibiotic susceptibility testing of microorganisms isolated from conventional media or blood cultures were performed using the automated Phoenix™ M50 system (Becton Dickinson, USA), applying susceptibility breakpoints valid for the date of isolations, according to the Clinical and Laboratory Standards Institute (CLSI) (27). The antibiotics analyzed for gram-positive cocci other than *S. aureus* included oxacillin, vancomycin, penicillin, ampicillin, clindamycin, trimethoprim-sulfamethoxazole, and erythromycin. For enteric gram-negative bacilli, cefepime, imipenem, meropenem, and ciprofloxacin were tested. In the case of *Pseudomonas aeruginosa* and other environmental gram-negative bacilli, the antibiotics included ceftazidime, cefepime, piperacillin-tazobactam, imipenem, meropenem, and ciprofloxacin. For *Mycobacterium tuberculosis*, rifampicin, isoniazid, and ethambutol were tested. In the case of *S. aureus*, susceptibility was assessed for vancomycin, oxacillin, clindamycin, erythromycin, and trimethoprim-sulfamethoxazole. Vancomycin susceptibility was categorized according to the reported minimum inhibitory concentrations (MICs): ≤ 0.5 $\mu\text{g/mL}$, 1 $\mu\text{g/mL}$, and >1 $\mu\text{g/mL}$ (27).

Data Collection and Statistical Analysis

Demographic, clinical, and microbiological data of the individuals of interest were collected. Age, sex, origin (metropolitan area of Medellín, eastern Antioquia, and others); type of health insurance affiliation; affected joint, bone, or both joint and bone; acute, subacute, or chronic infection; and medical history such as trauma to the affected limb, skin or soft tissue infection (furuncle, cellulitis, or abscess), infected trauma wound, orthopedic hardware, fractures, or no identifiable factor were recorded.

Microbiological information included the microorganism's name, susceptibility to the evaluated antibiotics, obtained cultures, sample types, and prior antibiotic use before culturing. The information was collected exclusively by one of the researchers and recorded using an Excel™ database. The open-source statistical package PSPP was used, applying descriptive statistics that included absolute frequencies and proportions for qualitative variables.

RESULTS

A total of 84 medical records with a diagnosis of septic arthritis or osteomyelitis according to the ICD were reviewed. Out of the 84 records, 21 patients were excluded due to diagnoses of non-infectious arthritis, 2 with less than 1 month, 12 with negative cultures for OAI (which were treated

as culture-negative infections), and 1 case due to incomplete information, resulting in a total of 48 medical records corresponding to an equal number of patients. The number of negative cultures was not included in the analysis.

Among the 48 patients, the majority were males (68.7%), and 50% had health insurance under the contributory regime. 58% of the patients were over 6 years old. 89.5% of the patients came from the eastern region of Antioquia or from other departments. Regarding the type of infection, acute OA predominated (37.5%), followed by acute septic arthritis (20.8%), acute osteomyelitis (16.7%), and chronic osteomyelitis (14.6%). The femur was the primary bone involved (29.2%), followed by the tibia (20.8%), and the most affected joints were the knee and the hip (14.6% each) (Table 1).

Table 1. Demographic and general clinical characteristics

Characteristics		n=48 (%)
Sex	Male	33 (68.7)
	Female	15 (31.2)
Health affiliation	Contributory	24 (50.0)
	Subsidised	23 (47.9)
	Other	1 (2.0)
Age		
	1-12 months	9 (18.7)
	1-5 years old	11 (22.9)
	6-10 years old	13 (27.0)
	11-15 years old	15 (31.2)
Origin		
	Medellín Metropolitan Area	5 (10.4)
	East of Antioquia	25 (52.0)
	Other	18 (37.5)

Table 1. Demographic and general clinical characteristics. Continued

Characteristics		n=48 (%)
Type of infection		
	SA	10 (20.8)
	OM	11 (22.9%)
	COM	7 (14.6)
	OA	20 (41.6%)
Bone and/or joint involved *		
	Femur	14 (29.2)
	Tibia	10 (20.8)
	Knee	7 (14.6)
	Hip	7 (14.6)
	Humerus	6 (12.5)
	Elbow	6 (12.5)
	Ankle	5 (10.4)
	Fibula	2 (4.2)
	Shoulder	2 (4.2)
	Ulna	2 (4.2)
	Other†	7 (14.6)

*Some patients presented more than one compromised anatomical site. †Hallux, phalanx, radius, metatarsal. OA: osteoarthritis; OM: osteomyelitis; SA: septic arthritis; COM: chronic osteomyelitis
Source: Author's own work

In 37 (77.0%) patients, some history related to OAI was identified. Skin and soft tissue infections were the main antecedent (20/48; 41.5%), followed by open fractures (6/48; 12.5%) and the use of orthopedic hardware (6/48; 12.5%). Methicillin-susceptible *S. aureus* (MSSA) and methicillin-resistant *S. aureus* (MRSA) were the predominant microorganisms in OAI related to skin and soft tissue infections, osteosynthesis material, limb trauma, and infections without an identifiable antecedent, accounting for 85% (17/20), 50% (3/6), 100% (3/3), and 63.3% (7/11) of cases, respectively. Environmental and enteric gram-negative bacilli predominated in cases of open fractures (Figure 1).




Figure 1. Clinical history and identified microorganisms

*Furuncles, cellulitis or abscess. MSSA: methicillin susceptibility *Staphylococcus aureus*; MRSA: methicillin resistant *Staphylococcus aureus*.

Source: Author's own work

When evaluating age and its relationship with etiology, *S. aureus* infections predominated in all age ranges: 6/31 (19.3%) under 12 month, 6/31 (19.3%) between 1-5 years, and 19/31 (61.2%) over 6 years. Infections with environmental gram-negative bacilli (*P. aeruginosa*, *Aeromonas hydrophila*, *Acromobacter xylosoxidans*) mostly occurred in those over 6 years old. In cases of enteric gram-negative bacilli, the ages with the most cases were under 12 months and over 6 years old (Table 2).

Table 2. Etiology and distribution by age

Microorganisms	Age	1-12 months	1-5 years old	6-10 years old	11-15 years old	Total*
MSSA		3	2	6	5	16
MRSA		3	4	3	5	15
<i>Staphylococcus capitis</i>		-	1	-	-	1
<i>S. epidermidis</i>		-	-	-	2	2
<i>Escherichia coli</i>		1	1	-	-	2
<i>Enterobacter asburiae</i>		-	-	-	1	1
<i>Citrobacter koseri</i>		1	-	-	-	1
<i>Mycobacterium tuberculosis</i>		-	-	1	-	1
<i>Salmonella spp.</i>		1	-	-	-	1
<i>Bacillus cereus</i>		-	1	-	-	1
<i>Pseudomonas aeruginosa</i>		-	1	2	-	3
<i>Aeromonas hydrophila</i>		-	-	-	2	2
<i>Enterobacter cloacae</i>		-	-	-	1	1
<i>Proteus vulgaris</i>		-	-	-	1	1
<i>S. sciuri</i>		-	-	-	1	1
<i>Serratia marcescens</i>		-	-	1	-	1
<i>Enterococcus faecalis</i>		-	1	1	-	2
<i>Acromobacter xylosoxidans</i>		-	-	1	-	1

*A total of 52 microorganisms were identified in 48 patients. In 4/6 open fractures, two microorganisms were isolated.

MSSA: methicillin susceptibility *Staphylococcus aureus*; MRSA: methicillin resistant *Staphylococcus aureus*

Source: Author's own work

The analysis of the evaluated antibiotics revealed that MSSA was more susceptible than MRSA (Table 3). For vancomycin, all *S. aureus* isolates were susceptible, with an MIC ≤ 1 μ g/mL. *Enterococcus faecalis* isolates were susceptible to ampicillin (100%), as were 100% of enteric and environmental gram-negative bacilli to carbapenems (Figure 2). Antibiotic susceptibility testing was not performed for *A. xylosoxidans* due to a lack of reference cut-offs.

Table 3. Antimicrobial susceptibility in *Staphylococcus aureus*

	No.	VAN / MIC (µg/mL)					
		CLI	ERY	SXT	≤0.5	1	>1
MSSA	16	16 (100%)	14 (87%)	16 (100%)	1 (7%)	15 (94%)	0
MRSA*	15	13 (86.6%)	11 (73%)	14 (93%)	1 (7%)	14 (93%)	0

* Positive D-test in one isolate. CLI: clindamycin; ERY: erythromycin; SXT: trimethoprim-sulfamethoxazole; VAN: vancomycin. MIC: minimal inhibitory concentration; MSSA: methicillin susceptibility *Staphylococcus aureus*; MRSA: methicillin resistant *Staphylococcus aureus*

Source: Author's own work




Figure 2. Identified microorganisms and their susceptibility to the analyzed antibiotics

**Staphylococcus epidermidis*: 2; *S. capitis*: 1; *S. sciuri*: 1. † *Escherichia coli*: 2; *Proteus vulgaris*: 1; *Citrobacter koseri*: 1; *Serratia marcescens*: 1; *Enterobacter cloacae*: 1; *E. asburiae*: 1; *Salmonella* spp.: 1. ‡ *Pseudomonas aeruginosa*: 3; *Aeromonas hydrophila*: 2. OXA: oxacillin; VAN: vancomycin; CLI: clindamycin; ERY: erythromycin; SXT: trimethoprim-sulfamethoxazole; AMP: ampicillin; GEN: gentamicin; CIP: ciprofloxacin; CAZ: ceftazidime; FEP: cefepime; TZP: piperacillin-tazobactam; IMI: imipenem; MER: meropenem

Source: Author's own work

Regarding the collected samples, primary cultures (bone, bone pus, synovial fluid, or tissue) were taken in all patients. Hemocultures were taken from 22 patients, with 100% positivity (Figure 3). Empirical antibiotics were initiated before the collection of primary cultures in 33 patients (68.7%).




Figure 3. Comparison among the different types of specimens cultured

Source: Author's own work

DISCUSSION

This study provided insights into the microbiology of OAI in a group of patients under 15 years old hospitalized at a high-complexity clinic in Rionegro, Antioquia. Most OAI cases typically occur in male patients under 5 years old (2,5,17,28). However, in the studied group, the majority of OAI cases occurred in males over 5 years old, consistent with recent publications by various authors (7,22,23). OAI in males over 5 years old may be related to a greater tendency for trauma because of the initiation of physical activity and contact sports in school-age children and adolescents (29,30). In our series, due to the low number of patients, OAI with negative cultures (not included in the analysis), and the fact that not all cases recorded the clinical history preceding the infection, this finding cannot be generalized.

At a local level, few studies describe the microbiology of OAI in children. Recently, in the city of Medellín, a descriptive retrospective study reported the microbiology of OAI in children from a university clinic (23). Of the 39 characterized cases, the main identified microorganisms were MSSA (53.8%), MRSA (20.5%), other gram-positive cocci (10%), *P. aeruginosa* (5.1%); *Enterobacter cloacae*, *Haemophilus haemolyticus*, *S. agalactiae*, *S. pyogenes*, *M. tuberculosis*, and *C. albicans*, each with one case (2.5%). In the same city, at San Vicente Foundation University Hospital, several researchers have reported the etiology of OAI at different times. Urrego JD *et al.* conducted a study to describe the epidemiological profile of patients aged 0 to 15 years with OAI. Of the 72 patients, the main identified microorganisms were MSSA (38.9%), MRSA (26.4%), *S. pyogenes* (2.5%), and *M. tuberculosis* (1.5%), among others (22). Atehortúa S *et al.* described the etiology in 60 patients under 12 years diagnosed with septic arthritis, with a positive culture (27/60; 45%). The distribution of isolates was as follows: MSSA (21.6%), MRSA (8.3%), *S. pyogenes* (3.3%), methicillin-resistant *S. epidermidis* (3.3%), *S. pneumoniae* (1.6%), and *N. meningitidis* (1.6%) (19). Sierra M *et al.* in a study on non-gonococcal septic arthritis in a group of patients over 12 years old (n=54) found that the main microorganism in synovial fluid culture was *S. aureus* in 75.5% of cases, followed by other gram-negative bacilli such as *Klebsiella* spp. (5.7%), *E. coli* (3.8%), *Salmonella* spp. (1.9%), and *Enterobacter* spp. (1.9%). In this latter study, a distinction between MSSA and MRSA was not made (20). Our results are similar to those reported locally, and it is noteworthy that no cases of *S. pneumoniae* or *H. influenzae* serotype B

were presented, which could be explained by the introduction of both conjugate vaccines into the expanded immunization program (8,9,11). It is also emphasized that no isolates of *K. kingae* were present, similar to the studies mentioned locally. Various hypotheses have been proposed to explain the difficulty in isolating *K. kingae* in conventional culture media, such as a low bacterial inoculum at the infection site, demanding nutritional requirements of the microorganism, and some growth-inhibiting factors present in traditional culture media (19). This last result contrasts with the recently published study by Cañete I *et al.*, who, in a study describing the characteristics of pediatric OAI in a hospital in Santiago, Chile, South America, found that in patients under 5 years old, the most frequent microorganism was *K. kingae* (n=12), followed by *S. aureus* (n=9). The researchers used different diagnostic techniques such as polymerase chain reaction (PCR), which could explain better performance in etiological diagnosis (8,19).

Regarding etiology, *S. aureus* was the most frequently identified microorganism. MRSA showed high susceptibility to trimethoprim-sulfamethoxazole, clindamycin, and vancomycin, suggesting a phenotype of community-acquired MRSA (CA-MRSA). The high proportion of positive isolates for MRSA (48.3%) is striking, representing almost half of the *S. aureus* isolates. When compared with recent reports of the microbiology of OAI in children locally, the identification of MRSA has ranged between 8.3 and 28% (19,22,23,31). In South America, reports of MRSA as a cause of OAI vary between 7.7% and 63.3% according to some authors (7-11). The data suggest an increase in OAI cases caused by MRSA, with a highly variable distribution even at local and regional levels. Finding a MRSA proportion of 48.3% in the studied group would indicate that initial therapeutic management should consider an effective antibiotic against MRSA (1,9,10,17,24,32).

When comparing different types of cultured samples, we found a very high proportion of primary cultures (bone, bone pus, synovial fluid, and tissue) and positive blood cultures (100%). Synovial fluid culture in conventional media had a lower positivity rate (68.1%). In OAI, cultures are typically positive in 40% to 60% of cases (6,10). In acute cases, blood cultures reach a positivity rate of 60% (9). Atehortua S *et al.* reported that, in confirmed cases of septic arthritis, 49% of blood cultures were positive, and in 36.9% of cases, synovial fluid cultures were negative, indicating the importance of routinely ordering blood cultures (19). Zuluaga AF *et al.* demonstrated, in cases of chronic osteomyelitis, that bone biopsy had a high diagnostic yield (94/100; 94%) and was superior to cultures from samples other than bone (21). Recently, it was reported that, in confirmed cases of IOA in children (n=39), bone, bone pus or synovial fluid cultures in blood culture bottles, and synovial tissue culture, showed the highest positivity rates at 96%, 87.5%, and 66.5%, respectively (23). The high proportion of positive cultures in our study could be explained by several reasons. First, the CS is an institution that receives patients referred from distant rural areas with complex and prolonged infectious processes that may take a long time to receive medical attention. Second, the orthopedic surgical team has prioritized the collection of samples with the best diagnostic performance in surgery, such as bone pus, bone and synovial tissue. Third, empirical antibiotics are initiated after blood cultures are taken in acute IOA cases. It has been mentioned that the use of antibiotics before sample collection in surgical culture may reduce diagnostic yield. Ratnayake K *et al.* in a study on acute osteomyelitis in children found that in 42/50 (84%) of cases, a positive primary culture was obtained after antibiotics were initiated (33). Other authors have reported similar findings (4,23,34). These data suggest that antibiotics in acute IOA could be initiated before surgical procedures, ensuring prior blood culture collection, proper sample collection in surgery, and the use of appropriate culture media. To increase microbiological recovery, the use of blood cultures to inoculate with pus or synovial fluid could be considered (19,28,35).

The study has several limitations, including the small sample size and its single-center design. Including patients retrospectively may mean that individuals with osteomyelitis or septic arthritis registered with different diagnoses may not have been identified. Information on the characteristics

of the synovial fluid cytological study was not included in cases of septic arthritis, and microbiological cultures were not uniformly taken in all patients. The lack of molecular microbiological diagnostic tests may limit the detection of difficult-to-isolate microorganisms such as *K. kingae*. The number of isolates and susceptibility to different antibiotics was low for microorganisms other than *S. aureus*, preventing the generalization of any recommendations for empirical treatment. However, this did not significantly impact the objectives and results of the research.

CONCLUSIONS

In this cohort, osteoarticular infections (IOA) predominated in school-age children and adolescents. *Staphylococcus aureus* was the most common etiological agent, with a significant proportion of MRSA. MRSA showed high susceptibility to trimethoprim-sulfamethoxazole and vancomycin. Enteric and environmental gram-negative bacilli predominated in open fractures. The high positivity of primary cultures suggests the importance of always obtaining them, prioritizing the collection of bone and joint tissue and bone pus. The results highlight the need for studies with a larger number of patients to better characterize the etiology of IOA in children, monitor antibiotic susceptibility over time, and establish clinical differences between infections caused by MSSA and MRSA.

FUNDING

None.

CONFLICT OF INTEREST

None to declare.

REFERENCES

1. Ju KL, Zurakowski D, Kocher MS. Differentiating between methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* osteomyelitis in children: an evidence-based clinical prediction algorithm. *J Bone Joint Surg Am* [Internet]. 2011;93(18):1693-1701. <https://doi.org/10.2106/JBJS.J.01154>
2. Pääkkönen M, Peltola H. Bone and joint infections. *Pediatr Clin North Am* [Internet]. 2013;60(2):425-436. <https://doi.org/10.1016/j.pcl.2012.12.006>
3. Grammatico-Guillon L, Maakaroun-Vermesse Z, Baron S, Gettner S, Rusch E, Bernard L. Paediatric bone and joint infections are more common in boys and toddlers: a national epidemiology study. *Acta Paediatr* [Internet]. 2013;102(3):e120-e125. <https://doi.org/10.1111/apa.12115>
4. Yi J, Wood JB, Creech CB, Williams D, Jimenez-Truque N, Yildirim I, *et al.* Clinical epidemiology and outcomes of pediatric musculoskeletal infections. *J Pediatr* [Internet]. 2021;234:236-244.e2. <https://doi.org/10.1016/j.jpeds.2021.03.028>
5. Autore G, Bernardi L, Esposito S. Update on acute bone and joint infections in paediatrics: a narrative review on the most recent evidence-based recommendations and appropriate antinfective therapy. *Antibiotics* [Internet]. 2020;9(8):486. <https://doi.org/10.3390/antibiotics9080486>
6. Kini AR, Shetty V, Kumar AM, Shetty SM, Shetty A. Community-associated, methicillin-susceptible, and methicillin-resistant *Staphylococcus aureus* bone and joint infections in children: experience from India. *J Pediatr Orthop B* [Internet]. 2013;22(2):158-166. <https://doi.org/10.1097/BPB.0b013e32835c530a>

7. Hänel A, Rodríguez P, Silva A, Meza G, Piñera C. Infecciones osteoarticulares en niños: experiencia de 5 años. *Rev Chilena Infectol* [Internet]. 2020;37(6):742-749. <http://dx.doi.org/10.4067/S0716-10182020000600742>
8. Cañeta-Campos I, Moller-Macherone F, Figueroa-Gatica MJ, Monge-Iriarte M, Le Corre-Pérez N, Vizcaya-Altamirano MC, *et al.* Infecciones osteoarticulares en población pediátrica: clínica y microbiología de los últimos 15 años. *Rev Chilena Infectol* [Internet]. 2022;39(6):706-712. <http://dx.doi.org/10.4067/S0716-10182022000600706>
9. Highton E, Pérez MG, Cedillo-Vilamagua C, Sormani MI, Mussini MS, Isasmendi A, *et al.* Infecciones osteoarticulares en un hospital pediátrico de alta complejidad: epidemiología y características clínicas asociadas con bacteriemia. *Arch Argent Pediatr* [Internet]. 2018;116(2):e204-e209. <http://dx.doi.org/10.5546/aap.2018.e204>
10. Rosanova MT, Berberian G, Bologna R, Giménez S, Sarkis C, Buchovsky A, *et al.* Estudio descriptivo de infecciones osteo-articulares en niños en tiempos de *Staphylococcus aureus* resistente a metilicina de la comunidad (SARM-Co). *Rev Chilena Infectol* [Internet]. 2015;32(3):321-325. <https://doi.org/10.4067/S0716-10182015000400010>
11. Zunino C, Vomero A, Pandolfo S, Gutiérrez C, Algorta G, Pérez MC, *et al.* Etiología y evolución de las infecciones osteo-articulares 2009-2015. Hospital Pediátrico del Centro Hospitalario Pereira Rossell, Uruguay. *Rev Chilena Infectol* [Internet]. 2017;34(3):235-242. <https://doi.org/10.4067/S0716-10182017000300005>
12. Jaña-Neto FC, Sartori-Ortega C, De Oliveira-Goiano E. Epidemiological study of osteoarticular infections in children. *Acta Ortop Bras* [Internet]. 2018;26(3):201-205. <https://doi.org/10.1590/1413-785220182603145650>
13. Álvarez-Olmos MI, Enríquez SP, Pérez-Roth E, Méndez Álvarez S, Escobar J, Vanegas N, *et al.* Pediatric cases from Colombia caused by a Panton-Valentine Leukocidin-positive community-acquired methicillin-resistant *Staphylococcus aureus* ST8-SCCmecIVc clone. *Pediatr Infect Dis J* [Internet]. 2009;28(10):935. <https://doi.org/10.1097/INF.0b013e3181b2102b>
14. Machuca MA, González CI, Sosa LM. *Staphylococcus aureus* resistente a metilicina causante de infecciones comunitarias y de infecciones asociadas a la atención en salud en pacientes pediátricos del Hospital Universitario de Santander. *Biomédica* [Internet]. 2014;34(Supl.1):163-169. <https://doi.org/10.7705/biomedica.v34i0.1648>
15. Sosa-Ávila LM, Machuca-Pérez MA, Sosa-Ávila CA, González-Rugeles CI. Infecciones por *Staphylococcus aureus* metilino resistente en niños de Bucaramanga Colombia. *Salud UIS* [Internet]. 2010;42(3):248-255. Disponible en: <https://revistas.uis.edu.co/index.php/revistasaluduis/article/view/1680>
16. Jiménez JN, Ocampo AM, Vanegas JM, Rodríguez EA, Garcés CG, Patiño LA, *et al.* Characterization of virulence genes in methicillin susceptible and resistant *Staphylococcus aureus* isolates from a paediatric population in a university hospital of Medellín, Colombia. *Mem Inst Oswaldo Cruz* [Internet]. 2011;106(8):980-985. <https://doi.org/10.1590/s0074-02762011000800013>
17. Congedi S, Minotti C, Giaquinto C, Da Dalt L, Donà D. Acute infectious osteomyelitis in children: new treatment strategies for an old enemy. *World J Pediatr* [Internet]. 2020;16:446-455. <https://doi.org/10.1007/s12519-020-00359-z>
18. Schmitt SK. Osteomyelitis. *Infect Dis Clin North Am* [Internet]. 2017;31(2):325-338. <https://doi.org/10.1016/j.idc.2017.01.010>
19. Atehortua S, Woodcock S, Naranjo J, Faraco F, Uribe-Ríos A. Diagnóstico microbiológico de artritis séptica en niños usando botellas de hemocultivos como un método alternativo. *Infectio* [Internet]. 2020;24(2):100-104. <https://doi.org/10.22354/in.v24i2.840>

20. Sierra M, Manjarres G, Ramírez LA, Felipe O, Uribe O. Artritis séptica no gonococcica en el Hospital Universitario San Vicente de Paúl (1984-1991). *Iatreia* [Internet]. 1994;7(1):29-33. <https://doi.org/10.17533/udea.iatreia.3538>
21. Zuluaga AF, Galvis W, Saldarriaga JG, Agudelo M, Salazar BE, Vesga O. Etiologic diagnosis of chronic osteomyelitis. A prospective study. *Arch Intern Med* [Internet]. 2006;166(1):95-100. <https://doi.org/10.1001/archinte.166.1.95>
22. Urrego-Rendón JD, Uribe-Ríos A. Perfil epidemiológico de los pacientes entre 0 y 15 años con infección osteoarticular en el Hospital Infantil San Vicente Fundación. *Iatreia* [Internet]. 2021;34(4):316-324. Disponible en: <https://revistas.udea.edu.co/index.php/iatreia/article/view/342923>
23. Hoyos-Orrego A, Mantilla-Mantilla LF. Microbiology of bone-joint infections in children seen at a university clinic in the city of Medellín, Colombia. *Infectio* [Internet]. 2023;27(1):29-35. <https://doi.org/10.22354/24223794.1116>
24. Saavedra-Lozano J, Falup-Pecurariu O, Faust SN, Girschick H, Harwig N, Kaplan S, *et al.* Bone and joint infections. *Pediatr Infect Dis J* [Internet]. 2017;36(8):788-799. <https://doi.org/10.1097/INF.0000000000001635>
25. World Health Organization. International Classification of Diseases, Eleventh Revision (ICD-11) [Internet]. World Health Organization (WHO); 2019/2021. [Consulted 01 Jul 2022]. Available from: <https://www.who.int/classifications/classification-of-diseases>
26. OPS/OMS. Manual para el diagnóstico bacteriológico de la tuberculosis: Normas y guía técnica. Parte I Baciloscopia [Internet]. Organización Panamericana de la Salud; 2008. [Consulted 01 Jul 2022]. Disponible en: <https://iris.paho.org/handle/10665.2/782>
27. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing [Internet]. 32nd ed. CLSI supplement M100. Clinical and Laboratory Standards Institute; 2022. Available from: https://clsi.org/media/wi0pmpke/m100ed32_sample.pdf
28. Arnold JC, Bradley JS. Osteoarticular infections in children. *Infect Dis Clin North Am* [Internet]. 2015;29(3):557-574. <https://doi.org/10.1016/j.idc.2015.05.012>
29. Dormans JP, Drummond DS. Pediatric Hematogenous Osteomyelitis: New Trends in Presentation, Diagnosis, and Treatment. *J Am Acad Orthop Surg* [Internet]. 1994;2(6):333-341. <https://doi.org/10.5435/00124635-199411000-00005>
30. Pääkkönen M, Kallio MJT, Lankinen P, Peltola H, Kalillo PE. Preceding trauma in childhood hematogenous bone and joint infections. *J Pediatr Orthop B* [Internet]. 2014;23(5):196-199. <https://doi.org/10.1097/BPB.0000000000000006>
31. Camacho-Cruz J, Gutierrez IF, Brand-López K, Sosa-Rodriguez YA; Vásquez-Hoyos P, Gómez-Cortés LC, *et al.* Differences between methicillin-susceptible versus methicillin-resistant *Staphylococcus aureus* infections in pediatrics. Multicenter cohort study conducted in Bogotá, Colombia, 2014-2018. *Pediatr Infect Dis J* [Internet]. 2022;41:12-19. <https://doi.org/10.1097/INF.0000000000003349>
32. Lima AL, Oliveira PR, Carvalho VC, Cimerman S, Savio E. Directrices panamericanas para el tratamiento de las osteomyelitis e infecciones de tejidos blandos. Recommendations for the treatment of osteomyelitis. *Braz J Infect Dis* [Internet]. 2014;18(5):526-534. <https://doi.org/10.1016/j.bjid.2013.12.005>
33. Ratnayake K, Davis AJ, Brown L, Young TP. Pediatric acute osteomyelitis in the postvaccine, methicillin-resistant *Staphylococcus aureus* era. *Am J Emerg Med* [Internet]. 2015;33(10):1420-1424. <https://doi.org/10.1016/j.ajem.2015.07.011>

34. Burnett RSJ, Aggarwal A, Givens SA, McClure TJ, Morgan PM, Barrack RL. Prophylactic antibiotics do not affect cultures in the treatment of an infected TKA: A prospective trial. *Clin Orthop Relat Res* [Internet]. 2010;468(1):127-134. <https://doi.org/10.1007/s11999-009-1014-4>
35. Saavedra-Lozano J, Calvo C, Huguet R, Rodrigo C, Nuñez E, Pérez C, *et al.* Documento de consenso SEIP-SERPE-SEOP sobre etiopatogenia y diagnóstico de la osteomielitis aguda y artritis séptica no complicadas. *An Pediatr (Barc)* [Internet]. 2015;83(3):216.e1-216.e10. <https://doi.org/10.1016/j.anpedi.2014.08.006>