

Functional disability and morbidities among the elderly people, according to socio-demographic conditions and indicative of depression

Bárbara Ferreira Agreli¹
Flavia Aparecida Dias²
Pollyana Cristina dos Santos Ferreira³
Nayara Cândida Gomes⁴
Darlene Mara dos Santos Tavares⁵

Disability and morbidity among the elderly, according to sociodemographic conditions and indicative of depression

Objectives. To investigate the association of functional incapacity and the number of morbidities with sociodemographic, economic and health variables.

Methods. This is a household survey, quantitative and transversal approach, conducted with 1,693 elderly in Uberaba-MG in 2012. We used the Mini-Mental State Examination, Katz Scale, Lawton and Brody Scale; Geriatric Depression Scale abbreviated and semi-structured instrument for evaluation of demographic data, economic and morbidities.

Results. Functional incapacity in basic activities of daily living was associated with older age, a greater number of morbidities and presence of indicative of depression. Regarding the instrumental activities of daily living, functional incapacity was related to age, lower education and income, the greater number of morbidities and have indicative of depression. The largest number of comorbidities was associated with female sex, older age, lower income and indicative of depression. **Conclusion.** This study highlights the need for health actions directed to the oldest old, with less education and income, and indicative of depression to minimize the dependency on the performance of functional capacity and impact on morbidities.

1 Physiotherapy student, Federal University of Triângulo Mineiro -UFTM, Brazil. e-mail: bazinhaf@hotmail.com

2 Nurse, Master. PhD student, UFTM, Brazil. e-mail: flaviadias_ura@yahoo.com.br

3 Nurse, Master. PhD student, UFTM, Brazil. e-mail: pollycris21@bol.com.br

4 Nurse, Master. PhD student. UFTM, Brazil. e-mail: nayaranha_gomes@hotmail.com

5 Nurse, Ph.D. UFTM, Brazil. e-mail: darlenetavares@enfermagem.uftm.edu.br

Article from the research: Morbidity, quality of life and functional capacity of the elderly people.

Financing: Fundação de Amparo à Pesquisa do estado de Minas Gerais (FAPEMIG) e Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

Conflict of interests: none.

Receipt date: Aug 13, 2016.

Approval date: Jan 31, 2017.

How to cite this article: Agreli BF, Dias FA, Ferreira PCS, Gomes NC, Tavares DMS. Disability and morbidity among the elderly, according to sociodemographic conditions and indicative of depression. Invest. Educ. Enferm. 2017; 35(1): 48-58

DOI: 10.17533/udea.iee.v35n1a06

Descriptors: aged; activities of daily living; chronic disease; health care.

Discapacidad y morbilidad entre los ancianos, de acuerdo con las condiciones sociodemográficas e indicativo de la depresión

Objetivos. Investigar la asociación de la incapacidad funcional y el número de morbilidades con variables sociodemográficas, económicas y de salud entre los ancianos. **Métodos.** Se trata de una encuesta de hogares, de enfoque cuantitativo y transversal, realizado con 1 693 personas mayores en Uberaba-MG en 2012. Se utilizaron los instrumentos: Mini Examen del Estado Mental, Escala de Katz, Escala de Lawton y Brody y la Escala de Depresión Geriátrica Abreviada. También, se recolectó información sobre datos demográficos, económicos y de morbilidad. **Resultados.** La incapacidad funcional en actividades básicas de la vida diaria se asoció con la edad, el número de comorbilidades y la presencia de depresión. En cuanto a las actividades instrumentales de la vida diaria, la incapacidad funcional se relaciona con la edad, la educación y los ingresos más bajos, además del mayor número de comorbilidades y de probable depresión. El mayor número de comorbilidades se asoció con el sexo femenino, la edad avanzada, menores ingresos y probable depresión. **Conclusión.** Este estudio pone de relieve la necesidad de que las acciones de salud dirigidas a los más ancianos, con menos educación e ingresos, e indicativa de la depresión, con el fin de reducir al mínimo la dependencia de las prestaciones de la capacidad funcional y el impacto en la morbilidad.

Introduction

The evaluation of the functional capacity of the elderly allows the multi-professional team to have a more accurate view of the impact and severity of the diseases in the physical, emotional and social aspects.¹ Thus, studies on this subject can subsidize the planning of public health policies for the elderly according to their health priorities.² The prevalence of functional disability for basic activities (BADLs) and instrumental (IADLs) of daily life varies in

Descritores: anciano; actividades cotidianas; enfermedad crónica; atención a la salud.

Incapacidade e morbidades entre idosos, segundo condições sócio-demográficas e indicativo de depressão

Objetivo. Verificar a associação da incapacidade funcional e do número de morbidades com variáveis sócio-demográficas, econômicas e de saúde. **Métodos.** Trata-se de inquérito domiciliar, de abordagem quantitativa e transversal, realizado com 1 693 idosos em Uberaba-MG no ano de 2012. Utilizou-se o Mini Exame de Estado Mental, Escala de Katz, Escala de Lawton e Brody, Escala de Depressão Geriátrica Abreviada e instrumento semiestruturado para avaliação dos dados sócio-demográficos, econômicos e morbidades. **Resultados.** A incapacidade funcional para atividades básicas da vida diária esteve associada à maior faixa etária, maior número de morbidades e presença de indicativo de depressão. Em relação às atividades instrumentais de vida diária, a incapacidade funcional relacionou-se à maior idade, menor escolaridade e renda, maior número de morbidades e ter indicativo de depressão. O maior número de morbidades esteve associado ao sexo feminino, maior idade, menor renda e indicativo de depressão. **Conclusão.** Este estudo evidencia a necessidade de ações em saúde direcionadas aos idosos mais velhos, com menor escolaridade e renda e indicativo de depressão, visando minimizar a dependência para o desempenho da capacidade funcional e impacto sobre as morbidades.

Descritores: idoso; atividades cotidianas; doença crônica; atenção à saúde.

national studies³⁻⁵ and international studies.^{6,7} As for its determinant factors, research have controversies.⁸⁻¹⁰ In a study in Bahia, functional disability in IADLs was related to income, marital status, occupation during free time, age, health problems and sequels. In the BADLs there was only the presence of health problems.⁸ In another national study from Minas Gerais, dependence on IADL and BADL was associated with ≥ 75 years old, the absence of a partner, a stroke, a heart disease and diabetes mellitus.⁵ The indication of depression has been considered a risk factor for

functional incapacity in the national literature.⁹ In international research with elderly individuals 65 years old or older, female and older individuals were associated with functional disability for BADLs, leading to dependency. However, there was no such association for IADLs.¹⁰ In Japan, the elderly living with their children alone had a significantly greater risk of functional disability than the three-generation family group.⁷

It should also be pointed out that the elderly may present an unsuccessful aging, with associated comorbidities and dependence on activities of daily living (ADLs).⁹ Also, the literature has described that some socioeconomic conditions such as female, older and lower income are related to the greater number of morbidities.^{11,12} Brazilian research has shown that the low socioeconomic and cultural level are related to the development of chronic diseases.¹³ In an international study the female, with advanced age, worse self-perception of health and inability to access to medical care.¹⁴ However, studies in Brazil have been carried out in Family Health Strategies^{9,13} or with elderly people aged 65 and over,¹¹ and there is a need for further research among those aged 60 and over in the community.

The evaluation of the functional incapacity by the nurse together with the identification of the factors associated with the non-performance of the ADLs is important to promote early interventions, favoring the maintenance of autonomy and independence of the elderly. It is a tool for planning and implementing nursing care. It is understood that nursing assessments and consultations, as well as home visits, should include questions related to the changes in functionality.¹ Thus, this study aimed to verify the association between functional disability with gender, age, education, income, number of morbidities and the indicative of depression of the elderly living in the urban area to broaden the discussions on this topic, to identify the factors related to the presence of functional disability and greater number of morbidities, and to verify the association between the number of morbidities with the gender, the education, the age group, the income and the indicative of depression.

Methods

This study integrates a longitudinal study, developed by the Group of Research in Collective Health of the Federal University of the Triângulo Mineiro (UFTM) with a domiciliary survey type, and of quantitative, analytical, cross-sectional and observational approach. The sample was performed in a previous study, according to the totality of the elderly in the municipality, by the Research Center in Collective Health, considering 95% confidence, 80% of test power, margin of error of 4% for interval estimates and a proportion estimated from 0.5 to the proportions of interest, resulting in 2 892 elderly. A list was obtained to compose the sample, together with the Zoonoses Control Center of the municipality, containing the full name and address of the elderly. The choice of this place is because it maintains the updated register of all the households in Uberaba-MG. For the selection of the elderly, the stratified proportional sampling technique was used. The neighborhoods of the municipality were considered as strata. Thus, in 2012, the sample consisted of 2 118 eligible elders considering the losses in previous years. Inclusion criteria were: people aged 60 years old or older, living in the urban area of the city of Uberaba and without cognitive decline. Starting from the sample and observing the established criteria, 1,693 elderly people were interviewed, 265 died, and 160 presented a cognitive decline. Data collection was performed at the elderly household from July 2012 to March 2013 and reviewed by field supervisors.

The cognitive decline was evaluated by the Mini Mental State Examination (MMSE), translated and validated in Brazil.^{15,16} The socio-demographic data and morbidities self-reported by the elderly were collected in an instrument built Health Research Group. Functional capacity was performed by Index of Katz, adapted to the Brazilian reality¹⁷ and by the Lawton and Brody Scale, adapted in Brazil.¹⁸ The index of depression was measured by the Geriatric Abridged Depression Scale, validated in Brazil, with a cutoff of six or more.¹⁹ The following variables were selected: socio-demographic: gender (Male and female), age group in years (60-69, 70-79,

80 years old and over), marital status (married or living with a partner, separated/divorced; widowed and single), education in years of study (illiterate, 1-3, 4-7, 8, 9-10 and 11 or more), per capita income, in minimum wages (without income; <1; 1; >1-3; >3-5; >5); self-reported morbidities: rheumatism, arthritis/arthrosis, osteoporosis, asthma/bronchitis, tuberculosis, embolism, high blood pressure, poor circulation, heart problems, diabetes, obesity, stroke, Parkinson's disease, urinary incontinence, fecal incontinence, constipation, sleeping problems, cataracts, glaucoma, spinal problems, kidney problem, accident/trauma sequel, malignant tumors, benign tumors, vision problems; Number of morbidities: 0, 1-6, 7 or more; Functional capacity in BADLs: bathing, dressing, toilet, transfer, sphincter control, feeding; Number of functional disability for BADLs: (0, 1-3, 4 or more); Functional capacity in the IADLs: using the telephone, shopping, going to distant places, eating, housework, doing household chores, washing and ironing, taking medicines properly and taking care of finances; Classification of functional disability in IADLs: total dependence (0 to 7 points), partial dependence (8 to 20 points) and independent (21 points); Indicative of depression: yes or no.

The data collected were processed in a microcomputer, by two people, in double entry, in Excel® program. Subsequently, it was transported to the software "Statistical Package for Social Sciences" (SPSS) version 17.0, for data analysis. A descriptive statistical analysis was performed, and the chi-square test ($p < 0.05$) was used to compare the variables. The project was approved by the Committee on Ethics in Research with Human Beings of UFTM, opinion nº 2265. Only after the consent of the interviewee and signing the Free and Informed Consent Term, the interview was conducted.

Results

There was a female elderly prevalence, aged 70-79 years old, married, living with their children, with 4-7 years of education and income of a minimum wage. Table 1 shows the socioeconomic and demographic variables of the elderly.

Most of the elderly were independent for BADLs (96.2%). Among those with functional disability, 3.2% had 1-3 disabilities, and 0.6% had four and more, with those related to urinary and fecal control (2.7%) followed by bathing (1.4%) and dressing (1.4%). Independence for IADLs was reported by 33.8% of the elderly, 65.6% of whom needed partial help and 0.6% were dependent. The main disabilities were related to travel (25.5%) and housework (19%).

In the three groups, those with 70-79 years old prevailed. However, the proportion of elderly individuals aged 80 years old and over was higher among those with 1-3 and 4 and more disabilities. A higher number of morbidities and the presence of an indication of depression were associated with a greater number of functional disabilities for BADLs among the elderly.

Table 3 shows the level of dependence in the IADL according to socioeconomic variables, the number of morbidities and indicative of depression. In the IADLs, the highest level of dependence was higher among the elderly with 80 years old and over, lower educational level, who received a minimum wage, had seven or more morbidities and had an indicative of depression.

Table 4 presents the number of morbidities according to socioeconomic variables and indicative of depression. The majority of the elderly had 1-6 morbidities (65.5%) followed by seven or more (31.1%). The elderly, female, 70-79 years old, who received a minimum wage and had an indication of depression presented more morbidities compared to the others.

Table 1. Frequency distribution of the socioeconomic and demographic variables of the 1,693 elderly people. Uberaba-MG, 2014

Variables	Values	n	%
Gender	Male	616	36.4
	Female	1077	63.6
Age group	60-69 years old	645	38.1
	70-79 years old	737	43.5
	80 or more	311	18.4
Marital status	Never married/lived with partner	89	5.3
	Married	730	43.1
	Widow	682	40.3
	Separate/divorced	191	11.3
Education	None	346	20.4
	1-3 years	437	25.8
	4-7 years	604	35.7
	8 years	104	6.1
	9-10 years	33	1.9
	11 years and more	168	9.9
Per capita income	None	115	6.8
	< 1	30	1.8
	1	809	47.8
	>1-3	573	33.8
	>3-5	104	6.1
	>5	59	3.5
Home	Live alone	306	18.1
	Only with a Professional Caregiver	7	0.4
	Only with a partner	379	22.4
	With others of their generation (with or without a partner)	122	7.2
	With their children (with or without a partner)	688	40.6
	With their grandchildren (with or without a partner)	115	6.8
	Other	76	4.5

Table 2. Number of functional disability in BADL, according to socio-demographic and economic variables, number of morbidities and indicative of depression. Uberaba-MG, 2014

Variables	Number of disabilities for BADL						X ²	p
	Without (n = 1629)		1-3 (n = 54)		4 or more (n = 10)			
	n	%	n	%	n	%		
Gender								
Male	598	36.7	16	29.6	2	20	2.29	0.317
Female	1031	63.3	38	70.4	8	80		
Age group								
60-69 years old	628	38.6	16	29.6	1	10	11.44	0.022
70-79 years old	711	43.6	20	37.0	6	60		
80 years old or more	290	17.8	18	33.3	3	30		
Education								
None	332	20.4	12	22.2	2	20	10.04	0.437
1-3 years	417	25.6	17	31.5	3	30		
4-7 years	579	35.6	20	37.0	5	50		
8 years	101	6.2	3	5.6	0	0		
9-10 years	33	2.0	0	0	0	0		
11 years or more	166	10.2	2	3.7	0	0		
Per capita income								
None	107	6.6	7	13.0	1	10	17.41	0.066
< 1	30	1.8	0	0	0	0		
1	773	47.5	28	51.9	8	80		
>1-3	554	34.1	18	33.3	1	10		
>3-5	103	6.3	1	1.9	0	0		
>5	59	3.6	0	0	0	0		
Morbidities								
None	58	3.6	0	0	0	0	33.22	<0.001
1-6	1086	66.7	20	37.0	3	30		
7 or more	485	29.8	34	63.0	7	70		
Indicative of depression								
Yes	410	25.2	30	55.6	7	70	30.02	<0.001
No	1218	74.8	24	44.4	3	30		

Table 3. Functional dependence level in the IADL according to socio-demographic and economic variables, the number of morbidities and indicative of depression. Uberaba-MG, 2014

Variables	Level of dependence for IADL						X ²	p
	Total		Partial		Independent			
	n	%	n	%	n	%		
Gender								
Male	0	0	404	36.5	212	36.8	5.76	0.056
Female	10	100	703	63.5	364	63.2		
Age group								
60-69 years old	2	20	356	32.2	287	49.8	94.94	<0.001
70-79 years old	3	30	490	44.3	244	42.4		
80 years old or more	5	50	261	23.6	45	7.8		
Education								
None	2	20	268	24.2	76	13.2	44.32	<0.001
1-3 years	3	30	288	26.0	146	25.3		
4-7 years	4	40	384	34.7	216	37.5		
8 years	0	0	59	5.3	45	7.8		
9-10 years	0	0	15	1.4	18	3.1		
11 years or more	1	10	92	8.3	75	13.0		
Per capita income								
None	1	10	75	6.8	39	6.8	32.15	<0.001
< 1	0	0	22	2.0	8	1.4		
1	6	60	575	52.0	228	39.7		
>1-3	2	20	341	30.8	230	40.1		
>3-5	1	10	54	4.9	49	8.5		
>5	0	0	39	3.5	20	3.5		
Morbidities								
None	0	0	25	2.3	33	5.7	63.31	<0.001
1-6	3	30	679	61.3	427	74.1		
7 or more	7	70	403	36.4	116	20.1		
Indicative of depression								
Yes	6	60	346	31.3	95	16.5	51.94	<0.001
No	4	40	761	68.7	480	83.3		

Table 4. Number of morbidities according to socio-demographic and economic variables, the number of morbidities and indicative of depression. Uberaba-MG, 2014

Variables	Number of morbidities						X ²	p
	0		1-6		7 or more			
	n	%	n	%	n	%		
Gender								
Male	40	69.0	463	41.7	113	21.5	90.85	<0.001
Female	18	31.0	646	58.3	413	78.5		
Age group								
60-69 years old	31	53.4	434	39.1	180	34.2	10.14	0.038
70-79 years old	20	34.5	470	42.4	247	47.0		
80 years old or more	7	12.1	205	18.5	99	18.8		
Education								
None	12	20.7	219	19.8	115	21.9	10.99	0.358
1-3 years	11	19.0	288	26.0	138	26.2		
4-7 years	22	37.9	383	34.6	199	37.8		
8 years	4	6.9	77	6.9	23	4.4		
9-10 years	2	3.4	23	2.1	8	1.5		
11 years	7	12.1	118	10.6	43	8.2		
Per capita income								
None	7	12.1	74	6.7	34	6.5	21.25	0.019
< 1	0	0	22	2.0	8	1.5		
1	21	36.2	510	46.1	278	52.9		
>1-3	24	41.4	376	34.0	173	32.9		
>3-5	4	6.9	77	7.0	23	4.4		
>5	2	3.4	47	4.2	10	1.9		
Indicative of depression								
Yes	6	10.3	215	19.4	226	43.0	107.05	<0.001
No	52	89.7	893	80.5	300	57.0		

Discussion

Socio-demographic and economic data are consistent with those of the national and international scientific literature.^{1,10,20} As for the dependence on the BADL, similar results were obtained in researches carried out in Rio Grande do Sul¹ and João Pessoa²⁰ since the findings related to the IADL are divergent of the survey conducted in Goiânia, with 14.9% of the elderly totally dependent and 45.7% needing partial help.⁹ Dependence may require the need for support from caregivers who need adequate preparation and help. The health services must attend to the patients in their needs, being relevant to know which are the activities of greater dependence, to the elaboration of a plan of action directed to the promotion of health, of prevention and treatment of these commitments.⁹ It is emphasized that it has family homes in which the elderly live with younger generations, becoming increasingly common.²¹ In this context, considering that the largest percentage of elderly in this study were children, with or without a partner, it is necessary to develop multi-professional interventions with a focus on strengthening this support relationship.

The association of age, morbidities, and indicative of depression with a functional disability for BADL is similar to a national finding.⁹ Another survey in Belo Horizonte observed higher disability percentages with increasing age.⁴ Considering that functional incapacity among the elderly contributes to an increase in the use of acute care services, such as outpatient, emergency and hospitalization,²² and that the presence of depressive symptoms together with the social implications.⁹ it is relevant to identify early cases to elaborate actions that favor the maintenance of autonomy and independence of the elderly, even with limitations. The association between age and functional disability in IADL is also consistent with a national study.⁹ In Belo Horizonte, it was also observed higher percentages of disability as age increased.⁴ Regarding education, it is emphasized that the longer the study time, the greater the chances that the individual will remain autonomous despite his or her age.²

It should be noted that low income is related to a series of unfavorable conditions, which may contribute to the loss of functional autonomy. Low-income elderly individuals are more likely to have some dependency on the performance of IADL than those in the most economically disadvantaged classes.² The presence of poly-morbidities was also related to functional disability for IADL among the elderly in João Pessoa-PB.²⁰ Aging without planning may increase the risk of disease incidence. In this way, the prevention of comorbidities becomes paramount. From this perspective, public health policies should be directed towards the elderly. It is believed that, in this way, strategies that would increase the possibility of aging with the quality of life, improvement of autonomy and functionality may be constituted.²⁰ The evaluation of functional capacity can contribute in this aspect by enabling nurses and other members of the health team a better view of the severity and impact of comorbidities.¹ As in this study, the indicative of depression was associated with functional disability in national and international studies.¹² Thus, the results obtained in this research reinforce the health professional's need to track the indicative of depression and take the necessary actions, such as diagnostic confirmation and therapeutic establishment.

The association of the female with the greatest number of morbidities has also been reproduced in research carried out with the elderly in Brazil¹¹ and South Africa.¹² In this context, it is relevant that the nursing develops educational and preventive strategies especially focused on women. Considering their greater longevity and comorbidities, it is believed that the monitoring of their health conditions can minimize the impact on their functionality. Different from the results of this study, a survey carried out by the FIBRA Network did not observe differences between the age groups with the number of diseases,¹¹ and a study carried out in South Africa showed that the greatest number of morbidities prevailed in the age group of 60-70 years old.¹²

The association of low income with the greater number of morbidities was also verified in a survey conducted by the FIBRA Network.¹¹ It is

possible that the elderly with low income have less access to health services and medicines and have difficulties in maintaining treatment until the end interfering with their quality of life.^{11,20} Depression contributes to the elderly restriction to the domestic environment, which decreases their chances of enjoying the advantages of social interaction and it is associated with an increased risk of mortality, morbidity, physical and cognitive disability, inactivity and depression.²³ It is emphasized that the functional incapacity, among other factors, that compromise the social involvement of the elderly contribute to the elderly dissatisfaction with life²⁴ and it can justify this result. In this sense, nursing can contribute by identifying with the elderly social activities that are of interest considering their socioeconomic conditions.

Conclusion

The functional disability for the BADLs was associated with greater age, a greater number of comorbidities and the indicative of depression. The IADLs were the largest age group, lower education, lower income, presence of a greater number of diseases and indicative of depression. The elderly with the highest number of diseases were female, 70-80 years old, with a minimum wage and an indication of depression. These results reinforce that the social inequalities and health status of the elderly may be related to functional capacity and the presence of comorbidities. In this way, the importance of periodic evaluation of the functionality and aspects related to health is highlighted, considering the factors that can impact them to maintain the independence and control of the comorbidities of the elderly as long as possible. In this context, it is highlighted that the nurse can perform such evaluations in nursing consultations. Although it is not possible to identify the cause and effect relationship between the variables, since this is a cross-sectional study, the results found a point to the need for greater attention to maintaining the functionality and morbidities of this population. The identification of factors associated with the functional disability

of the elderly provides important elements for prevention and intervention measures. In this sense, it is suggested that new studies be carried out with a longitudinal design to verify the possible causal relationship between the variables.

References

1. Santos SSC, Gautério DP, Vidal DAS, Rosa BM, Zortea B, Urquia BS. (In)dependência na realização de atividades básicas da vida diária em pessoas idosas domiciliadas. *Rev. RENE*. 2013; 4(3):579-87.
2. Kagawa CA, Corrente JE. Análise da capacidade funcional em idosos do município de Avaré-SP: fatores associados. *Rev. Bras. Geriatr. Gerontol*. 2015; 18(3):577-86 .
3. Pereira GN, Bastos GAN, Duca GFD, Bós AJG. Indicadores demográficos e socioeconômicos associados à incapacidade funcional em idosos. *Cad. Saúde Pública*, Rio de Janeiro. 2012; 28(11):2035-42.
4. Fialho CB, Lima-Costa MF, Filho AIL. Capacidade funcional e uso de serviços de saúde por idosos da Região Metropolitana de Belo Horizonte, Minas Gerais, Brasil: um estudo de base populacional. *Cad. Saúde Pública*, Rio de Janeiro. 2014; 30(3):599-610.
5. Barbosa BR, Almeida JM, Barbosa MR, Rossi-Barbosa LAR. Avaliação da capacidade funcional dos idosos e fatores associados à incapacidade. *Cien. Saude Colet*. 2014; 19(8):3317-25.
6. Hu YN, Hu GC, Hsu CY, Hsieh SF, Li CC. Assessment of individual activities of daily living and its association with self-rated health in elderly people of Taiwan. *Int. J. Gerontol*. 2012; 6(2):117-21.
7. Saito E, Ueki S, Yasuda N, Yamazaki S, Yasumura S. Risk factors of functional disability among community-dwelling elderly people by household in Japan: a prospective cohort study. *BMC Geriatrics*. 2014; 14:93.
8. Reis LA, Reis LA, Torres GV. Impacto das variáveis sociodemográficas e de saúde na capacidade funcional de idosos de baixa renda. *Ciênc. Cuid. Saúde*. 2015;14(1):847-54.
9. Nunes DP, Nakatani AYK, Silveira EA, Bachion MM, Souza MR. Capacidade funcional, condições socioeconômicas e de saúde de idosos atendidos por equipes de Saúde da Família de Goiânia (GO, Brasil). *Cienc. Saude Colet*. 2010; 15(6): 2887-98.
10. Millán-Calenti JC, Tubío J, Pita-Fernández S, González-Abraldes I, Lorenzo T, Fernández-Arruty T et al. Prevalence of functional disability in activities

- of daily living (ADL), instrumental activities of daily living (IADL) and associated factors, as predictors of morbidity and mortality. *Arch. Gerontol. Geriatr.* 2010; 50(3):306–10.
11. Pinto JM, Neri AL. Doenças crônicas, capacidade funcional, envolvimento social e satisfação em idosos comunitários: Estudo Fibrá. *Cienc. Saude Colet.* 2013; 18(12):3449-60.
 12. Phaswana-Mafuya N, Peltzer K, Chirinda W, Musekiwa A, Kose Z, Hoosain E. et al. Self-reported prevalence of chronic non-communicable diseases and associated factors among older adults in South Africa. *Glob Health Action.* 2013; 6:20936.
 13. Pimenta FB, Pinho L, Silveira MF, Botelho ACC. Fatores associados a doenças crônicas em idosos atendidos pela Estratégia de Saúde da Família. *Ciência & Saúde Coletiva.* 2015; 20(8):2489-98.
 14. Jerliu N, Toçi E, Burazeri G, Ramadani N, Brand H. Prevalence and socioeconomic correlates of chronic morbidity among elderly people in Kosovo: a population-based survey. *BMC Geriatr.* 2013; 13:22.
 15. Bertolucci PHF, Brucki SMD, Campacci SR, Juliano Y. O mini-exame do estado mental em uma população geral: impacto da escolaridade. *Arq. Neuropsiquiatr.* 1994; 52(1):1-7.
 16. Brucki SMD, Nitrini R, Caramelli P, Bertolucci PHF, Okamoto IH. Sugestões para o uso do mini-exame do estado mental no Brasil. *Arq. Neuro-Psiquiatr.* 2003; 61(3):777-81.
 17. Lino VTS, Pereira SRM, Camacho LAB, Filho STR, Buksman S. Adaptação transcultural da Escala de Independência em Atividades de Vida Diária (Escala de Katz). *Cad. Saude Publica.* 2008; 24(1):103-12.
 18. Santos RL, Virtuoso Júnior JS. Confiabilidade da versão brasileira da escala de atividades instrumentais da vida diária. *Rev. Bras. Promoc. Saude.* 2008; 21(4):290-6.
 19. Almeida OP, Almeida SA. Confiabilidade da versão brasileira da Escala de Depressão em Geriatria (GDS) versão reduzida. *Arq. Neuropsiquiatr.* 1999; 57(2-B): 421-6.
 20. Dias LD, Brito GEG, Forte FDS, Araújo KMB, Lucena EMF. Perfil sociodemográfico e de saúde de idosos do município de João Pessoa – PB. *Rev. Bras. Promoc. Saude.* 2012; 25(1):86-96.
 21. Luchesi BM, Brito TRP, Costa RS, Pavarini SCI. Suporte social e contato intergeracional: estudando idosos com alterações cognitivas. *Revista Eletrônica de Enfermagem.* 2016; 17(3). Available from: <https://www.fen.ufg.br/revista/v17/n3/pdf/v17n3a06.pdf>
 22. Wu CY, Hu HY, Li CP, Fang YT, Huang N, Chou YJ. The association between functional disability and acute care utilization among the elderly in Taiwan. *Arch. Gerontol. Geriatr.* 2013; 57(2):177-83.
 23. Lee CT, Yeh CJ, Lee MC, Lin HS, Chen VC, Hsieh MH et al. Leisure activity, mobility limitation and stress as modifiable risk factors for depressive symptoms in the elderly: results of a national longitudinal study. *Arch. Gerontol. Geriatr.* 2012; 54(2):221-9.
 24. Ozturk A, Simşek TT, Yümin ET, Sertel M, Yümin M. The relationship between physical, functional capacity and quality of life (QoL) among elderly people with a chronic disease. *Arch. Gerontol. Geriatr.* 2011; 53(3):278-83.