

Table 1. Implant survival in the follow-up period

No. of patients		Evaluated implants		Unsuccessful Implants		Survival rate	
Time	n	n	n	n	%		
0 months	30	60	0	0	100		
12 months	29	57	1	1	98.1		
24 months	28	56	0	0	98.1		
36 months	27	54	0	0	98.1		

Table 2. Average radiographic bone loss in the follow-up period (in mm)

Mesial right implant			Distal right implant			Mesial left implant			Distal left implant		
Baseline	1 Year	3 years	Baseline	1 year	3 years	Baseline	1 year	3 years	Baseline	1 year	3 years
(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)	(CI 95%)
0.32	0.54	0.59	0.44	0.81	0.90	0.32	0.82	0.89	0.29	0.66	0.71
(0.14- 0.51)	(0.29- 0.79)	(0.35- 0.83)	(0.25- 0.62)	(0.51- 1.12)	(0.60- 1.20)	(0.14- 0.50)	(0.54- 1.10)	(0.6- 1.18)	(0.12- 0.46)	(0.41- 0.91)	(0.46- 0.96)

Table 3. Probing depth at 36 months. Frequencies are grouped by side and zone.

Probing depth	Right				Left			
	n	Mesial	n	Distal	n	Mesial	n	Distal
0 mm	3	11.5%	3	11.5%	3	11.5%	3	11.5%
1 mm	8	30.8%	5	19.2%	7	26.9%	5	19.2%
2 mm	7	26.9%	11	42.3%	9	34.6%	10	38.5%
3 mm	8	30.8%	7	26.9%	7	26.9%	7	26.9%

Table 4. Other conditions of the state of overdentures

Observations	No. of cases	%
Need to change retentive elements (rubbers)	5	19.2
Need to change retention attachments (housing)	4	15.4
Need to repair incisal edge fractures	3	11.5
Need to rebase the prosthetic base	2	7.7
Pigmentations	2	7.7
Presence of calculus on prostheses	2	7.7