

Table 1. Results per group. Comparison of time changes between T1 and T2 by Wilcoxon test

Control Group			
	T1	T2	P value
	Average ± SD	Average ± SD	> z
Right anterior space	1.89 ± 0.59	2.04 ± 0.47	0.21
Left anterior space	2.21 ± 0.79	2.06 ± 0.67	0.33
Right upper space	2.69 ± 0.83	2.63 ± 0.78	0.88
Left upper space	2.81 ± 0.70	2.84 ± 0.60	0.94
Right posterior space	2.32 ± 0.87	2.09 ± 0.69	0.57
Left posterior space	2.09 ± 1.05	2.03 ± 0.48	0.73
Bite Turbo Group			
	T1	T2	P value
			> z
Right anterior space	2.00 ± 0.50	2.21 ± 0.80	0.26
Left anterior space	1.70 ± 0.66	1.95 ± 0.60	0.15
Right upper space	2.29 ± 0.78	2.43 ± 0.94	0.26
Left upper space	1.75 ± 0.25	1.54 ± 0.22	0.03*
Right posterior space	2.08 ± 0.59	2.21 ± 0.62	0.67
Left posterior space	1.75 ± 0.19	1.81 ± 0.27	0.26

Source: by the authors

Table 2. Level of satisfaction with the use of bite turbo

Bite turbo group		
		%
Esthetic Compromise	No	100.0%
	Yes	0.0%
Use bite turbo again	No	14.3%
	Yes	85.7%
Oral hygiene difficulty	No	100.0%
	Yes	0.0%
Communication difficulty	No	85.7%
	Yes	14.3%
Masticatory difficulty	No	57.1%
	Yes	42.9%
Satisfaction score	Completely unsatisfactory	0.0%
	Unsatisfied	0.0%
	Indifferent	14.3%
	Satisfactory	42.9%
	Completely Satisfactory	42.9%

Source: by the authors