

Subject index for volume 26, 2013

- Animal slaughter; 26(3):159-168
- Animal welfare; 26(1):24-29
- Antimicrobial therapy; 26(2):137-143
- Aquaculture; 26(3):211-218
- Biofiltration; 26(3):211-218
- Bio-modeling; 26(3):159-168; 177-185
- Bos indicus* heifers; 26(1):9-14
- Bos taurus taurus*; 26(3):219-225
- Bos taurus*; 26(3):159-168; 169-176
- Bovine mastitis; 26(4):280-287
- Brody; 26(3):169-176
- Bubalus bubalis*; 26(3):177-185
- Buffering value index; 26(1):37-41
- Carcass conformation; 26(3):159-168
- Carcass quality; 26(3):159-168
- Cattle behavior; 26(1):24-29
- Cattle; 26(1):37-41; 193-200, 26(3):169-176, 26(4):255-262, 26(4):263-269
- Cellular immune response; 26(4):280-287
- Cellulose; 26(1):15-23
- Cervid; 26(2):137-143
- Chicken; 26(4):270-279
- Chloride cells; 26(2):127-135
- Coagulase negative staphylococcus; 26(2):119-126
- Conjugated linoleic acid; 26(4):263-269
- Conservation aquaculture; 26(2):127-135
- Cornea; 26(1):31-36
- Creole bulls; 26(2):98-107
- Dairy cattle; 26(2):90-99; 26(3):193-200
- Dairy steers; 26(3):169-176
- Days open; 26(3):193-200
- Digestibility; 26(2):69-78, 26(4):255-262
- Energy reserves; 26(4):288-299
- Environmental temperature; 26(3):193-200
- Euryhaline; 26(2):127-135
- Fat bulks; 26(4):288-299
- Fatty acid composition; 26(2):79-89
- Feed intake; 26(4):243-254
- Fertility; 26(3):193-200
- Fish growth; 26(3):211-218, 26(4):288-299
- Follicular growth; 26(1):9-14
- Forage; 26(1):15-23
- Genetic correlations; 26(3):177-185
- Genetic evaluation; 26(2):90-99
- Geometric morphometrics; 26(3):201-210
- Gompertz; 26(3):169-176
- Growth measurements; 26(3):159-168
- Growth rate; 26(3):169-176
- Gut immunology; 26(2):108-118
- Hemicellulose; 26(1):15-23
- Heritability; 26(2):90-99
- Histological; 26(1):15-23
- Horse; 26(1):31-36
- Hydroxycinnamic acid; 26(4):255-262
- Intestinal villi; 26(2):108-118
- Intracellular bacteria; 26(3):219-225

- Keratomalacia; 26(1):31-36
- Lactation; 26(3):177-185
- Lactic acid bacteria; 26(4):280-287
- Legendre polynomials; 26(3):177-185
- Leukocytes; 26(4):280-287
- Lignin; 26(1):15-23, 26(4):255-262
- Linseed oil; 26(2):79-89
- Lipids; 26(2):69-78
- Macrolides; 26(2):119-126
- Mammary gland; 26(4):280-287
- Mastitis; 26(2):119-126
- Mathematical models; 26(3):169-176, 26(4):270-279
- Meta-analysis; 26(4):243-254
- Metabolic adjustments; 26(4):288-299
- Milk composition; 26(4):263-269
- Milk; 26(2):69-78
- Modeling; 26(3):186-192
- Monounsaturated; 26(4):263-269
- Mycotoxin; 26(4):243-254
- Non-linear models; 26(3):169-176
- Nutrition; 26(1):37-41, 26(2):69-78
- Ophthalmology; 26(1):31-36
- Ovulation; 26(1):9-14
- Phenotypic plasticity; 26(3):201-210
- Phospholipids; 26(2):79-89
- Pig weaning; 26(2):108-118
- Pig; 26(4):243-254
- Post-traumatic abscess; 26(2):137-143
- Pregnancy; 26(1):9-14
- Production cruves; 26(4):270-279
- Random regression models; 26(3):177-185
- Real time PCR; 26(3):219-225
- Recombinant bovine somatotropin; 26(3):193-200
- Regression analysis; 26(4):270-279
- Reptile; 26(3):201-210
- Resident macrophages; 26(3):219-225
- Ruminant; 26(4):255-262
- Salinity tolerance; 26(2):127-135
- Seminal quality; 26(2):98-107
- Services per pregnancy; 26(3):193-200
- Sexual dimorphism; 26(3):201-210
- Single nucleotide polymorphisms, SNP's 26(4):263-269
- Sperm cryopreservation; 26(2):98-107
- Standardized ileal digestibility; 26(3):186-192
- Sunflower oil; 26(2):79-89
- Susceptible host; 26(3):219-225
- Sustainable development; 26(3):211-218
- Sustained swimming; 26(4):288-299
- Swine; 26(3):186-192
- Triglycerides; 26(2):79-89
- Tumor necrosis factor alpha (TNF- α); 26(2):108-118
- Vocal characteristics; 26(1):24-29
- von Bertalanffy; 26(3):169-176
- Water use; 26(3):211-218
- Weight gain; 26(4):243-254