

Ivermectin resistance of three *Rhipicephalus microplus* populations using the larval immersion test[□]

Resistencia de tres poblaciones de Rhipicephalus microplus a la ivermectina por el ensayo de inmersión de larvas

Resistência de três populações de Rhipicephalus microplus à ivermectina pelo teste de imersão de larvas

David Villar^{1*}, MV, MSc, PhD; Jonathan Puerta¹, Est MV; Anderson López¹, MV; Jenny J Chaparro¹, MV, MSc, DrSc.

¹Grupo de Investigación VERICEL, Laboratorio de Parasitología, Escuela de Medicina Veterinaria, Facultad de Ciencias Agrarias, Universidad de Antioquia (UdeA), Medellín, Colombia.

(Received: March 19, 2015; accepted: July 07, 2015)

doi: 10.17533/udea.rccp.v29n1a06

Summary

Background: in Colombia, the control of tick infestation in cattle is almost exclusively performed with chemical acaricides. It is important to determine the degree of resistance of *Rhipicephalus microplus* field populations to ivermectins (IVM) as the first step to design strategies to increase the useful life of acaricides and decrease the reliance on chemical controls. **Objective:** to test the degree of resistance to IVM. **Methods:** three cattle farms in Antioquia (Colombia) where IVM had previously failed to control infestations of *R. microplus* ticks were studied. Ticks were collected several months apart in 2013, and the larval immersion test (LIT) was performed on the progeny of the adult females. Concentration-mortality data were subjected to probit analysis. **Results:** the three populations showed lethal concentrations (LC) 50 and 99 of ≥ 30 ppm and ≥ 400 ppm, respectively. There was no difference observed in the LC at different times of collection for each population studied. Such high values for LC, together with very low regression slopes (≤ 2), indicated a very heterogeneous response to increasing concentrations of IVM, which is common for resistant populations. At two farms, subcutaneous injection with a long-acting formulation of IVM at a dosage of 630 $\mu\text{g}/\text{Kg}$ was ineffective at eliminating existing infestations and protecting against reinfestations, confirming farmer suspicions of lost efficacy. **Conclusion:** epidemiological studies are necessary to assess the current status of resistance to IVM in this region of Colombia, and it is likely that the intensive use of IVM will aggravate this situation in the future.

Keywords: low efficacy, macrocyclic lactones, ticks.

[□] To cite this article: Villar D, Puerta J, López A, Chaparro JJ. Ivermectin resistance of three *Rhipicephalus microplus* populations using the larval immersion test. Rev Colomb Cienc Pecu 2016; 29:51-57.

* Corresponding author: David Villar. Facultad de Ciencias Agrarias, Universidad de Antioquia, Calle 70 No. 52-21, Medellín, Colombia. E-mail: davidvillar2003@yahoo.com

Resumen

Antecedentes: en Colombia, el control de la infestación de garrapatas en el ganado se ha realizado casi exclusivamente con acaricidas químicos y es importante determinar el grado de resistencia de las poblaciones de campo de *Rhipicephalus microplus* a ivermectinas (IVM) como el primer paso para diseñar tácticas que aumenten su vida útil y disminuyan la dependencia de los controles químicos. **Objetivo:** probar el grado de resistencia a IVM. **Métodos:** se evaluó la resistencia de *R. microplus* a IVM en tres fincas ganaderas en Antioquia (Colombia) donde se sospechaba que productos con IVM estaban fracasando para controlar infestaciones por garrapatas. Se recolectaron garrapatas con varios meses de diferencia durante el año 2013 y se empleó la prueba de inmersión de larvas (LIT) con la descendencia de las hembras adultas. Los datos de concentración-mortalidad fueron sometidos a análisis probit. **Resultados:** los resultados para las tres poblaciones estudiadas mostraron concentraciones letales (LC) 50 y 99 de ≥ 30 ppm y ≥ 400 ppm, respectivamente. No se observó diferencia en las LC obtenidas en diferentes momentos de recogida para cada población estudiada. Los altos valores de LC, junto con pendientes muy bajas en las rectas de regresión (≤ 2), indicaron una respuesta muy heterogénea a concentraciones crecientes de IVM, propio de poblaciones resistentes. En dos de las granjas, la administración de inyecciones subcutáneas con una formulación de acción prolongada de la IVM a dosis de 630 $\mu\text{g}/\text{Kg}$ fue incapaz de eliminar las infestaciones y proteger frente a reinfestaciones, confirmando las sospechas de los ganaderos de pérdida de eficacia. **Conclusión:** estudios epidemiológicos son necesarios para evaluar el estado actual de la resistencia a IVM en esta región de Colombia, y es probable que el uso intensivo de IVM agrave esta situación en el futuro.

Palabras clave: garrapatas, lactonas macrocíclicas, pérdida de eficacia.

Resumo

Antecedentes: na Colômbia, o controle de infestação com carrapatos em bovinos tem sido realizado quase exclusivamente com acaricidas químicos pelo qual se faz importante para determinar o grau de resistência de populações de campo de *Rhipicephalus microplus* a uma das drogas antiparasitárias de maior uso no manejo de gado, a ivermectina (IVM), como o primeiro passo para projetar táticas para aumentar a sua vida útil e diminuir a dependência de controles químicos. **Objetivo:** testar o grau de resistência à ivermectina. **Método:** foi avaliada a resistência de *R. microplus* à ivermectina em três fazendas de gado leiteiro localizadas no departamento de Antioquia (Colômbia) com suspeita de que o controle dos carrapatos com a utilização deste produto não estava sendo efetivo. Coletaram-se carrapatos em diferentes meses durante o ano 2013, e foi utilizado o teste de imersão de larvas (LIT) na descendência das fêmeas adultas, como método para detecção da resistência. Os dados de concentração-mortalidade foram submetidos à análise Probit. **Resultados:** todas as três populações avaliadas mostraram concentrações letais (LC) 50 de ≥ 30 ppm, e LC 99 de ≥ 400 ppm. Não foram observadas diferenças na LC entre as coletas feitas nos diferentes meses, em cada população amostrada. Os altos valores de LC encontrados junto com uma baixa regressão linear (≤ 2), indicam uma resposta muito heterogênea as concentrações crescentes de ivermectina, próprio de populações resistentes. Em duas das fazendas, a administração de injeções subcutâneas de ivermectina com uma formulação de longa ação, em doses de 630 $\mu\text{g}/\text{Kg}$, não foram capazes de eliminar as infestações e proteger contra a reinfestação, confirmando as reclamações dos fazendeiros para a perda de eficácia da droga antiparasitária. **Conclusões:** os estudos epidemiológicos são necessários para avaliar o estado atual da resistência à ivermectina nesta região da Colômbia, e é provável que o uso intensivo da ivermectina agrave ainda mais esta situação no futuro.

Palavras chave: carrapatos, lactonas macrocíclicas, perda de eficácia.

Introduction

The cattle tick *Rhipicephalus (Boophilus) microplus* is considered the main external parasite affecting cattle productivity in the tropical countries of America (Grise *et al.*, 2014). In Colombia, control of this parasite is almost exclusively performed with chemical acaricides (García, 2011).

However, resistance to most acaricides is a growing phenomenon worldwide; therefore, it is necessary to identify strategies to increase their useful life and to decrease reliance on chemical controls. The first cases of ivermectin resistance were reported in Brazil 14 years ago (Martins and Furlong, 2001), but, as the compound was introduced in most countries in the early 1980s, it is likely that resistance developed

earlier. There are many types of generic ivermectin (IVM) in the Colombian market, with at least 29 different names registered. The farms analyzed here use IVM routinely, regardless of real or perceived needs.

Confirmation of resistance is routinely determined using dose-response bioassays and/or assessment of the field efficacy of acaricides according to methods approved by the FAO (FAO, 2004) or the World Association for the Advancement of Veterinary Parasitology (WAAVP; Holdsworth *et al.*, 2006). The standard *in vitro* bioassays recommended in the FAO guidelines are the larval packet test (LPT) and the adult immersion test (AIT). Although it remains difficult to predict field acaricide efficacy based on these resistance bioassays, they can diagnose resistance before control failures are obvious. The LPT is laborious and modifications such as the larval immersion test (LIT), initially developed in Brazil (Klafke *et al.*, 2006) is more sensitive in discriminating between resistant and susceptible strains against macrocyclic lactones (Sabatini *et al.*, 2001; Klafke *et al.*, 2012). Numerous reports using the LIT in the last decade show *R. microplus* resistance to ivermectin in Latin American countries, including Mexico (Perez-Cogollo *et al.*, 2010a; 2010), Uruguay (Castro-Janer *et al.*, 2011), and Brazil (Klafke *et al.*, 2010; 2012). To complement the above *in vitro* assays, field studies using the WAAVP guidelines should be used simultaneously to confirm a reduction in therapeutic efficacy and protective period (Holdsworth *et al.*, 2006).

The objective of this study was to determine the degree of resistance of *R. microplus* field populations to IVM at three cattle farms in the province of Antioquia (Colombia). This study complements a field study conducted at two farms in Antioquia and reported elsewhere (López *et al.*, 2015).

Materials and methods

The authors considered that this study did not require approval of an Ethics Committee for Animal Experimentation.

Type of study

A descriptive study was performed.

Selection of farms

Three Antioquian farms with failures to control tick infestation, that previously claimed a lack of satisfactory control despite of the use of one or more commercial trade versions of IVM, were visited on three separate occasions for the collection of ticks at approximately two-month intervals between March and October of 2013. On each occasion, the samples were taxonomically identified as *R. microplus*. Except for the Tarso farm, herds were typical of the predominant production system and management practices of their area. The tick from San Jerónimo was obtained at a family-run mixed dairy farm with approximately 30 Simmental, Normandy and Holstein crosses. The Yarumal ticks originated from a small dairy farm of approximately 32 cattle of Jersey and Holstein crosses. The Tarso ticks were collected from a beef farm of highly selected Brangus imported from Texas, and comprising approximately 250 cattle. IVM was used for an uncertain number of years at the three locations, and producers had failed to clear the initial invasion of engorging ticks after use of one or several trade versions of IVM.

Larval immersion test (LIT)

A control solution containing 1% ethanol and 0.02% Triton X-100 in distilled water was used to prepare all IVM immersion dilutions and to test control larvae. A commercial 1% IVM (Ivomec – Merial Saúde Animal, Brazil. Batch number BE314/11, expiration date 11/2016) was used to prepare all serial dilutions with the control solution. The analysis of this particular product reported a concentration of 0.85% ivermectin (López *et al.*, 2015). For the LIT, ten different concentrations of IVM dilutions were prepared, ranging from 10 to 300 ppm, in a final immersion volume of 4 mL; larvae arising from eggs of at least 25 ticks were first mixed in distilled water to avoid exposing only those originating from a few ticks. Approximately 300 - 400 larvae were then immersed in different concentration solutions for 10 minutes under gentle agitation. They were immediately placed in Petri dishes and were transferred with a paintbrush to filter papers (850 x 750 mm) that were folded and sealed with sticky tape, forming a packet.

Three packets containing approximately 100 larvae each were used for each IVM dilution. The packets were placed in the incubator for 24 h, after which they were opened under a desk lamp and the larvae were gently transferred with a paintbrush to a white sheet of paper. Mortality was immediately determined using a magnifying lens because the larvae are incapable of locomotion. To assist with counting, those that moved spontaneously were removed with sticky tape and considered alive. Bioassays for each farm were replicated at least three times.

Data analysis

The number of dead versus exposed larvae was introduced into IBM SPSS Statistics 21 software (version 2003) and submitted to probit analysis to calculate the LC50 and LC99% with their respective confidence intervals (CI95%), chi-square tests for homogeneity, and slopes of the regression line.

Results

Concentration-mortality regressions, lethal concentrations LC50 and LC99, their 95% confidence intervals and slopes for each population are shown in Table 1. There was no difference in the LC50 and LC99 estimates for each of the three populations studied when bioassays were performed at different times of collection. Mortality in the control groups exposed to the ethanol-Triton X solution was zero; therefore, no correction was necessary for the ivermectin-exposed groups. Figure 1 represents the

probit mortality plots with the regression lines for each of the populations studied. The small slopes of the regression (≤ 2.5) and the distant values for the LC50 and LC99 are indicative of a very heterogeneous response to increasing concentrations of IVM in all three populations.

Discussion

The objective of this study was to investigate farmer reports of failure to control tick infestation at three farms in Antioquia; in each case IVM formulations were no longer able to eliminate tick infestations from cattle. The evaluation of *R. microplus* resistance to IVM was performed using the LIT bioassay with larvae arising from ticks collected at the sampled farms, and further tested using *in vivo* studies that are reported elsewhere (López *et al.*, 2015). Although the LIT has been used as a successful method to detect resistance (Sabatini *et al.*, 2001; Klafke *et al.*, 2006), ideally, it should be run in parallel to a susceptible strain to calculate resistance ratios and determine discriminating doses. Despite the limitation of not having a susceptible strain in this study, the results showed LC and concentration-mortality slopes similar to those reported for the most resistant tick populations from Veracruz (Fernández-Salas *et al.*, 2012), Yucatán (Pérez-Cogollo *et al.*, 2010a; 2010b) and Sao Paulo (Klafke *et al.*, 2012). For example, comparison of LC50 values to those of a study that determined the status of resistance to IVM in 53 randomly selected field populations of *R. microplus* in Mexico (Fernández-Salas *et al.*, 2012) demonstrates that the level of resistance was equal to that of the four

Table 1. Lethal concentration (50 and 99%) estimates for ivermectin obtained using the larval immersion test (LIT) for field populations of *Rhipicephalus microplus* from several municipalities in the province of Antioquia, Colombia.

Population	Slope \pm SE	Chi ² (df)	LC50 (ppm)	CI 95% LC50	LC99 (ppm)	CI 95% LC99
Yarumal ^a	1.992 \pm 0.068	83.4 (8)	45.9	34.1-58.8	676.1	400.8-1580.0
Yarumal ^b	1.888 \pm 0.051	66.2 (10)	47.1	39.4-54.9	801.6	559.1-1297.8
San Jerónimo ^a	2.185 \pm 0.064	134.4 (11)	34.2	27.2-41.7	397.3	257.2-771.9
San Jerónimo ^b	2.341 \pm 0.076	90.1 (9)	36.1	27.8-45.2	356.2	233.2-681.7
Tarso ^a	1.501 \pm 0.065	113.9 (8)	27.8	15.8-40.0	985.5	418.7-6172.5
Tarso ^b	1.768 \pm 0.059	137.1 (11)	32.2	24.1-41.3	674.4	380.6-1742.7

df: degrees of freedom; LC: lethal concentration; ppm: parts per million; CI: confidence interval.

^{a, b}Data are from separate bioassays.




Figure 1. Concentration vs mortality plots from three field populations of *Rhipicephalus microplus* in Antioquia subjected to the larval immersion test with ivermectin.*

most resistant populations, with LC₅₀s in the 30-50 ppm range. In general, low slope values (1.2-1.9) are related to high LC₅₀ (>35 ppm) and high LC₉₉ (>600 ppm) values, as reported here, and are hallmarks of very high resistance compared to values for susceptible *R. microplus* strains (slopes \approx 5, LC₅₀s \approx 5 ppm for the Deutch strain). Low -slopes for the concentration-mortality lines could be interpreted as a result of a large heterogeneity with different levels of resistance in the field populations studied. To our knowledge, no studies of a highly resistant IVM strain have been described with high-slopes that would indicate complete loss

of heterogeneity and consequently of the remaining susceptible genes in a population.

To relate the results of the LIT bioassay to the field situation, a long-acting formulation of IVM was used on two of the sampled farms. That clearly showed that the efficacy of IVM in terms of clearing the initial infestation load and protecting against reinfestation was markedly reduced from target expectations (López *et al.*, 2015). At 10 days post-injection, a reduction in the number of adult standard ticks was only 40 and 75% at the San Jeronimo and

* This figure appears in full color in the online version of this article.

Tarso farms, respectively (López *et al.*, 2015). These results are worrisome compared to field studies showing that 3.15% ivermectin formulations had a therapeutic and persistent efficacy of 95% at 56 days post-treatment (Arieta-Román *et al.*, 2010). In a study conducted at an USDA-ARS quarantine facility, the therapeutic efficacy of a similar long-acting ivermectin formulation was 99.9% against all stages of a susceptible strain of *R. microplus* at the time of treatment (Davey *et al.*, 2010). Furthermore, the protective period against larval re-infestation was 14 d if a level of control $\geq 99\%$ was desired, and dropped to 70.4% for animals exposed artificially to larvae at 28 d post-treatment. Because ticks are unable to complete development in <18 d, the study concluded that cattle could be treated with this long-acting IVM at 31 d intervals without the risk of viable ticks detaching from the infested animals. In the present study, the high degree of resistance based on the LIT was clearly demonstrated by the ineffective elimination of existing infestations and the loss of protection against reinfestations.

In conclusion, the present study confirmed the presence of *R. microplus* ticks that are highly resistant to IVM to the point of having lost efficacy for eliminating existing tick infestations. The sampled farms completely relied on chemical control to fight tick infestations. Therefore, it is imperative to apply integrated approaches for parasite control in farms that now have multiresistant strains of *R. microplus*.

Acknowledgements

The authors thank the Biogénesis Research group Sustainability Project 2014-2015 (Estrategia de Sostenibilidad CODI 2014-2015).

Conflicts of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

References

- Arieta Román RJ, Rodríguez Vivas RI, Rosado Aguilar JA, Ramirez Cruz GT, Basto Estrella G. Persistent efficacy of two macrocyclic lactones against natural *Rhipicephalus (Boophilus) microplus* infestations in cattle in the Mexican tropics. *Rev Mex Cienc Pecu* 2010; 1: 59-67.
- Castro Janer E, Rifran L, Gonzalez P, Niell C, Piaggio J, Gil A, Schumaker TTS. Determination of the susceptibility of *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae) to ivermectin and fipronil by larval immersion test (LIT) in Uruguay. *Vet Parasitol* 2011; 178:148-55.
- Davey RB, Pound JM, Miller JA, Klavons JA. Therapeutic and persistent efficacy of a long-acting (LA) formulation of ivermectin against *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae) and será concentration through time in treated cattle. *Vet Parasitol* 2010; 169:149-56.
- FAO. Working Group on Parasite Resistance. "Acaricide resistance: diagnosis, management and prevention", in Guidelines Resistance Management and Integrated Parasite Control in Ruminants 2004, pp. 25-77.
- Fernández Salas A, Rodríguez Vivas RI, Alonso Diaz MA, Basurto Camberos H. Ivermectin resistance status and factors associated in *Rhipicephalus microplus* (Acari:Ixodidae) populations from Veracruz, Mexico. *Vet Parasitol* 2012; 190:210-5.
- García Paz JL. Evaluación de las propiedades acaricidas de *Piper crassinervium* Kunth. *Piper aequale* Vahl. (*Piperaceae*) sobre larvas de *Rhipicephalus (Boophilus) microplus*. Trabajo de grado para optar a título de maestría. Universidad Nacional. Colombia. 2011
- Grise L, Leite RC, Martins JR, Barros AT, Andreotti R, Perez de Leon AA, Pereira JC, Silva H. Reassessment of the potential economic impact of cattle parasites in Brazil. *Braz J Vet Parasitol Jaboticabal* 2014; 23(2):150-6.
- Holdsworth PA, Kemp D, Green P, Peter RJ, De Bruin C, Jonsson NN, Letonja T, Rehbein S, Vercruyse J. World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) guidelines for evaluating the efficacy of acaricides against ticks (Ixodidae) on ruminants. *Vet Parasitol* 2006; 136:29-43.
- Klafke GM, Sabatini GA, Albuquerque TA, Martins JR, Kemp DH, Miller RJ, Schumaker TTS. Larval immersion tests with ivermectin in populations of the cattle tick *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae) from State of Sao Paulo, Brazil. *Vet Parasitol* 2006; 142:386-90.
- Klafke GM, Albuquerque TA, Miller RJ, Schumaker TTS. Selection of an ivermectin-resistant strain of *Rhipicephalus microplus* (Acari: Ixodidae) in Brazil. *Vet Parasitol* 2010; 168:97-104.

Klafke GM, Castro Janer E, Mendes MC, Namindome A, Schumaker TTS. Applicability of *in vitro* bioassays for the diagnosis of ivermectin resistance in *Rhipicephalus microplus* (Acari: Ixodidae). *Vet Parasitol* 2012; 184:121-220.

López Arias A, Villar D, Miller R, Chaparro J, Pérez de León, A. Reduced efficacy of commercial acaricides against populations of resistant cattle tick *Rhipicephalus microplus* from two municipalities of Antioquia, Colombia. *Libertas Académica*, 2015 (in press).

Martins JR, Furlong J. Ivermectin resistance of the cattle tick *Boophilus microplus* in Brazil. *Vet Rec* 2001; 149:64.

Pérez Cogollo LC, Rodríguez Vivas RI, Ramírez Cruz GT, Rosado-Aguilar JA. Survey of *Rhipicephalus microplus* resistance to ivermectin at cattle farms with history of macrocyclic lactones use in Yucatan, Mexico. *Vet Parasitol* 2010a; 172:109-113.

Pérez Cogollo LC, Rodríguez Vivas RI, Ramírez Cruz GT, Miller RJ. First report of the cattle tick *Rhipicephalus microplus* resistant to ivermectin in Mexico. *Vet Parasitol* 2010b; 168:165-9.

Sabatini GA, Kemp DH, Hughes S, Nari A, Hansen J. Tests to determine LC50 and discriminating doses for macrocyclic lactones against the cattle tick, *Boophilus microplus*. *Vet Parasitol* 2001; 95(1):53-62.