

Comparison of laboratory methods to assess fiber contents in feedstuffs[□]

Comparación de métodos de laboratorio para determinar el contenido de fibra en alimentos para animales

Comparação de metodologias laboratoriais para determinação de fibras em alimentos

Maria do Socorro N Lourenço¹, PhD; Juliana D Messana^{2*}, PhD; Ana Paula de O Sader², PhD; Roberta C Canesin³, PhD; Euclides B Malheiros⁴, Pablo S Castagnino², MSc; Telma T Berchielli^{2,5}, PhD.

¹*Departamento de Química e Biologia, Universidade Estadual do Maranhão, São Luís, MA, Brasil.*

²*Departamento de Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista, Jaboticabal, SP, Brasil.*

³*Instituto de Zootecnia, Centro de Pesquisas em Pecuária de Corte, Sertãozinho, São Paulo, Brasil.*

⁴*Departamento de Ciências Exatas, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista, Jaboticabal, SP, Brasil.*

⁵*Membro INCT/CA, Universidade Federal de Viçosa – Departamento de Zootecnia, Viçosa, MG, Brasil.*

(Received: September 12, 2014; accepted: September 13, 2016)

doi: 10.17533/udea.rccp.v30n1a03

Abstract

Background: Peter Van Soest proposed the conventional method of fiber analysis using neutral detergent fiber (NDF) and acid detergent fiber (ADF). The main advantage of this method is the precision of results; however, reagents are costly and laboratory work requires long runtime and labor. **Objective:** to compare analytical data from different methodologies used to assess NDF and ADF contents. **Methods:** means obtained with the conventional method were compared through Dunnett's test ($\alpha = 5\%$) with values from alternative methods using autoclave as the digester system. A completely randomized design in a 4 x 2 factorial arrangement was used. **Results:** NDF content through alternative methods was assessed for Tifton 85 hay, babassu meal and sugarcane, whereas ADF was only determined for babassu meal. NDF and ADF did not differ ($p \geq 0.05$) between non-sequential or sequential analysis for all the feedstuffs and methods studied, except for ADF in

□ To cite this article: Lourenço MS, Messana JD, Sader AP, Canesin RC, Malheiros EB, Castagnino PS, Berchielli TT. Comparison of laboratory methods to assess fiber contents in feedstuffs. Rev Colomb Cienc Pecu 2017; 30:21-29.

* Corresponding author: Juliana D Messana. Departamento de Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista. Via de Acesso Prof Paulo Donato Castellane,s/n, CEP:14884-900, Jaboticabal, SP, Brasil. Tel.:+551632092682. E-mail: duarte_juliana@hotmail.com

corn silage. **Conclusion:** alternative methodologies allow reducing operating costs and time but lack uniformity and accuracy for analyzing ADF in corn silage.

Keywords: acid detergent fiber, babassu meal, neutral detergent fiber, sugarcane, Tifton 85 hay.

Resumen

Antecedentes: Peter Van Soest propuso el método tradicional de análisis de los valores de fibra detergente neutra (FDN) y fibra detergente ácida (FDA). La principal ventaja de este método es la precisión de los resultados, sin embargo, los reactivos son costosos y el trabajo de laboratorio es largo y dispendioso. **Objetivo:** comparar diferentes metodologías propuestas para el análisis de las FDN y FDA mediante la medición de los datos analíticos. **Métodos:** las medias obtenidas por el método tradicional fueron comparadas con las medias de métodos alternativos que usan autoclave como sistema digestor por el test de Dunnette ($\alpha = 5\%$). Un diseño completamente aleatorio, con arreglo factorial 4×2 fue adoptado. **Resultados:** los métodos alternativos fueron utilizados para las determinaciones de FDN en el heno de Tifton 85, salvado de harina de babasú y caña de azúcar, mientras que las determinaciones de los valores de FDA apenas en el salvado de harina de babasú. Los resultados de los análisis de FDN y FDA no fueron significativamente diferentes ($p \geq 0.05$) entre los análisis no secuencial y secuencial en todos los alimentos y métodos estudiados, a excepción de la determinación de FDA en el ensilaje de maíz. **Conclusión:** las metodologías alternativas permitieron la reducción de costos operacionales y del tiempo de análisis, pero sin uniformidad en la precisión en el análisis de FDA en el ensilado de maíz.

Palabras clave: caña de azúcar, fibra detergente ácida, fibra detergente neutra, harina de babasú, heno de Tifton 85.

Resumo

Antecedentes: Peter Van Soest propôs o método tradicional de análise dos valores de fibra em detergente neutro (FDN) e fibra em detergente ácido (FDA). A principal vantagem deste método é a precisão dos resultados, no entanto, são caros reagentes e trabalho de laboratório longos e caros. **Objetivo:** comparar diferentes metodologias propostas para análise da FDN e FDA, através da verificação dos dados analíticos. **Métodos:** as médias obtidas pelo método convencional foram comparadas com as médias de métodos alternativos que utilizam a autoclave como sistema digestor, pelo teste de Dunnette ($\alpha = 5\%$). O delineamento adotado foi o inteiramente casualizado, em um esquema fatorial 4×2 . **Resultados:** os métodos alternativos foram usados para determinação de FDN no feno de Tifton 85, farelo de babaçu e cana-de-açúcar, enquanto para as determinações dos teores de FDA, apenas no farelo de babaçu. A precisão observada nas análises dos teores de FDN e FDA não diferiram significativamente ($p \geq 0.05$) quanto a análise não sequencial e sequencial, em todos os alimentos e métodos estudados, com exceção da determinação de FDA na silagem de milho. **Conclusão:** as metodologias alternativas permitiram a redução dos custos operacionais e do tempo da análise, mas sem uniformidade na precisão das análises de FDA em silagem de milho.

Palavras chave: cana-de-açúcar, farelo de babaçu, feno de Tifton 85, fibra detergente ácida, fibra detergente neutra.

Introduction

There is an increase challenge to develop efficient and low-cost analytical methodologies (Bialowas *et al.*, 2006; Gerbase *et al.*, 2006). To make sure that an analytical method generates reliable information from a sample, it must be validated through a process starting from the planning of the analytical strategy and goes on through its practical development (Inmetro, 2007). Analytical determination of neutral detergent fiber (NDF) and acid detergent fiber

(ADF) contents are important variables analyzed in a ruminant laboratory. The conventional method by Van Soest has a large feedstuffs database with precise results (Jung, 1997; Silva and Queiróz, 2002). However, reagents are costly and laboratory routine requires long runtimes due to manual steps. In an attempt to optimize this methodology, several alternative methods have been adopted in laboratory routines, but often without ensuring reliability of the results, which requires a systematic evaluation of the analytical procedure to demonstrate its precision

and accuracy. The use of autoclaves instead of the conventional digester is an alternative method recommended for NDF and ADF analyses (Pell and Schofield, 1993; Deschamps, 1999; Senger *et al.*, 2008). Through this system, it is possible to analyze the samples collectively, without requiring individual control of the samples. In this system, samples can be weighed both in filter crucibles or small bags, and the analyses can be conducted in a sequential or non-sequential form (Komarek, 1993). Some authors have suggested that fiber analyses should be performed sequentially on high-pectin forages (Van Soest *et al.*, 1991). Additionally, TNT (non-woven textile) bags have been preferentially used to reduce the costs of analysis (Casali *et al.*, 2009). The aim of this study was to evaluate different methodologies for NDF and ADF quantification by assessing the analytical data.

Materials and methods

The experiment was conducted at the Laboratory of Animal Nutrition of UNESP, Campus Jaboticabal—SP, Brazil. Six feedstuffs were evaluated: five roughages—Tifton 85 hay (*Cynodon* spp.), sugarcane (*Saccharum officinarum* L.), corn silage (*Zea mays* L.), xaraes grass (*Brachiaria brizantha* cv. *Xaraés*) and marandu grass (*Brachiaria brizantha* cv. *Marandu*)—and one protein concentrate—babassu meal (*Orbignya phalerata*). Except for babassu meal and Tifton 85 hay, all samples were pre-dried, following the procedure described by Silva and Queiroz (2002).

The NDF and ADF solutions were prepared following the methodology proposed by Van Soest *et al.* (1991). Decalin and sodium sulfite were not used. For measuring amylase-treated NDF of corn silage, 50 µL/g of DM of term stable alpha-amylase were used (Novozymes, Araucária, PR, Brazil). In each analytical method, 15 L of each detergent solution were prepared separately and at once, aiming at the precision of results and minimizing possible errors during quantitative determinations. Acetone and deionized water were used as solvents for washing the samples. Repeatability was calculated to verify precision of the analyses, representing the concordance between results from consecutive measurements of the same method performed under the same measuring

conditions (repeatability conditions): same procedure, same analyst, same instrument used under the same conditions and place (Inmetro, 2000).

The analyses developed by the conventional methodology followed the method modified by Van Soest *et al.* (1991). In each determination, around half gram of sample was weighed, adding 100 mL of detergent solution (acid or neutral) in every digestion step, and lead to boil for one h. The analytical results were obtained considering sample weight—being a gravimetric quantitative determination—through the following formula (1) for the determination of the NDF or ADF contents:

$$\% \text{ NDF or ADF} = (\text{WF-T}) / \text{WS} \times 100 \quad (1)$$

Where:

WS = dry matter weight of the sample in grams.

WF = weight (g) of the crucible plus detergent fiber residue after digestion and drying.

T = tare (initial weight) of the crucible (g).

The three alternative methodologies using autoclave (Pell and Schofield, 1993; Deschamps, 1999; Senger *et al.*, 2008) are simplifications of the original analytical procedures, without altering the principles of the method proposed by Van Soest (1963; 1967). The alternative methods differed as to the material used for conditioning the samples during analysis. For the purpose of organization, the alternative methods were named as follows:

Alternative method 1 - autoclave/ANKOM bags.

Alternative method 2 - autoclave/TNT (non-woven textile) bags.

Alternative method 3 - autoclave/filter crucibles.

Alternative methods 1 and 2, using small bags for sample conditioning and autoclave as digester, followed the recommendations by Komarek (1993) as to use bags instead of filter crucibles, and by Pell and Schofield (1993), Deschamps (1999), and Senger *et al.* (2008), concerning the use of autoclave. The time and temperature in the autoclave followed the best result proposed by Senger *et al.* (2008).

The ANKOM bags were acquired ready for use, whereas the non-woven textile (TNT) bags were produced manually, using 100-micron (μ) TNT and a mold with the dimensions of the ANKOM bags (5 x 5 cm).

The digestion step in alternative methods 1 and 2 was done in autoclave for 40 min, at a temperature of 110 °C (Senger *et al.*, 2008). In this stage all sealed bags containing the samples were conditioned in a single plastic beaker (capacity of 2 L) and immersed in 600 mL detergent solution (neutral or acid). All bags were washed in a beaker (2 L capacity) with hot (90–100 °C) water three times (5 min each). Bag residues were soaked two times in acetone for 5 min. All bags were collectively washed. Concentration of NDF and ADF was calculated using the following formula (2):

$$\% \text{ NDF or ADF} = (\text{WF-T}) / \text{WS} \times 100 \quad (2)$$

Where:

WS = dry matter weight of the sample (g).

WE = weight (g) of the bag plus residue of detergent fiber after digestion and drying.

T = tare (initial weight) of the bag (g).

B = blank value (grams; final weight of the bag after drying/initial weight of the bag).

The alternative method 3 used filter crucibles for sample conditioning (Van Soest *et al.*, 1991). Around half gram of each sample was weighed in the filter crucibles in triplicate, which were coupled to individual plastic beakers, adding 600 mL detergent solution (neutral or acid), inside the autoclave containing water for the digestion process. Digestion occurred in 40 min. Then, crucibles were immediately washed. Washing, drying and weighing the crucibles with the residue followed the same procedure as the conventional method. NDF and ADF contents were calculated as the difference between the tare of the crucible and the crucible weight plus detergent fiber residue after digestion and drying, using the formula defined in (1).

The analyzes performed by the conventional or alternative methods were conducted considering two different laboratory sequences: the first, known as non-sequential order, involves two weightings of the same sample and proceeding to the NDF and ADF analysis separately; while in the second sequence, known as sequential, a single sample is weighed to determine NDF. Then, we used the NDF residue to determine ADF content by washing, filtering and oven-drying.

Statistical analysis

The design was completely randomized, in a 4 x 2 factorial arrangement (4 methodologies and 2 sequences of analysis). The statistical model was:

$$Y_{ijk} = m + MT_i + FM_j + (MT*FM)_{ij} + \varepsilon_{ijk}$$

Where:

Y_{ijk} = NDF and ADF contents.

m = overall effect of the mean.

MT_i = effect of method i.

FM_j = effect of sequence j.

$(MT*FM)_{ij}$ = effect of the interaction between method i and sequence j.

ε_{ijk} = residual error.

The normality test of error used was Cramer-von-miser's ($\alpha = 5\%$), and the homoscedasticity test was Levene's ($\alpha = 5\%$). The data were subjected to analysis of variance through General Linear Models (GLM) of the SAS, version 9.1[®] (Statistical Analysis System Institute, Cary, NC, USA, 2002). Means were compared by Dunnett's test, having the conventional method as "control" (CMT, $\alpha = 5\%$).

Results

After obtaining the analytical results, the means by each alternative method were compared with those obtained by the conventional method.

There was significant difference between methods for all the studied feedstuffs (Table 1; $p < 0.01$). The sequence of analytical procedure (non-sequential or sequential) did not differ ($p \geq 0.05$). No significant interaction was observed ($p \geq 0.05$) between method and sequence of analysis for all feedstuffs (Table 1). Variation observed in NDF content in hay was 78.84 (alternative method 1) and 81.82% (alternative method 3). NDF content in sugarcane varied from 49.73 (alternative method 2) to 53.68% (alternative method 3). NDF in corn silage was different from

the conventional method (NDF = 42.9%; $p = 0.01$) compared to alternative method 1 (NDF = 55.11%), alternative method 2 (NDF = 48.21%), and alternative method 3 (NDF = 49.54%).

ADF content differed among methods in all feedstuffs studied ($p < 0.01$; Table 2), but did not differ in the analytical procedure (non-sequential or sequential) for all feedstuffs ($p < 0.01$). As for the interaction between method and sequence of analysis only corn silage was different ($p < 0.01$).

Table 1. Means observed in the analysis of variance in feedstuffs, sequence (SE) and methods studied for evaluating NDF contents.

Feedstuff	Methods ¹					Mean ²	CV ⁴ (%)	P-value
	SE	CMT	AMT ₁	AMT ₂	AMT ₃			
Tifton hay	NS	78.90	78.76	79.72	81.99	79.73 ^a	2.01	0.01
	S	79.06	78.91	78.51	81.63	79.53 ^a		
	Mean ³	78.98	78.84	79.11	81.82 [*]			
Sugarcane	NS	54.08	50.22	49.96	53.42	51.92 ^a	2.83	0.01
	S	52.89	50.24	49.50	53.94	51.64 ^a		
	Mean ³	53.48	50.23 [*]	49.73 [*]	53.68			
Corn silage	NS	41.72	55.69	48.39	50.08	48.97 ^a	7.64	0.01
	S	44.07	54.52	48.03	49.01	48.91 ^a		
	Mean ³	42.90	55.11 [*]	48.21 [*]	49.54 [*]			
Babassu meal	NS	70.94	62.81	67.77	65.07	66.73 ^a	5.92	0.01
	S	65.84	64.96	67.07	70.31	67.04 ^a		
	Mean ³	68.39	63.88 [*]	67.42	67.99			
Xaraes grass	NS	68.96	72.28	70.20	71.48	70.69 ^a	2.49	0.01
	S	68.33	71.34	69.75	71.30	70.18 ^a		
	Mean ³	68.64	71.81 [*]	69.98 [*]	71.39 [*]			
Marandu grass	NS	72.44	77.02	74.49	75.79	74.91 ^a	2.34	0.05
	S	72.57	75.21	74.21	75.97	74.48 ^a		
	Mean ³	72.51	76.15 [*]	74.35 [*]	75.89 [*]			

¹SE = sequence (NS = non-sequential; S = sequential); CMT = conventional method (block digester/filter crucibles); AMT₁ = alternative method 1 (autoclave/ANKOM); AMT₂ = alternative method 2 (autoclave/TNT); AMT₃ = alternative method 3 (autoclave/filter crucibles).

²Means in column of sequence (NS and S) followed by the same letter do not differ by the F test ($\alpha = 0.05$).

³Means in rows of methods AMT₁, AMT₂, and AMT₃ followed by asterisks (*) differ from CMT by Dunnett's test ($\alpha = 0.05$).

⁴Coefficient of variation.

The means of ADF content obtained in the analyses of the six feedstuffs studied were calculated, considering the analytical methods and sequences (Table 2).

The same analytical pattern was observed in ADF determinations for Tifton hay, sugarcane,

xaraes grass, and marandu grass. All alternative methods differed ($p < 0.05$) from the conventional method, especially alternative method 2, which had greater mean compared to the other methods. Alternative method 1 was recommended for determining ADF contents in babassu meal, with no loss of analytical precision.

Table 2. Means obtained in the analysis of variance of feedstuffs, sequences (SE) and methods studied at the evaluation of ADF contents.

Feedstuff	Methods ¹					Mean ²	CV ⁴ (%)	P-value
	SE	CMT	AMT ₁	AMT ₂	AMT ₃			
Tifton hay	NS	38.78	47.58	63.10	47.21	49.17 ^a	8.65	0.05
	S	41.32	43.64	67.12	44.32	48.80 ^a		
	Mean ³	40.05	45.61 [*]	65.11 [*]	45.77 [*]			
Sugarcane	NS	30.67	35.39	42.61	34.73	46.08 ^a	9.23	<0.01
	S	31.38	32.23	43.19	34.15	46.03 ^a		
	Mean ³	31.02	33.81 [*]	42.91 [*]	34.44 [*]			
Corn silage	NS	26.27 ^a	32.10 ^{*a}	35.87 ^{tb}	31.54 ^{*a}	31.20	8.56	<0.01
	S	25.94 ^a	28.46 ^{tb}	39.65 ^{*a}	30.07 ^{*a}	31.03		
	Mean ³	26.11	30.28	37.97	30.78			
Babassu meal	NS	37.82	43.66	56.83	46.01	46.08 ^a	11.75	<0.01
	S	42.78	40.99	56.84	43.52	46.03 ^a		
	Mean ³	40.30	42.33	56.84 [*]	44.77 [*]			
Xaraes grass	NS	36.74	46.83	57.19	42.68	45.86 ^a	10.21	<0.01
	S	37.12	41.28	58.10	39.84	44.09 ^a		
	Mean ³	36.93	44.06 [*]	57.65 [*]	41.26 [*]			
Marandu grass	NS	38.42	49.12	60.44	44.66	48.16 ^a	11.42	0.05
	S	39.71	43.72	61.50	41.82	46.69 ^a		
	Mean ³	39.06	46.42 [*]	60.97 [*]	43.24 [*]			

¹SE = sequence (NS = non-sequential; S = sequential); CMT = conventional method (block digester/filter crucibles); AMT₁ = alternative method 1 (autoclave/ANKOM); AMT₂ = alternative method 2 (autoclave/TNT); AMT₃ = alternative method 3 (autoclave/filter crucibles).

²Means in column of sequence (NS and S) followed by the same letter do not differ by the F test ($\alpha = 0.05$).

³Means in rows of methods AMT₁, AMT₂, and AMT₃ followed by asterisks (*) differ from CMT by Dunnett's test ($\alpha = 0.05$).

⁴Coefficient of variation.

Corn silage was the only sample presenting significant interactions between method and sequence of analysis ($p < 0.05$). In all non-sequential and sequential analyses there was significant difference ($p < 0.05$) between alternative and the conventional method. There was no difference ($p > 0.05$) as to the sequences of analysis within alternative method 3, but there was loss of analytical precision. Therefore, the use of alternative methods (1 and 2) is not recommended for the assessment of ADF contents in corn silage.

The results from all analyses did not differ as to the sequence (non-sequential or sequential), except for ADF determination in corn silage by alternative methods 1 and 2, which use autoclave digestion, and ANKOM and TNT bags, respectively.

Discussion

We observed that alternative methods depend on the feedstuff analyzed when compared to the conventional method. For NDF determinations, alternative method 1 is recommended for the analyses of Tifton hay; alternative method 2 for analyses of Tifton hay and babassu meal; and alternative method 3 for sugarcane and babassu meal. The NDF in hay was similar to that reported by Rodrigues *et al.* (2006) in plants at 28 days (80.80%) and 70 days of age (80.70%). NDF content in sugarcane was similar to that reported by Santos *et al.* (2008) at different cutting ages, 11 months (48.60%) and 24 months (56.88%), and by Rodrigues *et al.* (1997), who found variations from 45 to 56% in 11 sugarcane isolates.

The NDF values in corn silage are in accordance with the 55.44% observed by Valadares Filho *et al.* (2006). Fox *et al.* (1990), Van Soest (1994) and Carvalho (1995) reported NDF contents in corn silage varying from 46 to 63.2%, which is a wider variation compared with the present study.

The NDF content in babassu meal varied from 63.88 (alternative method 1) to 68.39% (conventional method). The contents observed by Rocha Júnior *et al.* (2003), Cavalcante *et al.* (2005) and Vieira *et al.* (2005; 64.50 to 78.70%) indicate a threshold that comprises the results obtained in this study. The NDF variation

obtained in samples of Xaraes grass was from 68.64 (conventional method) to 71.81% (alternative method 1), which is slightly lower to the content (73.40%) reported by Euclides (2002).

The mean NDF content in marandu grass varied from 72.51 (conventional method) to 76.15% (alternative method 1), differing from the variation reported by Araujo (2005), 69.80 to 74.31%, in a sub-humid region during the dry period with the use of irrigation.

The mean ADF contents for Tifton hay varied from 40.05 (conventional method) to 65.11% (alternative method 2), which is greater than the values reported by Gonçalves *et al.* (2003): 35.60% in hay at a cutting age of 25 days.

In sugarcane, variation from 31.02 (conventional method) to 42.91% (alternative method 2) was observed. Pate *et al.* (2001), analyzing the nutritional value of 66 commercial varieties of sugarcane planted in the south of Florida, observed a wide variation in ADF (28.30 to 41.50%), which are similar to values observed in our study.

The analyses of corn silage through the non-sequential test revealed ADF variations from 26.26 (conventional method) to 35.87% (alternative method 2). The ADF mean variation was 25.94 (conventional method) to 39.65% (alternative method 2). The mean ADF observed by Valadares Filho *et al.* (2006) was 30.80%, analyzing 112 samples of corn silage.

The results for babassu meal, 40.30 (conventional method) to 56.84% (alternative method 2), are close to those mentioned by Cavalcante *et al.* (2005) and Vieira *et al.* (2005), who reported values between 32.90 and 53.80%. The 48.30% mean ADF was also observed by Moreira Filho (2008), studying the chemical composition of six native species of babassu meal grazed by goats.

The variation of means obtained for xaraes grass ranged from 36.93 (conventional method) to 57.65% (alternative method 2). The mean of the results obtained in alternative method 2 is the highest compared with the means for the other methods

analyzed. Pereira *et al.* (2008) obtained a 38.96% mean value for ADF in xaraes grass, close to the value observed by the conventional method.

In marandu grass, the variation in the means was 39.06 (conventional method) to 60.97% (alternative method 2). The values observed in the alternative method 1 (46.42%) and alternative method 3 (43.24%) were near to those reported (46.42%) by Pereira *et al.* (2008); whereas, the mean (39.06%) for the conventional method was greater than the mean content (34.20%) reported by Santos *et al.* (2008), who studied marandu grass cultivars subjected to three different levels of fertilization. We also observed that the average results obtained with alternative method 2 were higher compared with the other methods studied.

During all ADF analysis, alternative method 2 had the greatest means and divergence from the conventional method. Casali *et al.* (2009) reported that TNT, in spite of being a fabric similar to ANKOM, does not present pores. This characteristic is given to TNT during the manufacturing process, in which part of the surface is sealed by heat. This fabric structure could broaden the estimates of analytical results. Thus, the results obtained allow us to suggest that it is still necessary to better investigate the use of bags fabricated with TNT in NDF and ADF analyses.

Only corn silage presented difference ($p < 0.05$) between sequences of analysis in alternative methods 1 and 2. The validation of alternative methodologies, which use the autoclave as digester system, to analyse NDF and ADF contents was not obtained for all the feedstuffs used; it was proven for Tifton 85 hay, babassu meal and sugarcane for NDF, but only in babassu meal at the determination of ADF.

In conclusion, the non-sequential and sequential forms of analysis and all analytical methods can be applied, with no loss of analytical precision for assessing NDF and ADF in tifton hay, sugar cane, babassu meal, Xaraes and Marandu grass. However, ADF quantification is not appropriate for corn silage using alternative method 2 (autoclave/TNT) and alternative method 3 (autoclave/filter crucible).

Conflicts of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

References

- Araujo DLC. Avaliação dos capins tifton (*Cynodon* spp), Tanzânia (*Panicum maximum*) e marandu (*Braquiária brizanta*) e terminação de ovinos em pastagens cultivadas com uso de suplementação. Dissertação (Mestrado em Ciência Animal) - Universidade Federal do Piauí, Teresina; 2005. 66f.
- Bialowas YD, Sullivan EC, Schneller RD. Designing a low-cost pollution prevention plan to pay off at the University of Houston. *J Air Waste Manag Assoc* 2006; 56:1320-1324.
- Carvalho MP. *Citrus*. In: Simpósio sobre nutrição de bovinos 6. Anais Piracicaba: FEALQ; 1995. p.171-241.
- Casali AO, Detmann E, Valadares Filho SC, Pereira JC, Cunha M. da, Detmann KSC, Paulino MF. Estimação de teores de componentes fibrosos em alimentos para ruminantes em saco de diferentes tecidos. *Rev Bras Zoot* 2009; 38:130-138.
- Cavalcante RR, Figueirêdo AV, Carvalho MAM, Lopes JB, Almeida MM. Digestibilidade aparente de nutrientes de rações balanceadas com alimentos alternativos para cutias (*Dasyprocta prymnolopha*) em crescimento. *Cienc Anim Bras* 2005; 6:163-171.
- Deschamps FC. Implicações do Período de crescimento na composição química e digestão dos tecidos de cultivares de capim-elefante (*Pennisetum purpureum* Schumach.). *Rev Bras Zoot* 1999; 28:1358-1369.
- Euclides VPB. Novidades em forrageiras para pecuária em regiões Tropicais. In: Seminário de pasturas y suplementacion estratégica em ganado bovino IICA. Asunción. Universidad Nacional de Asuncion - Facultad de Ciencias Veterinárias; 2002. p.12.
- Fox DG, Sniffen CJ, O'Connor JD, Russel JB, Van Soest PJ. The Cornell net carbohydrate and protein system for evaluating diets. Part I- A model for predicting cattle requirements and feedstuff utilization. Search: Agriculture. Agricultural Experimental Station, New York: Cornell University; 1990. 128p.
- Gerbase AE, Gregório JR, Calvete T. Gerenciamento dos resíduos da disciplina química inorgânica II do curso de química da Universidade Federal do Rio Grande do Sul. *Quím Nova* 2006; 29:397-403.
- Geredes L, Werner JC, Colozza MT, Possenti RA, Schammas EA. Avaliação de características de valor nutritivo das gramíneas forrageiras marandu, setária e tanzânia nas estações do ano. *Rev Bras Zoot* 2000; 29:955-963.

- Gonçalves GD, Santos GT, Jobim CC, Damasceno JC, Cecato U, Branco AF. Determinação do consumo, digestibilidade e frações protéicas e de carboidratos do feno de tifton 85 em diferentes idades de corte. *Rev Bras Zoot* 2003; 32:804-813.
- Instituto Nacional de Metrologia, Normalização e Qualidade Industrial. INMETRO. Vocabulário internacional de termos fundamentais e gerais de metrologia. 2. ed. Brasília; 2000. 75p.
- Instituto Nacional de Metrologia, Normalização e Qualidade Industrial. INMETRO. Orientações sobre validação de métodos de ensaios químicos: documento de caráter orientativo. Brasília; 2007. (DOQ-CGCRE-008 – revisão 02 de junho/2007).
- Jung HG. Analysis of forage fiber and cell walls in ruminant nutrition. *J Nutrition* 1997; 127:810S-813S.
- Komarek AR. A filter bag procedure for improved efficiency of fiber analysis. *J Dairy Sci* 1993; 76(Sup):1- 250.
- Moreira Filho MA. Composição bromatológica de seis espécies nativas do estado do Piauí consumidas por caprinos. *Pubvet* 2008; 2:34- art. 219.
- Pate FM, Alvarez J, Phillips JD, Eiland BR. Sugarcane as cattle feed: production and utilization. Florida: University of Florida/ Cooperative Extension Service; 2001. 25p.
- Pell AN, Schofield P. Computerized monitoring of gas production to measure forage digestion *in vitro*. *J Dairy Sci* 1993; 76:1063-1073.
- Pereira RC, Ribeiro KG, Pereira OG, Rigueira JPS, Silva JL, Santos JM. Composição químico-bromatológica em cultivares de *Brachiaria*. Anais. In: Simpósio nacional cerrado, Simpósio Internacional de Savanas Tropicais; 2008.
- Rocha Júnior VR, Valadares Filho SC, Borges AM, Magalhães KA, Ferreira CCB, Valadares RFD, Paulino MF. Determinação do valor energético de alimentos para ruminantes pelo sistema de equações. *Rev Bras Zoot* 2003; 32:473-479.
- Rodrigues AA, Primavesi O, Esteves SN. Efeito da qualidade de variedades de cana de-açúcar sobre seu valor como alimento para bovinos. *Pesq Agrop Bras* 1997; 32:1333-1338.
- Rodrigues LRA, Rodrigues TJD, Reis RA, Soares Filho CV. Produção de massa seca e composição química de cinco cultivares de *Cynodon*. *Anim Sci* 2006; 28:251-258.
- Santos LC, Bonomo P, Silva CCF, Pires AJV, Veloso CM, Patês NMS. Produção e composição química da *Brachiaria brizantha* e *Brachiaria decumbens* submetidas a diferentes adubações. *Cienc Anim Bras* 2008; 9:856-866.
- Senger CCD, Kozloski GV, Snachezn LMB, Mesquita FR, Alves TP, Castagnino DS. Evaluation of autoclave procedures for fibre analysis in forage and concentrate feedstuffs. *Anim Feed Sci and Techn* 2008; 146:169-174.
- Silva DJ, Queiróz AC. Análise de alimentos: métodos químicos e biológicos. 2. ed. Viçosa, MG: Universidade Federal de Viçosa; 2002. 235 p.
- SAS Institute. SAS/STAT User's guide: Statistics. Version 9.1®. 4th ed. SAS Inst. Inc., Cary, NC. 2002. 525p.
- Valadares Filho SC, Machado PAS, Chizzotti ML, Amartal HF, Magalhães KA, Rocha Junior VR, Capelle ER. Tabelas brasileiras de composição de alimentos para bovinos. Viçosa: Universidade Federal de Viçosa; 2006. 502 p.
- Van Soest PJ. Use of detergents in the analysis of fibrous feeds. II - a rapid method for determination of fiber and lignin. *J Assoc Offic Anal Chem* 1963; 46:829-835.
- Van Soest PJ. Development of a comprehensive system of feed analyses and its application to forages. *J Anim Sci* 1967; 26:119-128.
- Van Soest PJ, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci* 1991; 74:3583-3597.
- Van Soest PJ. Nutritional ecology of the ruminant. 2. ed. Ithaca: Comstock Publishing Association; 1994. 476 p.
- Vieira MMM, Cavalcante MAB, Neiva JNM. Valor nutritivo de silagens de capim elefante (*Pennisetum purpureum*) contendo diferentes níveis de farelo de babaçu (*Orbignya* sp.). In: Reunião Anual da Sociedade Brasileira de Zootecnia, Goiânia, 2005. Anais Goiânia: SBZ; 2005. 1 CD-ROM.