

SHORT COMMUNICATION

Genetic effect of season on the preweaning growth of beef cattle: A first approach on Retinta calves

Efecto genético de la estación climática sobre el crecimiento predestete del ganado de carne: Un primer estudio en terneros de raza Retinta

Efeito genético de estações climáticas no crescimento pré-desmame de bovinos de corte: uma primeira abordagem para os bezerros Retinta

Rosa M Morales^{1*} ; Alberto Menéndez-Buxadera¹ ; Sebastián Demyda-Peyrás² ; Antonio Molina¹ .

¹Departamento de Genética, Facultad de Veterinaria, Universidad de Córdoba, España

²IGEVET- Instituto de Genética Veterinaria "Ing. Fernando N Dulout", Universidad Nacional de La Plata, Consejo Nacional de Investigaciones Científicas y Técnicas (UNLP-CONICET LA PLATA), Facultad de Ciencias Veterinarias UNLP, La Plata, Argentina.

To cite this article:

Morales RM, Menéndez-Buxadera A, Demyda-Peyrás S, Molina A. Genetic effect of season on the preweaning growth of beef cattle: A first approach to Retinta calves. Rev Colomb Cienc Pecu 2020; 33(2): 134-143. DOI: <https://doi.org/10.17533/udea.rccp.v33n2a01>

Abstract

Background: Heat stress derived from global warming is causing major economic losses in the livestock industry. **Objective:** To develop a novel methodological approach for determining the influence of climatic factors on the estimation of genetic parameters for growth traits in Retinta cattle breed by using reaction-norm models. **Methods:** Live weight records (n=7,753) from 3,162 Retinta calves born from 1,249 dams and 85 sires and raised in the Andalusian region (Spain) were analyzed. The effect of heat stress was measured using the temperature-humidity index, calculated with climatological data obtained from four weather stations. A bivariate-random-regression reaction-norm model was used to estimate the (co)variance components of weight until weaning in two different climatic seasons corresponding to warm and cold months. **Results:** The heritability pattern of individuals reared under diverse environments during the first 90 days of age was different. However, differences were not significant at the end of the growing period. Weaned calves reared during the cold season showed greater growth from 70 to 160 days in comparison with those reared during the warm season. **Conclusions:** Overall, this assessment did not show significant effects of the genotype-environment interaction. However, highly significant evidence of genotype-climatic condition interaction was found during the calf's first three months of age.

Keywords: *beef cattle, genotype-environment interaction, genetic effects, global warming, heat stress, preweaning growth, random regression, reaction-norm, Retinta breed, THI.*

Received: May 21, 2018; accepted: August 9, 2019

*Corresponding Author. Ctra. Madrid-Cádiz, km 396, 14071 Córdoba. España. Tel.: +34 957 21 10 70; Fax: +34 957 21 87 07. E-mail: y22mocer@uco.es



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Resumen

Antecedentes: El estrés térmico derivado del calentamiento global está causando pérdidas económicas en la industria ganadera. **Objetivo:** Desarrollar un enfoque metodológico para estimar la influencia de algunos factores climáticos sobre la estimación de parámetros genéticos en las variables de crecimiento de la raza bovina Retinta. **Métodos:** Se analizaron registros de peso vivo ($n=7.753$) de 3.162 terneros hijos de 1.249 vacas y 85 toros, criados en la región de Andalucía (España). El efecto del estrés térmico se midió mediante un índice de temperatura-humedad obtenido a partir de los datos de cuatro estaciones meteorológicas. Se usó un modelo bivariado de regresión aleatoria de “norma-reacción” para estimar la (co)varianza del peso hasta el destete en dos diferentes épocas climáticas correspondientes a los meses cálidos y fríos. **Resultados:** Se encontraron diferencias en el patrón de heredabilidad de individuos criados en diferentes ambientes durante los primeros 90 días de vida. Sin embargo, tales diferencias no fueron significativas al final del periodo de crecimiento. Los terneros destetados en la época fría mostraron un mayor crecimiento entre 70 y 160 días en comparación con los criados en la temporada cálida. **Conclusiones:** Se encontró una evidencia altamente significativa de la interacción genotipo-condición climática durante los primeros tres meses de crecimiento del ternero.

Palabras clave: calentamiento global, crecimiento predestete, efectos genéticos, estrés climático, ganado de carne, interacción genotipo-ambiente, norma-reacción, raza Retinta, regresión aleatoria, THI.

Resumo

Antecedentes: O estresse térmico devido ao aumento da temperatura média está produzindo grandes perdas econômicas na indústria pecuária. **Objetivo:** Realizar uma nova abordagem metodológica para estimar a importância dos fatores climáticos em parâmetros genéticos em variáveis de crescimento da população da raça Retinta. **Métodos:** Foram analisados registros de peso vivo ($n=7.753$) de 3.162 Retinta bezerros, nascidos de 1.249 vacas e 85 touros criados na região da Andaluzia (Espanha). O efeito do estresse térmico foi medido utilizando o índice de temperatura-umidade obtido com dados climatológicos de quatro estações meteorológicas. Um modelo de regressão aleatória bivariado de norma-reação foi usado para estimar os componentes da (co)variância do peso à desmama em duas estações climáticas com correspondência aos meses quentes e frios. **Resultados:** Diferenças foram encontradas no padrão de herdabilidade de indivíduos criados em diferentes ambientes nos primeiros 90 dias de idade. No entanto, essas diferenças não foram significativas no final do período de crescimento. Os bezerros desmamados globalmente na estação fria apresentaram um crescimento maior dos 70 aos 160 dias em comparação com os indivíduos criados na estação quente. **Conclusões:** Uma evidência altamente significativa da interação genótipo-ambiente foi encontrada durante os primeiros três meses de crescimento do bezerro.

Palavras-chave: aquecimento global, bovino de corte, crescimento pré-desmame, efeitos genéticos, estresse por calor, interação genótipo-ambiente, norma de reação, raça Retinta, regressão aleatória, THI.

Introduction

Extensive livestock production is highly dependent on weather conditions. Nowadays, cattle reared under extensive conditions are exposed to increasing heat stress, which is expected to worsen in the next decades (IPCC, 2014). Such an adverse situation is partially solved by using autochthonous breeds since they are better adapted to the local environment.

In cattle, heat stress has been associated with decreased productivity and economic losses (Mader, 2003; Nienaber and Hahn, 2007; St-Pierre *et al.*, 2003), normally induced by reduced feed intake (Collier and Zimbelman, 2007). In extreme cases, heat stress has been associated with negative energy balance and body weight loss (Lacetera *et al.*, 1996), as well as with a reduced immune system response (Silanikove, 2000). The first experimental approach to select animals based on heat stress tolerance was proposed by Misztal (1999). This researcher evaluated the productive reaction of individuals subjected to increasing heat loads (reaction-norm model) to determine the genotype-environment interaction and detect the best genetically adapted individuals to an increased temperature range (Santana *et al.*, 2016). Later, heat stress tolerance was included as a secondary trait in dairy selection schemes focused on increased production (Bohmanova *et al.*, 2005; Brugemann *et al.*, 2011; Carabano *et al.*, 2017; Ravagnolo and Misztal, 2000; Sanchez *et al.*, 2009). However, few studies have been performed in *Bos taurus* (Bradford *et al.*, 2016) and *Bos indicus* (Santana *et al.*, 2016) grazing cattle, and -to our knowledge- no studies have been performed in Spanish breeds.

Retinta is an autochthonous breed widely used in the south of Spain due to its excellent adaptation to the extreme environment that prevails in that region (Morales *et al.*, 2013). These individuals are raised in an extensive regime characterized by pasturelands with limited production, steep hills and a dry and hot

climate (Rodero-Serrano *et al.*, 2013). Under these herd conditions, the effect of heat stress is expected to rapidly manifest as weight loss and reduced reproductive potential in poorly adapted individuals (Vercoe and Frisch, 1980).

Therefore, the objective of this study was to develop a novel methodological approach to determine the influence of weather variables on the estimation of genetic parameters for growth traits in Retinta cattle by random regression (reaction-norm) models.

Materials and methods

Animals and growth records

Growth records of 3,162 calves born from 85 sires and 1,249 cows were collected between birth and 245 days of age. All calves were weighed at least once between birth and 4 months of age, and at least another time between 4 to 8 months of age. The animals belonged to 14 Retinta herds located in the south of Spain and were reared during a nine-year period (two-generation intervals). Farms were selected to maximize the pedigree completeness, more than 3 mean equivalent generations (Maignel *et al.*, 1996), production records and the genetic links among herds, which was achieved by using insemination with 45 connecting sires. Weight records outside the range of ± 3 standard deviations from the average were excluded, resulting in 7,753 live weight records from calves included in this study (Table 1). The relationship matrix (A) included a pedigree with a total of 6,053 animals and their ancestors. This genealogical records information was obtained from the herdbook of the Retinta Breeder's Association. Mean maximum generation was 11.70, mean complete generation was 3.79, and mean equivalent generation was 6.27, as determined by ENDOG (Gutiérrez and Goyache, 2005).

Table 1. Raw dataset and selected population.

Data type	Raw dataset	Selection Criteria			
		Genetic links among herds (by 45 connecting sires)	Mean equivalent generations >3	Calves weighed twice (between birth and 4 months of age, and between 4 and 8 months of age)	Weight records outside the range of ± 3 SD*
Farms	44	17	14	14	14
Calves	13,640	4,712	3,382	3,197	3,162
Cows	5,403	1,677	1,521	1,281	1,249
Sires	309	111	94	89	85
Weight records	13,828	9,327	8,726	7,813	7,753

*SD: standard deviations.

Climate data

Climate data were obtained from four weather stations of the Spanish Meteorological State Agency (Agencia Estatal de Meteorología—AEMET), located near the farms. Weather stations collect climatic information (temperature, relative humidity, degree of insolation, solar radiation, precipitation, and speed and wind direction) every hour. From this information, the average temperature and humidity were calculated to estimate the temperature and humidity index (THI), according to Finocchiaro *et al.* (2005) as:

$$\text{THI} = \text{Tdb} - (0.55 - 0.0055 \times \text{RH}) \times (\text{Tdb} - 14.4)$$

Where Tdb = dry bulb temperature ($^{\circ}\text{C}$) and RH = relative humidity of the air (%).

Solar radiation (SR; MJ/m² per day) was also registered to define the annual variation and determine the existence of weather seasons based on systematic records.

The herd heat load was estimated as the average THI 30 days before each weight assessment.

Reaction-norm model

Individual live weight results recorded during the high heat stress (HHS) and low heat stress (LHS) seasons were considered as two different traits. The components of variance through the trajectory of the pre-weaning age intra-stress zone were estimated applying a bi-character random regression model:

$$y_{ijkl:z} = \text{fixed} + \sum_{r=0}^3 \Phi_r \beta_{r:z} + \left[\sum_{r=0}^3 \Phi_r \lambda_{rk} a_{ir} \right]_{-z} + Z_1 p_i + e_{ijkl}$$

where $y_{ijkl:Z}$ is the i^{th} records of live weight in each z stress zone; **fixed** is a set of fixed effects: sex of the calf with two levels (S_j ; $j=2$ level), parity effect of the dam with ten levels (CNk, $k = 1, 2, \dots, 6$ or more calvings) and combined herd-year with 355 levels. Age effects were analyzed by a fixed covariable intra-stress zone (z) by a Legendre polynomial (Φ) of order $r = 3$.

Genetic effects (a_i) were calculated by a random regression matrix (λ_{rk}) intra-stress zone with a Legendre polynomial (Φ) of order $r = 1$; Z_1 is an incidence matrix of permanent

environmental random effects (p_i) due to repetitions of the same trait in the animal, and e_{ijkl} is a random error common to all observations. A residual intra-stress zone variance was evaluated, but it did not contribute to an increase in the model fit.

The significance of the effects included in the model was determined with the Wald test (Kenward and Roger, 1997). All the statistical analyses were carried out using the Asreml software (Gilmour *et al.*, 2009).

In this reaction-norm animal model, the variance can be estimated as follows:

$$\text{var}(y) = \Phi_1 (\lambda_{k;z} \otimes A) \Phi_1' + I_p \sigma_p^2 + I_n \sigma_e^2$$

Where A is the numerator relationship matrix, $I_p \sigma_p^2$ and $I_n \sigma_e^2$ are the variances of permanent environmental random effects and residual effect, with matrices of identity I_p and I_n , respectively. These variance components are given directly by Asreml, while the expected (co)variance components through different ages of the cow for each stress-zone can be represented by a further random regression matrix manipulation $\lambda_{k;z}$ with the following four sub-matrices:

$$\lambda_k = \left[\begin{array}{cc} \lambda_h = \begin{pmatrix} \sigma_{ih}^2 & \sigma_{is,h} \\ \sigma_{si,h} & \sigma_{sh}^2 \end{pmatrix} & \lambda_{hl} = \begin{pmatrix} \sigma_{ih,il} & \sigma_{ih,sl} \\ \sigma_{sc,ib} & \sigma_{sh,sl} \end{pmatrix} \\ \lambda_{lh} = \begin{pmatrix} \sigma_{il,ih} & \sigma_{il,sh} \\ \sigma_{ls,ih} & \sigma_{sl,sh} \end{pmatrix} & \lambda_l = \begin{pmatrix} \sigma_{il}^2 & \sigma_{is,l} \\ \sigma_{sl,l} & \sigma_{sl}^2 \end{pmatrix} \end{array} \right] \otimes A$$

The diagonal sub-matrix elements ($\lambda_{h,y}$ λ_l) are used to estimate the variances of live weight over pre-weaning age under HHS and LHS environmental conditions, symbolized as **h** and **l**, respectively, while the correspondent (co)variances are contained in $\lambda_{hl} = \lambda_{lh}$. The (co)variance estimates were obtained applying the procedure proposed by Jong (1995). Details of practical application with data of

this breed were recently presented by Morales *et al.* (2013). Heritability at j age and genetic correlations between any pair of ages in both stress zones were determined by classical methodologies.

Results

The analysis of the weather data collected during a nine-year period showed two well-differentiated periods with minimal variations within each period: a warm season from April to September (HHS) and a cold season from October to March (LHS). The former was characterized by a 40% increased SR, and 20% increased THI compared with LHS (Figure 1).




Figure 1. Progression of temperature, humidity (THI), and solar radiation (SR) index during pre-weaning growth of Retinta cattle in two climatic categories, warm or high heat stress (HHS) from months 4 to 9, and cold or low heat stress (LHS) from month 10 to 3 years of age.

All the included effects were highly significant ($p < 0.001$, Wald test), as well as weather-season-age of the calf interaction. Differences in the progression of live weight during the pre-weaning age trajectory of Retinta calves and each climatic class were significant ($p < 0.001$, Wald test) (Figure 2).




Figure 2. Progression of live weight of Retinta calves during pre-weaning.

Live weight variance components and heritabilities were estimated in each climatic season during the pre-weaning period, showing a wide range of variation throughout the calf's age. Heritability differences in the HHS and LHS zones were highly significant during the first 90 days of growth ($p < 0.001$, Wald test). After this point (end of the third month), differences were not significant ($p > 0.05$), converging to zero after 120 days of growth (Figure 3). Likewise, the genetic correlations of live weights in the different climatic periods showed the same trend, with values lower than 1 during the first three months of age, and close to 1 thereafter (Figure 3).




Figure 3. Heritability and genetic correlations of live weight through the pre-weaning trajectory in each climatic class.

Discussion

In this study, we aimed to demonstrate the flexibility and potentiality of reaction-norm random regression models to evaluate the genotype-environment interaction in extensive cattle production systems. Despite of not including some environmental factors in this analysis (e.g., nutritional management, livestock load per acre, or grazing strategies) and the weather analyzed included only the Mediterranean climate range (CSA, according to the Köppen-Geiger climate classification; Kottek *et al.*, 2006) our results showed the existence of a genetic influence on weaning weight under stressful environmental conditions. In addition, climatic conditions could have a strong influence on weight increase, particularly during the first months of life. However, the genetic correlation with climatic periods lower than 1 in the first three months of age suggests the existence of a highly significant interaction between the environment and the calf genotype in that specific period.

In this study, we used an analytical methodology previously reported in dairy cattle (Aguilar *et al.*, 2009; Ravagnolo and Misztal, 2000) and swine (Fragomeni *et al.*, 2015; Zumbach *et al.*, 2008) raised in confined systems, in which the environmental conditions were tightly controlled, with minimum variation among farms. On the other hand, Retinta cattle are raised in the typical environment of the south of Spain and the Andalucía region, characterized by harsh weather, cold winters and dry and hot summers, with abrupt changes between seasons. In our case, reaction-norm models proved to be valid to evaluate the heat-stress tolerance of grazing animals under such conditions, which is extremely important since those individuals are more prone to be exposed to heat stress (Tucker *et al.*, 2008).

In a previous study, Bradford *et al.* (2016) determined a heritability close to 0.24 for weaning weight in Angus cattle using reaction-norm models with intermediate heat loads. On the

contrary, the results obtained with calves raised under more favorable environmental conditions and in zebu cattle were higher (Cardoso and Tempelman, 2012; Santana *et al.*, 2016). In our study, heritabilities showed a wide range of variation throughout the calves' age, which could be partially explained by differences in heat stress tolerance among individuals' age or by differences in the availability and quality of pasturelands that influences milk production. However, it has also been demonstrated that milk production is severely affected by individual heat stress tolerance, suggesting that those animals could be affected not only by the heat stress tolerance of their genotype, but also by the genotype of their mother (Aguilar *et al.*, 2009; Bohmanova *et al.*, 2005; Carabano *et al.*, 2014; Sanchez *et al.*, 2009). Likewise, it is noteworthy that genetic correlations between climatic periods were extremely high during the last period of growth of the calf, in which the dependence on maternal milk is much lower.

Despite the overall interaction was not significant, genetic differences between climatic seasons were significant at specific growth periods. This phenomenon was also recently described by Santana *et al.* (2016), who observed substantial heterogeneity in the (co)variance components for weaning weight analyzing different environmental conditions, suggesting that it was an evidence of the genotype-environment interaction. Moreover, those researchers demonstrated that the best-fitted animals detected in the least stressful environments were not necessarily the most suitable ones, and even not suitable at all, to be raised in the most stressful environments. In our study, individuals that grew up in HHS suffered a highly stressful environment in comparison with the group reared in LHS. Nevertheless, the number of records analyzed in each category was homogeneous, suggesting that Retinta calves are well adapted to the environmental conditions observed in HHS.

Based on the current knowledge, low response to selection for weaning weight in populations

reared under stressful environments is expected, since greater growth appeared from 70 to 160 days in animals raised during the cold season (Figure 2). However, differences were not significant when they were considered altogether, suggesting the existence of a compensatory growth that equalizes calf weights after this age. It is also known that the dam begins to reduce milk production during pre-weaning, thus forcing the calf to consume grass, thus the genotype-environment interaction would be more evident during this specific period. Therefore, our results suggest that the effect of calving season and heat load could be exerted through the grazing behavior of the calf, which is clearly associated with thermal stress (Santana *et al.* 2016).

In conclusion, differences observed in the heritability pattern of individuals reared in diverse environments during the first 90 days of age were not significant at the end of the growing period. Our results confirm a clear variability among individuals since animals raised during the cold season showed a greater growth from 70 to 160 days. This information could be used as a baseline for future studies including ample range of rearing environments as well as the nutritional impact produced by environmental differences.

Declarations

Acknowledgments

The authors thank the National Retinta Breeders Association and the CEAG for data contribution and financial support. The provision of meteorological records by the Spanish Meteorological State Agency (Agencia Estatal de Meteorología—AEMET) is also appreciated. Thanks are also due to A. Di Maggio for manuscript correction.

Funding

National Retinta Breeders Association and CEAG (Experimental Agriculture and Livestock Centre of Jerez de la Frontera, Rural Council of Cadiz).

Conflicts of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

Author contribution

Rosa M Morales: Project administrator, data acquisition and curation, formal analysis, data visualization and writing of the original manuscript.

Alberto Menéndez-Buxadera: Experimental design and formal analysis of the data.

Sebastián Demyda-Peyrás: writing of the original manuscript. Critical reviewing and writing of the final draft.

Antonio Molina: Experimental design, investigation, formal analysis of the data and critical reviewing.

All the authors read and approved the final manuscript.

References

- Aguilar I, Misztal I, Tsuruta S. Genetic components of heat stress for dairy cattle with multiple lactations. *J Dairy Sci* 2009; 92:5702-11. DOI: <https://doi.org/10.3168/jds.2008-1928>
- Bohmanova J, Misztal I, Tsuruta S, Norman HD, Lawlor TJ. National genetic evaluation of milk yield for heat tolerance of United States Holsteins. *Interbull Bulletin* 2005; 33:160-2. DOI: [https://doi.org/10.3168/jds.S0022-0302\(04\)73216-6](https://doi.org/10.3168/jds.S0022-0302(04)73216-6)
- Bradford HL, Fragomeni BO, Bertrand JK, Lourenco DA, Misztal I. Genetic evaluations for growth heat tolerance in Angus cattle. *J Anim Sci* 2016; 94:4143-50. DOI: <https://doi.org/10.2527/jas.2016-0707>
- Brugemann K, Gernand E, von Borstel UU, König S. Genetic analyses of protein yield in dairy cows applying random regression models with time-dependent and temperature x humidity-dependent covariates. *J Dairy Sci* 2011; 94:4129-39. DOI: <https://doi.org/10.3168/jds.2010-4063>
- Carabano MJ, Bachagha K, Ramon M, Diaz C. Modeling heat stress effect on Holstein cows under hot and dry conditions: selection tools. *J Dairy Sci*. 2014;97(12):7889-904. Epub 2014/09/30. DOI: <https://doi.org/10.3168/jds.2014-8023>
- Carabano MJ, Ramon M, Diaz C, Molina A, Perez-Guzman MD, Serradilla JM. Breeding and genetics symposium: Breeding for resilience to heat stress effects in dairy ruminants. A comprehensive review. *J Anim Sci* 2017; 95:1813-26. DOI: <https://doi.org/10.2527/jas.2016.1114>
- Cardoso FF, Tempelman RJ. Linear reaction norm models for genetic merit prediction of Angus cattle under genotype by environment interaction. *J Anim Sci* 2012; 90:2130-41. DOI: <https://doi.org/10.2527/jas.2011-4333>
- Collier RJ, Zimbelman RB. Heat stress effects on cattle: what we know and what we don't know. Proc of the Southwest Nutrition and Management Conference, The University of Arizona, Tucson, February 23rd. 2007.
- Finocchiaro R, van Kaam JB, Portolano B, Misztal I. Effect of heat stress on production of Mediterranean dairy sheep. *J Dairy Sci* 2005; 88:1855-64. DOI: [https://doi.org/10.3168/jds.S0022-0302\(05\)72860-5](https://doi.org/10.3168/jds.S0022-0302(05)72860-5)
- Fragomeni B, Tsuruta S, Lourenco D, Gray K, Huang Y, Misztal I. Genomic mitigation of seasonality effect on carcass weight in commercial pigs. *J Anim Sci* 2015; 93:847.
- Gilmour AR, Gogel BJ, Cullis BR, Thompson R. ASReml User Guide Release 3.0. United Kingdom: VSN International Ltd; 2009.
- Gutiérrez JP, and Goyache F, A note on ENDOG: a computer program for analysing pedigree information. *J Anim Breed Genet* 2005; 122:172-176. DOI: <https://doi.org/10.1111/j.1439-0388.2005.00512.x>

- IPCC. Climate Change 2014 Synthesis Report: Fifth Assessment Report. In: Meyer RKPALA, editor. Geneva, Switzerland 2014. p. 151.
- Jong G. Phenotypic plasticity as a product of selection in a variable environment. *Amer Nat* 1995;145:493-512.
- Kenward MG, Roger JH. The precision of fixed effects estimates from restricted maximum likelihood. *Biometrics* 1997; 53:983-997.
- Kottek M, Grieser J, Beck C, Rudolf B, Rubel F. World map of the Köppen-Geiger climate classification updated. *Meteorol Z* 2006; 15:259-63.
- Lacetera N, Bernabucci U, Ronchi B, Nardone A. Body condition score, metabolic status and milk production of early lactating dairy cows exposed to warm environment. *Riv Agric Subtrop Trop* 1996; 90(1):43-55.
- Mader TL. Environmental stress in confined beef cattle. *J Anim Sci* 2003; 81:110-9. DOI: https://doi.org/10.2527/2003.8114_suppl_2E110x
- Maignel L, Boichard D, Verrier E. Genetic variability of French dairy breeds estimated from pedigree information. *Interbull Bull* 1996; 14:49-54.
- Misztal I. Model to study genetic component of heat stress in dairy cattle using national data. *J Dairy Sci* 1999;82:32.
- Morales R, Menéndez-Buxadera A, Avilés C, Molina A. Direct and maternal genetic effects for preweaning growth in Retinta cattle estimated by a longitudinal approach throughout the calving trajectory of the cow. *J Anim Breed and Genet* 2013;130:425-34. DOI: <https://doi.org/10.1111/jbg.12038>
- Nienaber JA, Hahn GL. Livestock production system management responses to thermal challenges. *Int J Biometeorol* 2007;52:149-57. DOI: <https://doi.org/10.1007/s00484-007-0103-x>
- Ravagnolo O, Misztal I. Genetic component of heat stress in dairy cattle, parameter estimation. *J Dairy Sci* 2000;83:2126-30. DOI: [https://doi.org/10.3168/jds.S0022-0302\(00\)75095-8](https://doi.org/10.3168/jds.S0022-0302(00)75095-8)
- Rodero-Serrano E, Demyda-Peyrás S, González-Martínez A, Rodero-Franganillo A, Moreno-Millán M. The rob(1;29) chromosome translocation in endangered Andalusian cattle breeds. *Livestock Science* 2013;158:32-9. DOI: <https://doi.org/10.1016/j.livsci.2013.10.001>
- Sanchez JP, Misztal I, Aguilar I, Zumbach B, Rekaya R. Genetic determination of the onset of heat stress on daily milk production in the US Holstein cattle. *J Dairy Sci* 2009; 92:4035-45. DOI: <https://doi.org/10.3168/jds.2008-1626>
- Santana ML, Jr., Bignardi AB, Eler JP, Ferraz JB. Genetic variation of the weaning weight of beef cattle as a function of accumulated heat stress. *J Anim Breed and Genet* 2016; 133:92-104. DOI: <https://doi.org/10.1111/jbg.12169>
- Silanikove N. Effects of heat stress on the welfare of extensively managed domestic ruminants. *Livest Prod Sci* 2000; 67:1-18. DOI: [https://doi.org/10.1016/S0301-6226\(00\)00162-7](https://doi.org/10.1016/S0301-6226(00)00162-7)
- St-Pierre NR, Cobanov B, Schnitkey G. Economic Losses from Heat Stress by US Livestock Industries1. *J Dairy Sci* 2003;86, Supplement:E52-E77. DOI: [https://doi.org/10.3168/jds.S0022-0302\(03\)74040-5](https://doi.org/10.3168/jds.S0022-0302(03)74040-5)
- Tucker CB, Rogers AR, Schütz KE. Effect of solar radiation on dairy cattle behaviour, use of shade and body temperature in a pasture-based system. *Appl Anim Behav Sci* 2008; 109:141-54. DOI: <https://doi.org/10.1016/j.applanim.2007.03.015>
- Vercoe JE, Frisch JE. Animal breeding and genetics with particular reference to beef cattle in the tropics. 4th World Conference on Animal Production. Buenos Aires, Argentina 1980. p. 452-63.

Zumbach B, Misztal I, Tsuruta S, Sanchez JP, Azain M, Herring W, *et al.* Genetic components of heat stress in finishing pigs: Parameter estimation. *J Anim Sci* 2008; 86:2076-81. DOI: <https://doi.org/10.2527/jas.2007-0282>