













Influence of the storage period on the nutritional and microbiological value of sun-dried brewer's grains

Influencia del período de almacenamiento sobre el valor nutricional y microbiológico de los granos de cervecería secados al sol

Influência do período de estocagem sobre o valor nutricional e microbiológico de grãos de cervejaria secos ao sol

Andressa Faccenda^{1*} ; Maximiliane A Zambom¹ ; André S Avila¹ ; Deise D Castagnara² ;
Ricardo Dri¹ ; Maria L Fischer¹ ; Rodrigo C R Tinini¹ ; Jéssica G Dessbesell¹ ; Ana-Ruth E Almeida¹ ;
Kleves V Almeida¹ .

¹Universidade Estadual do Oeste do Paraná, Paraná, Brazil.

²Universidade Federal do Pampa, Rio Grande do Sul, Brazil.

To cite this article:

Faccenda A, Zambom MA, Avila AS, Castagnara DD, Dri R, Fischer ML, Tinini RCR, Dessbesell JG, Almeida ARE, Almeida KV. Influence of storage period on the nutritional and microbiological value of sun-dried brewer's grains. Rev Colomb Cienc Pecu 2021; 34(4): 254–266. DOI: <https://doi.org/10.17533/udea.rccp.v34n4a02>

Abstract

Background: Brewer's grains, a by-product of the brewery industry, can be included in the diet of ruminants. However, its high humidity makes it difficult to store and preserve. **Objective:** To evaluate the efficiency of sun dehydration of wet brewer's grains (WBG) and the effect of storage period on its nutritional and microbiological quality. **Methods:** A completely randomized experimental design was used to evaluate WBG dehydration efficiency, with treatments corresponding to 0, 1, 2, 4, 6, 8, 10, 12, 14 and 16 hours of sun exposure. A second experiment was carried out using also a completely randomized design to evaluate the effect of storage with the following treatments: 0, 10, 20, 30, 60, 90, 120, 150 and 180 days of storage of the dry by-product. **Results:** Dry matter (DM) content linearly increased with dehydration period. The chemical composition of the dried brewer's grains had no effect as a function of storage period. Indigestible protein (C fraction) increased linearly but did not compromise the cumulative gas production and the *in vitro* digestibility of DM and protein. Storage time had no effect on fungus population. The maximum aflatoxin value was 45.5 µg/kg, and remained within acceptable limits for bovine feed. **Conclusion:** Dehydration of WBG in the sun is efficient to guarantee conservation and makes it possible to store the by-product. The storage of the dry by-product for 180 days does not compromise its nutritional or microbiological quality.

Received: March 4, 2019; accepted: September 20, 2020

*Corresponding author. Rua Pernambuco 1777, Marechal Cândido Rondon, Paraná, Brasil. CEP:85960-000. Tel.: +55 45 991067625. Email: andressafaccenda@hotmail.com



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Keywords: *barley; brewery by-product; brewer's grains; digestibility; dried brewer's grains feed conservation; feed by-product; fungi; nutritional value; ruminants; storage time.*

Resumen

Antecedentes: Los granos de cervecería son un subproducto de la industria cervecera que puede ser incluido en la dieta de rumiantes; sin embargo, su alta humedad dificulta el almacenamiento y conservación de ese producto. **Objetivo:** Evaluar la eficiencia de la deshidratación al sol de los granos húmedos de cervecería (WBG) y el efecto del período de almacenamiento sobre su calidad nutricional y microbiológica. **Métodos:** Para evaluar la eficiencia de la deshidratación de los WBG se utilizó un diseño experimental completamente al azar, con tiempos de tratamiento de 0, 1, 2, 4, 6, 8, 10, 12, 14, y 16 horas de exposición al sol. En un segundo experimento, también con diseño experimental completamente al azar, se evaluó el efecto del almacenamiento comparando los siguientes tratamientos: 0, 10, 20, 30, 60, 90, 120, 150 y 180 días de almacenamiento del subproducto seco. **Resultados:** La materia seca (DM) del WBG presentó un efecto lineal creciente con el proceso de deshidratación. La composición química de los granos secos de cervecería no tuvo efecto en función de los tiempos de almacenamiento. La proteína no digestible (fracción C) aumentó linealmente; sin embargo, no comprometió la producción acumulativa de gas y la digestibilidad *in vitro* de la DM y de la proteína. El tiempo de almacenamiento no tuvo efecto sobre la población de hongos. El valor máximo de aflatoxina obtenido fue de 45,5 µg/kg y permaneció dentro de los límites aceptables para alimentación de bovinos. **Conclusión:** La deshidratación de WBG al sol es eficiente para garantizar la conservación del material y viabilizar su almacenamiento. El almacenamiento por 180 días de este subproducto seco no compromete su calidad nutricional y microbiológica.

Palabras clave: *cebada; conservación de alimentos; digestibilidad; granos de cervecería; granos secos de cervecería; hongos; periodo de almacenamiento; rumiantes; subproducto alimenticio; subproductos de cervecería; valor nutricional.*

Resumo

Antecedentes: Os grãos de cervejaria são um subproduto que podem ser incluídos na dieta de ruminantes, no entanto sua forma úmida dificulta o armazenamento e a conservação desse produto. **Objetivo:** Avaliar a eficiência de desidratação ao sol dos grãos úmidos de cervejaria (WBG) e o efeito do período de armazenamento sobre a qualidade nutricional e microbiológica. **Métodos:** Para avaliar a desidratação do WBG, utilizou-se um delineamento experimental inteiramente casualizado, sendo os tratamentos os tempos de 0 e 1, 2, 4, 6, 8, 10, 12, 14 e 16 horas de exposição ao sol. Um segundo experimento foi realizado em delineamento inteiramente casualizado. Os tratamentos foram 0, 10, 20, 30, 60, 90, 120, 150 e 180 dias de armazenamento do subproduto seco. **Resultados:** A matéria seca (DM) do WBG apresentou efeito linear crescente com o processo de desidratação. A composição química dos grãos secos de cervejaria não apresentou efeito em função dos tempos de armazenagem. A proteína indigestível (Fração C) aumentou linearmente, no entanto não comprometeu a produção cumulativa de gás e a digestibilidade *in vitro* da DM e da proteína. A população de fungos não apresentou efeito com o tempo de armazenamento. O valor máximo de aflatoxina obtido foi de 45,5 µg/kg e permaneceu dentro dos limites aceitáveis para a alimentação de bovinos. **Conclusão:** A desidratação do WBG ao sol foi eficiente em garantir a conservação do material e viabilizar o seu armazenamento. A estocagem desse subproduto seco por 180 dias não comprometeu a sua qualidade nutricional e microbiológica.

Palavras-chave: *cevada; conservação; digestibilidade; fungos; grãos de cervejaria; grãos secos de cervejaria; ruminantes; subproduto de alimentação; subproduto de cervejaria; tempo de armazenamento; valor nutricional.*

Introduction

Wet brewer's grains are an agroindustry residue generated during the brewing process and represents approximately 85% of the total residue obtained in this process (Mussatto *et al.*, 2006). The high protein content of brewer's grains makes this ingredient an alternative protein source for ruminants and can replace traditional feedstuffs such as soybean meal, with several advantages: lower cost, does not compete with human nutrition, and contributes to sustainability of animal production (Negesse *et al.*, 2009).

However, the high moisture content of this residue is the most limiting factor in its use because it makes it difficult to transport, store, and preserve, making it more susceptible to development of molds, yeasts, and deterioration (Wang *et al.*, 2014). Thus, dehydration is an alternative for the preservation of brewer's grains. Dehydration is interesting because, in addition to avoiding quality loss, it decreases the volume of the material facilitating storage. However, dehydration using high temperature is not economically viable. Moreover, even when the dehydration temperature used is less than 60 °C there is a risk of toasting the grains (Mussatto *et al.*, 2006) and reducing its nutritional value due to complexation of proteins with cell wall compounds, rendering them unavailable (Pereira *et al.*, 1998). Alternatively, sun dehydration offers advantages, such as cost reduction and drying at lower temperatures.

After dehydration, chemical and physical modifications of the residue during storage are possible, especially if the storage site lacks adequate ventilation and protection against moisture. In addition, storage time itself may interfere with quality due to the aging of the product. Therefore, monitoring waste quality is necessary to determine the ideal storage period, thus ensuring that it can be used safely in animal feed.

In this context, the objective of this study was to evaluate the efficiency of dehydration of wet brewer's grains (WBG) in the sun,

as well as the effect of storage period on the nutritional and microbiological quality of the dried byproduct (DBG).

Materials and Methods

Ethical considerations

The experimental protocol for the use of cannulated ruminants was approved in 2013 by the Unioeste Animal Use Ethics Committee (Protocol 35/13).

Location

This study was divided into two experiments conducted at Universidade Estadual do Oeste do Paraná (Unioeste), Marechal Cândido Rondon - Brazil. In the first experiment, the wet dehydration process of WBG was evaluated. The second study investigated the influence of storage period on the nutritional and microbiological quality of the DBG.

Experiment 1

A total of 6,000 kg WDG was acquired from a brewery located in Toledo, Paraná- Brazil, with 206 g/kg dry matter (DM) content. The drying of the residue was conducted under the sun by uniformly distributing it on a concrete floor, forming a layer of approximately 1.5 cm, with a density of 0.998 kg of dry matter/m². The material was scattered and turned every hour. The WBG remained dispersed for dehydration for two-day and one-night period periods, totaling 31 hours. The residue remained exposed to the sun for 16 hours, and only those hours were considered for the dehydration rate due to the presence of sunlight. The period of sunlight exposure ranged from the moment that sun's rays began to directly fall on the brewer's grains until sunlight ended in the evening. At night, the residue was covered with a black tarpaulin to prevent reabsorption of moisture.

A completely randomized experimental design was used to determine dehydration. The treatments consisted of collection times during drying, with 10 repetitions. The sampling times

were: WBG was considered as time 0 and 1, 2, 4, 6, 8, 10, 12, 14, and 16 hours of dehydration, with the schedule of 9:00, 10:00, 11:00, 13:00, and 15:00 on the first day of drying, and at 8:00, 10:00, 12:00, 14:00, and 16:00 on the second day of drying. The particle size of the collected material was from 3 to 8 mm. The material was placed in paper bags and pre-dried in an oven at 55 °C for 72 hours to allow subsequent grinding. Afterwards, it was ground in a knife mill (Tecnal model TE 650/1, Tecnal Laboratory Equipment, Piracicaba, SP, Brazil) and analyzed for DM content according to AOAC (1990; method 934.01). During sampling, temperature, relative air humidity, solar radiation, and wind speed during the drying days were obtained from the Meteorological Station of Marechal Cândido Rondon, Paraná, Brazil.

Statistical analysis

Normality of the variables was verified using the Shapiro-Wilk test. Dehydration data were analyzed by PROC REG of SAS software, version 9.0 (SAS Institute Inc., Cary, NC, USA; 2002) testing linear and quadratic models. Significance was declared as $p < 0.05$.

Experiment 2

After dehydration under the sun, a second test was carried out with the purpose of evaluating possible changes caused by storage time on the DBG product. For this, a completely randomized design with nine storage times was used.

After dehydration, the DBG was collected, packed in polypropylene braided bags (Copagril, Marechal Cândido Rondon, PR, BR) and stored in a 15 x 20 m covered shed. Residue samples collected after dehydration were considered as 0 storage day. Four samples (repetitions) were collected for 10, 20, 30, 60, 90, 120, 150, and 180 days of DBG storage. Each sample was composed of 5 subsamples collected from different parts in the bag. The samples were ground in a 1 mm sieve (Star FT-80/2, Fortinox, Piracicaba, SP, BR) and analyzed for DM, ash, crude protein (CP), and ethereal extract (EE) according to AOAC

(1990) by the methods 934.01, 938.08, 981.10, and 920.85, respectively. Neutral detergent fiber was analyzed according to Van Soest *et al.* (1991) and adapted for the Ankom 2000 Fiber Analyzer (Ankom Technology, Macedon, NY, USA) and later corrected for ash and protein (NDFap). Total carbohydrates (TC), non-fibrous carbohydrates (NFC), and carbohydrate fractionation were calculated according to the equations proposed by Sniffen *et al.* (1992). Protein fractionation was performed according to Licitra *et al.* (1996).

Animals, location, and in vitro procedures

For *in vitro* procedures, ruminal fluid was collected from three cannulated Jersey steers from the Unioeste Experimental Farm. The animals were kept in Tifton grass picket (*Cynodon* sp.) receiving feed in the corn silage trough and concentrate composed of ground corn, soybean meal, DBG, and mineral mix.

In vitro gas production was obtained using the methodology of Theodorou *et al.* (1994) and modified by Mauricio *et al.* (1999) in a computerized wireless system (Ankom RF Gas Production System, Ankom Technology, Macedon, NY, USA). For this, 500 mg of sample was introduced into a glass flask (295 mL), along with 100 ml buffer solution (pH 6.8) and 25 mL rumen fluid. For the variation adjustments, flasks considered blank were incubated, containing the incubation solutions without substrate. Gas pressure produced in the flask was measured every 10 minutes for 48 hours. Accumulated gas volume was corrected for the fermented DM. To estimate ruminal fermentation kinetic parameters of each treatment, the bicompartamental logistic model was adjusted to the cumulative gas production curves as proposed by Schofield *et al.* (1994).

In vitro dry matter digestibility (IVDMD) was estimated by the technique described by Tilley and Terry (1963) and adapted by Holden (1999), using incubator model TE-150 Tecnal (Tecnal Laboratory Equipment, Piracicaba, SP, Brazil). *In vitro* crude protein digestibility (IVCPD) was

determined by the residual protein analysis after incubation. The technique of Goering and Van Soest (1970) was used for the determination of *in vitro* cell wall digestibility (IVCWD).

Microbiological analysis

For fungi analyses, 25 g of sample was added in 225 mL sterile distilled water and maintained with continuous stirring for five seconds. From this solution 1 mL was pipetted in successive dilutions of 10^{-1} to 10^{-5} using test tubes containing 9 mL distilled water. Subsequently, from the diluted extracts, surface seeding was performed on the plates using 0.1 mL inoculum per plate. The inoculum was spread on the plate with the aid of a Drigalski loop.

Potato dextrose agar (PDA, Acumedia, Lansing, MI, USA) was used as culture medium at pH 3.5 and acidified with tartaric acid (10:100 w/v; Synth, Diadema, SP, BR). The plates were incubated at 30 °C for 3-5 days. Subsequently, the fungal colonies were counted being considered only plaques with less than 100 colonies (APHA, 1992). Identification of the genus was performed according to Samson *et al.* (1996) through the observation of microscopic characteristics of the spores in each colony. Microscope slides were prepared by the scraping method, using cotton-blue as dye. Total aflatoxins was determined in a fluorimeter (Vican Series-4X fluorimeter, Vican, Miliford,

MA, USA) using the AflaTest® methodology (AflaTest Micotoxin Testing System, Vican, Miliford, MA, USA).

Statistical analysis

Normality of variables was verified using the Shapiro-Wilk test. The data of fungi genus population were transformed into Log base 10. Data were analyzed by PROC REG of SAS software, version 9.0 (SAS Institute Inc., Cary, NC, USA; 2002), testing linear and quadratic models. Significance was declared as $p < 0.05$.

Results

Experiment 1

Average ambient temperature and relative humidity during WBG dehydration were 25.6 °C and 64.2%, respectively (Table 1). Dry matter content of WBG increased linearly ($p < 0.05$) with dehydration (Figure 1).

Experiment 2

The DM content of DBG presented changes ($p < 0.05$) during storage (Figure 2). Contents of ash, CP, EE, NDFap, and NFC had no effect ($p > 0.05$) as a function of the storage times tested (Figure 3).

Table 1. Temperature of wet brewer's grains (TWBG, °C), ambient temperature (AT, °C), relative humidity (RH, %), solar radiation (SR, kJ/m²) and wind speed (WV, m/s) during the dehydration process of wet brewer's grains.

Hours of dehydration	H/day	TWBG (°C)	AT (°C)	RH (%)	SR (kJ/m ²)	WV (m/s)
0	09:00/1st	27.0	21.9	76	1047	4.1
1	10:00/1st	27.3	24.2	65	1714	5.4
2	11:00/1st	29.5	26.5	61	2131	3.8
4	13:00/1st	31.1	27.6	56	2257	1.3
6	15:00/1st	27.7	27.3	58	1746	2.2
8	08:00/2nd	24.9	20.0	84	228	3.2
10	10:00/2nd	29.4	25.6	65	1718	3.5
12	12:00/2nd	32.2	28.2	55	2644	3.2
14	14:00/2nd	32.1	27.4	59	1464	2.4
16	16:00/2nd	32.0	27.4	63	910	1.6




Figure 1. Dry matter (DM) contents during sun dehydration of brewer's grains.




Figure 2. Dry matter (DM) of dried brewer's grains (DBG), ambient temperature (AT) and relative humidity (RH) during storage period of DBG.




Figure 3. Ash content, crude protein (CP), ether extract (EE), neutral detergent fiber corrected for ash and protein (NDFap), and non-fibrous carbohydrate (NFC) of dried brewer's grains during storage.

Storage days of DBG did not influence ($p>0.05$) fractions A, B1, B2 and B3 of the protein (Table 2). Fraction C, considered indigestible, increased linearly ($p<0.05$) with storage time. Carbohydrate fractions A+B1, B2 and C had no effect with increasing storage time of DBG ($p>0.05$).

The gas volume of the fast degradation fraction (A) showed a quadratic effect as a

function of storage time, with the maximum estimated at 88 days, while degradation rate of non-fibrous carbohydrates (B) showed linear increasing effect ($p<0.05$) (Table 3). Time of colonization (C), gas volume (D), and degradation rate of the slow fraction (E) as well as total gas production (A+D) had no effect as a function of storage time ($p>0.05$). Dried brewer's grains storage time did not influence ($p>0.05$) *IVDMD*, *IVCPD*, and *IVCWD*.

Table 2. Fractionation of proteins (CP) and carbohydrates (TC) from dried brewer's grains submitted to different storage times.

Variables	Storage time (days)				SEM	p-value		R ²
	0	30	90	180		Linear	Quadratic	
Fractionation of proteins (g/kg de CP)								
A	161	141	126	134	13.3	0.208	0.194	-
B1	19.8	26.7	17.4	24.0	6.71	0.921	0.738	-
B2	454	447	445	438	14.2	0.480	0.941	-
B3	263	280	296	274	11.2	0.589	0.062	-
C ¹	101	103	113	128	5.21	0.002	0.780	0.99
Fractionation of carbohydrates (g/kg de TC)								
A+B1	145	125	111	132	11.8	0.533	0.075	-
B2	640	645	658	637	14.7	0.890	0.317	-
C	213	228	229	230	9.75	0.339	0.445	-

SEM: Standard error of the mean. Fractionation of proteins: A: soluble; B1: rapid ruminal degradation; B2: intermediate ruminal degradation; B3: slow ruminal degradation; C: indigestible. Fractionation of carbohydrates: A+B1: rapidly degradable; B2: potentially degradable fibrous carbohydrates; C: non-degradable carbohydrates. $\hat{Y} = 99.9755 + 0.1540x$.

Storage time did not affect DBG fungal population ($p>0.05$). The most common genotypes of fungi found in all evaluation times were *Cladosporium sp.* and *Fusarium sp.*

(Figure 4a). Total aflatoxin concentration (Figure 4b) showed a quadratic effect as a function of storage time ($p<0.05$) with the maximum estimated at 96 days.

Table 3. *In vitro* cumulative gas production and *in vitro* digestibility of dried brewer's grains submitted to different storage times.

Variables	Storage time (days)				SEM	p-value		R ²
	0	30	90	180		Linear	Quadratic	
Gas production kinetics (mL/100 mg fermented dry matter)								
A (mL) ¹	6.46	7.41	8.42	6.32	0.503	0.696	0.010	0.99
B (/h) ²	0.13	0.14	0.15	0.18	0.009	0.008	0.623	0.98
C (h)	0.68	1.46	0.76	1.67	0.371	0.210	0.590	-
D (mL)	10.4	9.08	10.4	11.1	0.714	0.199	0.491	-
E (/h)	0.031	0.029	0.029	0.034	0.0014	0.061	0.100	-
A+D (mL)	16.9	16.4	18.8	17.4	0.77	0.330	0.170	-
<i>In vitro</i> digestibility (g/kg dry matter)								
IVDMD	690	717	717	724	14.2	0.199	0.471	-
IVCPD	935	925	922	930	8.79	0.840	0.330	-
IVCWD	600	643	650	645	15.3	0.136	0.124	-

SEM: Standard error of the mean; A: gas volume from the fast degraded carbohydrate fraction (non-fibrous carbohydrate); B: rate of degradation of non-fibrous carbohydrate; C: latency or colonization time; D: gas volume from the slow-degraded carbohydrate fraction (fibrous carbohydrate); E: rate of degradation of fibrous carbohydrate; IVDMD: *in vitro* dry matter digestibility; IVCPD: *in vitro* crude protein digestibility; IVCWD: *in vitro* cell wall digestibility; ¹ $\hat{Y} = 6.4038 + 0.0442x - 0.00025x^2$; ² $\hat{Y} = 0.1337 + 0.00025x$.




Figure 4. Occurrence of fungal genus (a) and total aflatoxin (AF) (b) of dried brewer's grains (DBG) submitted to different storage times.

Discussion

Experiment 1

Average ambient temperature was 25.6 °C during dehydration, which may be considered low and suggests that wind speed, solar rays, and low relative air humidity also contributed to the drying of the material. Brewer's grains may reduce protein availability during dehydration, as drying temperature increases due to condensation of some proteins with indigestible cell wall materials (Pereira *et al.*, 1998). However, as it was dehydrated under the sun, the temperature was low so no change in composition was expected (Pereira *et al.*, 1998).

Dehydration of WBG showed increasing linear effect on DM content of the material. The DM content increased 49.82 g/kg per hour of drying. With this dehydration rate WBG resulted in a DM matter content of 949.85 g/kg at the end of the dehydration period. The DM obtained at the end of dehydration was sufficient to ensure the preservation of the residue, and exceeds the values obtained by Schone *et al.* (2016), who observed DM content around 880 g/kg when dehydrating brewer's grains in the sun in a 2 cm-thick layer. Industrial dehydration of WBG results in DM content between 900 g/kg (Aliyu and Bala, 2011) and 923 g/kg (Dihman *et al.*, 2003).

Experiment 2

The DM content of DBG presented changes during storage. This behavior is associated to air humidity at each storage time. Sun dehydration was conducted during the month of April, so collection for the 10, 20, 30, and 60 storage days occurred when air humidity was high, causing reabsorption of atmospheric moisture by the residue. The reduction of air humidity that occurred in the following months was responsible for the new increase in DM content of the material.

Storage time did not influence the chemical composition of DBG; however, some oscillations observed in nutrient concentration were mostly

due to the heterogeneity of the product, which is caused by grain variety, harvest time, maceration conditions and malting (Robertson *et al.*, 2010), type of beer produced, cereals added together with barley such as corn, wheat and rice (Mussatto *et al.*, 2006), production method, and waste recovery (Westendorf *et al.*, 2014). Additionally, DBG does not have a defined quality standard for commercialization because it is an agroindustry residue.

Storage days did not influence carbohydrate fractionation of fractions A, B1, B2, and B3 of the protein, but fraction C increased with storage time. For each day of storage this fraction increased 0.16 g/kg of CP, so part of the previously digestible protein was bound to the cell wall compounds, becoming unavailable.

Gas volume of the fast degradation fraction (A) showed a quadratic effect as a function of storage time, while degradation rate of non-fibrous carbohydrates (B) showed a linear increasing effect. Although these variables were significant, the results may also have occurred due to heterogeneity of the product. Gas volume of the rapid degradation fraction was low for all storage times since soluble carbohydrates and barley starch are largely solubilized during the malting and brewing stage, generating low-grade brewer's grains of fermentable carbohydrates.

Gas volume of the slow fraction (D), total gas production (A+D), and *in vitro* digestibility of DM, CP and cell wall had no effect as a function of storage time. High gas production of the slow degradation fraction occurred due to high concentration of fibrous carbohydrates, since DBG is mainly composed of barley, which is rich in cellulose, non-cellulosic polysaccharides, and lignin (Mussatto *et al.*, 2006).

Fungal population of DBG was not altered with increasing storage time. The most common genotypes of fungi found in all evaluation times were *Cladosporium* spp. and *Fusarium* spp. Genus *Rhizopus* spp. was the lowest in all storage times, while genus *Phoma* spp. was found only after 30 days of storage. Keller *et al.* (2012) obtained larger populations of

Cladosporium spp., *Aspergillus* spp. and *Mucor* spp., but also less frequently isolated fungi *Eurotium* spp., *Penicillium* spp., *Alternaria* spp., and *Fusarium* spp. Gonzalez-Pereyra *et al.* (2011) observed greater contamination by *Fusarium* spp. (30%), *Mucor* spp. (30%), and *Aspergillus* spp. (27.3%). Other genera, such as *Penicillium* spp., *Cladosporium* spp., *Geotrichum* spp., and *Alternaria* spp. were also isolated.

Among the main genera of fungi observed, it is noticeable that *Fusarium* and *Cladosporium* are field fungi, which attack cereals during grain development, causing damage before harvest (Pezzini *et al.*, 2005; Hofer *et al.*, 2016). In addition, genera *Aspergillus* and *Penicillium*, considered the main post-harvest fungi (Morcia *et al.*, 2013), were not detected, allowing to infer that most fungal populations present in DBG comes from contamination within the field, and the storage conditions were favorable for the preservation of the product. This statement was corroborated by Simas *et al.* (2007), who also pointed out that *Aspergillus* and *Penicillium* develop under inadequate storage conditions. When evaluating wet brewer's grains storage, these authors observed the presence of *Aspergillus* spp. as the most frequent genus (42.5%), followed by *Mucor* spp. (32.5%), *Rhizopus* spp. (32.5%), *Penicillium* spp. (7.5%), and *Fusarium* spp. (2.5%).

Total aflatoxin concentration showed a quadratic effect as a function of storage time. Values ranged from 35 to 45.5 µg aflatoxin/kg DBG, but remained within the maximum tolerance limits of 50 µg/kg for animal feed and concentrate (Brasil, 1988). Simas *et al.* (2007) analyzed 80 samples of brewer's grains finding that 33.75% of the samples tested positive for aflatoxin, with values between 1 and 3 µg/kg. Gonzalez-Pereyra *et al.* (2011) detected concentrations of 19 to 44.52 µg aflatoxin B1/kg in 18% of brewer's grains samples tested, while Keller *et al.* (2012) detected concentrations between 10 and 47 µg aflatoxin B1/kg in 4% of the samples.

Aflatoxin is produced by the fungi *Aspergillus* sp., but this genus was not detected in the present study. Gonzalez-Pereyra *et al.* (2011) pointed out that detection of mycotoxins in a given substrate implies contamination by toxigenic species that may or may not be present at sampling time. Therefore, it is believed that contamination by *Aspergillus* sp. can occur through the barley grains during the field phase or during malting (Morcia *et al.*, 2013). During brewing, subsequent to malting, the mash is heated to 78 °C for several hours (Gupta *et al.* 2010), which may have caused the disappearance of *Aspergillus* sp., thus maintaining the mycotoxins it produces, since they are thermostable (Terzi *et al.*, 2014). Therefore, the absence of *Aspergillus* spp. and the maintenance of aflatoxin concentration in DBG within the acceptable limits during the storage time allow us to affirm that the storage conditions were favorable for the preservation of the product.

In conclusion, dehydration of WBG for a total period of 31 hours, with 16 hours of direct exposure to the sun and a density of 0.998 kg DM/m², is efficient to preserve the material and makes its storage feasible. Storing DBG for 180 days does not compromise its nutritional and microbiological quality. More research is needed on the use of technologies that provide renewable and low-cost energy, such as solar radiation, for the dehydration of this by-product on a large scale, considering its potential long-term storage for use in ruminant feed.

Declarations

Acknowledgments

The authors would like to thank Cervejaria Colônia, Toledo, PR, Brazil for their collaboration in the study.

Funding

This study was financed by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Grant - PNPd 2891/2010.

Conflict of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

Author contributions

Andressa Faccenda: design and conception of the study, data collection, analysis and data interpretation, writing, review and editing of the paper. Maximiliane A Zambom: design and conception of the study, project administration, funding acquisition, critical reading and review of the paper. André S Avila: data collection, analysis and data interpretation, writing and review of the paper. Deise D Castagnara: design and conception of the study, data collection and critical reading of the paper. Ricardo Dri: data collection, critical reading and review of the paper. Maria L Fischer: data collection, critical reading and review of the paper. Rodrigo C R Tinini: data collection, critical reading and review of the paper. Jéssica G Dessbesell: data collection, critical reading and review of the paper. Ana-Ruth E Almeida: data collection, critical reading and review of the paper. Kleves V de Almeida: data collection, critical reading and review of the paper.

References

- Aliyu S, Bala M, Brewer's spent grain: A review of its potentials and applications. *African J Biotechnol* 2011; 10(3):324–331.
- Association of Official Analytical Chemists (AOAC). Official methods of analysis of Association of Official Analytical Chemists. 15th ed. Arlington: Association of Official Analytical Chemists; 1990.
- American Public Health Association (APHA). Compendium of methods for the microbiological examination of foods. 3rd ed. Washington: American Public Health Association; 1992.
- Brasil. Ministério da Agricultura. Portaria n. 07, de 09 de novembro de 1988. Fixa padrões de tolerância para aflatoxinas em alimentos para consumo animal: matérias primas e rações. *Diário Oficial da União*, p.21.968, 09 nov. 1988. Seção I. 1988.
- Dhiman TR, Bingham HR, Radloff HD. Production response of lactating cows fed dried versus wet brewers' grain in diets with similar dry matter content. *J Dairy Sci* 2003; 86(9):2914–2921. DOI: [https://doi.org/10.3168/jds.S0022-0302\(03\)73888-0](https://doi.org/10.3168/jds.S0022-0302(03)73888-0)
- Goering HK, Van Soest PJ. Forage fiber analysis (apparatus, reagents, procedures, and some applications). *Agriculture handbook*, 379. Washington DC: Agricultural Research Service; 1970.
- Gonzalez-Pereyra ML, Rosa CAR, Dalcero AM, Cavaglieri LR. Mycobiota and mycotoxins in malted barley and brewer's spent grain from Argentinean breweries. *Lett Appl Microbiol* 2011; 53(6):649–655. DOI: <https://doi:10.1111/j.1472-765X.2011.03157.x>
- Gupta M, Abu-Ghannam N, Gallagher E, Barley for brewing: Characteristic changes during malting, brewing and applications of its by-products. *Compr Rev Food Sci Food Saf* 2010; 9(3):318–328. DOI: <https://doi.org/10.1111/j.1541-4337.2010.00112.x>
- Hofer K, Geißinger C, König C, Gastl M, Hückelhoven R, Heß M, Coleman AD. Influence of Fusarium isolates on the expression of barley genes related to plant defense and malting quality. *J Cereal Sci* 2016; 69:17–24. DOI: <https://doi.org/10.1016/j.jcs.2016.02.005>
- Holden LA. Comparison of methods of *in vitro* dry matter digestibility for ten feeds. *J Dairy Sci* 1999; 82(8):1791–1794. DOI: [https://doi.org/10.3168/jds.S0022-0302\(99\)75409-3](https://doi.org/10.3168/jds.S0022-0302(99)75409-3)
- Keller LAM, Pereyra CM, Cavaglieri LR, Dalcero AM, Rosa CAR. Fungi and mycotoxins from pre- and poststorage brewer's grain intended for bovine intensive rearing. *ISRN Vet Sci* 2012; 2012:396590. DOI: <https://doi.org/10.5402/2012/396590>

- Licitra G, Hernandez TM, Van Soest PJ. Standardization of procedures for nitrogen fractionation of ruminant feeds. *Anim Feed Sci Technol* 1996; 57(4):347–358. DOI: [https://doi.org/10.1016/0377-8401\(95\)00837-3](https://doi.org/10.1016/0377-8401(95)00837-3)
- Mauricio RM, Mould FL, Dhanoa MS, Owen E, Channa KS, Theodorou MK. A semi-automated *in vitro* gas production technique for ruminant feedstuff evaluation. *Anim Feed Sci Technol* 1999; 79(4):321–330. DOI: [https://doi.org/10.1016/S0377-8401\(99\)00033-4](https://doi.org/10.1016/S0377-8401(99)00033-4)
- Morcía C, Rattotti E, Stanca AM, Tumino G, Rossi V, Ravaglia S, Germeier CU, Herrmann M, Polisenka I, Terzi V. Fusarium genetic traceability: Role for mycotoxin control in small grain cereals agro-food chains. *J Cereal Sci* 2013; 57(2):175–182. DOI: <https://doi.org/10.1016/j.jcs.2012.09.016>
- Mussatto SI, Dragone G, Roberto IC. Brewers' spent grain: Generation, characteristics and potential applications. *J Cereal Sci* 2006; 43(1):1–14. DOI: <https://doi.org/10.1016/j.jcs.2005.06.001>
- Negesse T, Makkar HPS, Becker K. Nutritive value of some non-conventional feed resources of Ethiopia determined by chemical analyses and an *in vitro* gas method. *Anim Feed Sci Technol* 2009; 154(3-4):204–217. DOI: <https://doi.org/10.1016/j.anifeedsci.2009.09.010>
- Pereira JC, Carro MD, Gonzalez J, Alvir MR, Rodriguez CA. Rumen degradability and intestinal digestibility of brewers' grains as affected by origin and heat treatment and of barley rootlets. *Anim Feed Sci Technol* 1998; 74(2):107–121. DOI: [https://doi.org/10.1016/S0377-8401\(98\)00168-0](https://doi.org/10.1016/S0377-8401(98)00168-0)
- Pezzini V, Valduga E, Cansian RL. Incidence of fungi and mycotoxins in stored corn under different conditions. *Rev Inst Adolfo Lutz* 2005; 64(1):91–6.
- Robertson JA, I'Anson KJA, Treimo J, Faulds CB, Brocklehurst TF, Eijsink VGH, Waldron KW. Profiling brewers' spent grain for composition and microbial ecology at the site of production. *LWT Food Sci Technol* 2010; 43(6):890–896. DOI: <https://doi.org/10.1016/j.lwt.2010.01.019>
- Samson RA, Hoekstra ES, Frisvad JC, Filtenborg O. Introduction to food-borne fungi. 5th ed. The Netherlands: Centralalbureau voor Schimmelcultures; 1996.
- Schofield P, Pitt RE, Pell AN. Kinetics of fiber digestion from *in vitro* gas production. *J Anim Sci* 1994; 72(11):2980–2991. DOI: <https://doi.org/10.2527/1994.72112980x>
- Schone RA, Zambom MA, Nunes RV, Fernandes T, Frank R, Oliveira TMM, Castagnara DD. Secagem de resíduo de cervejaria em camadas de diferentes espessuras. *Sci Agrar Parana* 2016; 15(2):127–131. DOI: <http://dx.doi.org/10.18188/1983-1471/sap.v15n2p127-131>
- Simas MMS, Botura MB, Correa B, Sabino M, Mallmann CA, Bitencourt TCBS, Batatinha MJM. Determination of fungal microbiota and mycotoxins in brewers grain used in dairy cattle feeding in the State of Bahia, Brazil. *Food Control* 2007; 18(5):404–408. DOI: <https://doi.org/10.1016/j.foodcont.2005.11.007>
- Sniffen CJ, O'Connor JD, Van Soest PJ, Fox DG, Russell J. A net carbohydrate and protein system for evaluating cattle diets: II Carbohydrate and protein availability. *J Anim Sci* 1992; 70(11):3562–3577. DOI: <https://doi.org/10.2527/1992.70113562x>
- Statistical Analysis System (SAS). SAS user's guide. Version 9.0. Cary, NC, USA: SAS Institute Inc., 2002.
- Terzi V, Tumino G, Stanca AM, Morcía C. Reducing the incidence of cereal head infection and mycotoxins in small grain cereal species. *J Cereal Sci* 2014; 59(3):284–293. DOI: <https://doi.org/10.1016/j.jcs.2013.10.005>
- Theodorou MK, Williams BA, Dhanoa MS, Mcallan AB, France J. A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Anim Feed Sci Technol* 1994; 48(3-4):185–197. DOI: [https://doi.org/10.1016/0377-8401\(94\)90171-6](https://doi.org/10.1016/0377-8401(94)90171-6)

Tilley JMA, Terry RA. Two-stage technique for the *in vitro* digestion of forage crops. *Grass Forage Sci* 1963; 18(2):104–111. DOI: <https://doi.org/10.1111/j.1365-2494.1963.tb00335.x>

Van Soest PJ, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci* 1991; 74(10):3583–3597. DOI: [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)

Wang B, Luo Y, Myung KH, Liu JX. Effects of storage duration and temperature on the chemical composition, microorganism density, and *in vitro* rumen fermentation of wet brewers grains. *Asian Australas J Anim Sci* 2014; 27(6):832–840. DOI: <https://doi.org/10.5713/ajas.2013.13668>

Westendorf ML, Wohlt JE, Sniffen CJ, Ward RT. Nutrient content of brewers grains produced at a commercial brewery: Variation in protein/nitrogen, fiber, carbohydrate, fat, and minerals. *Prof Anim Sci* 2014; 30(4):400–406. DOI: <https://doi.org/10.15232/pas.2013-01272>