

## Subject index for Volume 33, 2020

- Additives; 33(4): 264-272.
- Aged meat; 33(2): 96-109.
- Amino acid supplementation; 33(4): 239-251.
- Animal stress; 33(1): 6-31.
- Annatto seed; 33(2): 110-120.
- Antibiotic alternatives; 33(3): 195-201.
- Antibiotics replacement; 33(3): 195-201.
- Antimicrobial peptide; 33(1): 32-43.
- Antioxidants; 33(1): 6-31.
- Aquaculture; 33(1): 32-43.
- Astrocaryum murumuru*; 33(4): 264-272.
- Product; 33(4): 264-272.
- Bacterial resistance; 33(3): 195-201.
- Beef cattle; 33(2): 134-143.
- Beef; 33(2): 121-133.
- Bioavailability; 33(4): 217-227.
- Biofloc; 33(1): 32-43.
- Bixa orellana*; 33(2): 110-120.
- Bovine serum albumin; 33(3): 172-181.
- Bovine; 33(2): 121-133.
- Brilliant cresyl blue; 33(3): 172-181.
- Broiler; 33(3): 195-201.
- Calves; 33(1): 6-31.
- Campylobacter fetus venerealis*; 33(3): 149-158.
- Carcass characteristics; 33(4): 228-238.
- Carcass traits; 33(2): 96-109; 33(4): 217-227.
- Carcass yield; 33(2): 96-109.
- Cassava; 33(4): 204-216.
- Castration; 33(2): 96-109.
- Cattle growth; 33(1): 6-31.
- Cattle; 33(2): 121-133; 33(3): 149-158.
- Cereal grains; 33(4): 204-216.
- Chili pepper; 33(3): 195-201.
- Citrus pulp; 33(4): 204-216.
- Coccidiosis challenge; 33(3): 195-201.
- Complex diet; 33(3): 159-171.
- Conservation; 33(3): 182-194.
- Conservation; 33(4): 264-272.
- Corn; 33(2): 121-133; 33(4): 204-216.
- Coturnix japonica*; 33(2): 110-120.
- Cow milk; 33(4): 252-263.
- Cow; 33(4): 204-216.
- Creep feed; 33(3): 159-171.
- Culture; 33(1): 5-15.
- Curcumin; 33(1): 6-31.
- Cysteine; 33(4): 228-238.
- Dairy cattle; 33(4): 204-216.
- Devil fish; 33(4): 252-263.
- Diet complexity; 33(3): 159-171.
- dose-response; 33(4): 239-251.

- Dual purpose cows; 33(4): 252-263.
- Ear canal; 33(1): 5-15.
- Economic return; 33(4): 239-251.
- Effective population size; 33(1): 44-59.
- Effluents; 33(4): 264-272.
- Egg quality; 33(2): 110-120.
- Egg; 33(2): 110-120.
- Eimeria*; 33(1): 6-31.
- Elephant grass; 33(4): 264-272.
- Energy source; 33(4): 204-216.
- Ensilage; 33(3): 182-194.
- Ensiled grain; 33(2): 121-133.
- Ewes; 33(2): 81-95.
- Feed byproducts; 33(2): 110-120.
- Feed cost; 33(4): 239-251.
- Feed efficiency; 33(4): 204-216.
- Feed supplement; 33(4): 252-263.
- Fermentation; 33(3): 182-194; 33(4): 264-272.
- Fetal bovine serum; 33(3): 172-181.
- Fiber; 33(3): 182-194.
- Fish silage; 33(4): 252-263.
- fodder; 33(4): 264-272.
- Forage; 33(3): 182-194.
- Forage; 33(4): 264-272.
- Gene expression; 33(1): 32-43.
- Gene flow; 33(1): 44-59.
- Generation interval; 33(1): 44-59.
- Genetic correlation; 33(1): 60-70.
- Genetic diversity; 33(1): 44-59.
- Genetic drift; 33(1): 44-59.
- Genetic effects; 33(2): 134-143.
- Genetic parameters; 33(1): 60-70.
- Genotype-environment interaction; 33(2): 134-143.
- Gestation; 33(2): 81-95.
- Global warming; 33(2): 134-143.
- Grain; 33(2): 121-133.
- gross margin; 33(4): 239-251.
- Growth performance; 33(3): 159-171; 33(4): 217-227; 33(4): 228-238; 33(4): 239-251.
- Haemonchus contortus*; 33(2): 81-95.
- Heat stress; 33(2): 134-143.
- Heifers; 33(3): 149-158.
- Hematology; 33(2): 81-95.
- Heritability; 33(1): 60-70.
- Histopathology; 33(3): 149-158.
- Holstein; 33(1): 60-70.
- Horses; 33(1): 5-15.
- Identification; 33(1): 5-15.
- In vitro embryo production; 33(3): 172-181.
- Inbreeding; 33(1): 44-59.
- Japanese quail; 33(2): 110-120.
- Laboratory diagnosis; 33(2): 81-95.
- Lamb; 33(2): 96-109.
- Lameness; 33(4): 204-216.
- Lipid oxidation; 33(2): 96-109.
- Litter performance; 33(3): 159-171.

- Losses; 33(3): 182-194.
- Lymphocytic inflammation; 33(3): 149-158.
- Maize; 33(2): 121-133.
- Malassezia*; 33(1): 5-15.
- Marbling; 33(2): 121-133.
- Mastitis; 33(1): 60-70.
- Meat color; 33(2): 121-133.
- Meat maturation; 33(2): 96-109.
- Meat quality; 33(2): 121-133; 33(2): 96-109.
- Meat; 33(2): 96-109.
- Meiotic competence; 33(3): 172-181.
- Methionine; 33(4): 228-238.
- Milk quality; 33(1): 60-70.
- Milk quality; 33(4): 252-263.
- Milk yield; 33(1): 60-70.
- Multi-trait model; 33(1): 60-70.
- Murumuru meal; 33(4): 264-272.
- Nellore; 33(2): 121-133.
- Nematodes; 33(2): 81-95.
- Non-fibrous carbohydrates; 33(4): 204-216.
- Normal microbiota; 33(1): 5-15.
- Nursery diet; 33(3): 159-171.
- Nutrient digestibility; 33(2): 110-120.
- Nutrient requirements; 33(4): 217-227.
- Nutritional requirements; 33(4): 228-238.
- Oleoresins; 33(3): 195-201.
- Oocyte maturation; 33(3): 172-181.
- Oocyte quality; 33(3): 172-181.
- Oocyte recovery; 33(3): 172-181.
- Oviduct; 33(3): 149-158.
- Ovine; 33(2): 96-109.
- Ovis aries*; 33(2): 81-95.
- Parasites; 33(2): 81-95.
- Parasitology; 33(1): 6-31.
- PCR; 33(1): 5-15.
- Pedigree; 33(1): 44-59.
- Penaeidin; 33(1): 32-43.
- Pennisetum purpureum*; 33(4): 264-272.
- Performance; 33(3): 195-201.
- Peripartum; 33(2): 81-95.
- Phytogenic supplements; 33(3): 195-201.
- Pig; 33(4): 217-227.
- Piglet; 33(3): 159-171.
- Pigs; 33(4): 228-238.
- Plant pigment; 33(2): 110-120.
- Population structure; 33(1): 44-59.
- Post-mating infection; 33(3): 149-158.
- Poultry nutrition; 33(2): 110-120.
- Poultry production; 33(3): 195-201.
- Poultry; 33(4): 239-251.
- Prewaning growth; 33(2): 134-143.
- Probability of gene origin; 33(1): 44-59.
- Protected methionine; 33(4): 217-227.
- Protein source; 33(4): 252-263.
- Protein supplement; 33(4): 252-263.

- Protein; 33(3): 182-194; 33(4): 217-227; 33(4): 228-238.
- Protozoa resistance; 33(3): 195-201.
- Random regression; 33(2): 134-143.
- Ras; 33(1): 32-43.
- Reaction-norm; 33(2): 134-143.
- Retinta breed; 33(2): 134-143.
- Romosinuano cattle; 33(1): 44-59.
- Rumen degradability; 33(3): 182-194.
- Ruminal acidosis; 33(4): 204-216.
- Ruminant; 33(3): 182-194.
- Santa Ines breed; 33(2): 96-109.
- Santa Ines; 33(2): 81-95.
- Sarcomere; 33(2): 121-133.
- Segmented model; 33(4): 239-251.
- Shear force; 33(2): 96-109.
- Sheep; 33(2): 81-95; 33(2): 96-109.
- Shrimp immune system; 33(1): 32-43.
- Shrimp; 33(1): 32-43.
- Silage; 33(2): 121-133; 33(3): 182-194; 33(4): 264-272.
- Silo; 33(4): 264-272.
- Skin; 33(1): 5-15.
- Somatic cell score; 33(1): 60-70.
- Sorghum; 33(2): 121-133; 33(4): 204-216.
- Stability; 33(4): 264-272.
- Starch; 33(4): 204-216.
- Subcutaneous fat; 33(2): 121-133.
- Suckling; 33(3): 159-171.
- Sugarcane; 33(3): 182-194.
- Sulfur amino acids; 33(4): 228-238.
- Supplementation; 33(1): 6-31.
- Swine nutrition; 33(4): 228-238.
- Swine; 33(4): 217-227.
- Synthetic amino acids; 33(4): 217-227.
- THI; 33(2): 134-143.
- Turmeric; 33(3): 195-201.
- Urea; 33(3): 182-194; 33(4): 217-227.
- Uterus; 33(3): 149-158.
- Vagina; 33(3): 149-158.
- Variance components; 33(1): 60-70.
- Venereal disease; 33(3): 149-158.
- Vitamin E; 33(2): 96-109.
- Weaning; 33(3): 159-171.
- Yeast; 33(1): 5-15.
- Yolk pigmentation; 33(2): 110-120.
- Zebu; 33(2): 121-133.
- $\alpha$ -tocopherol; 33(2): 96-109.