

Subject index for Volume 34, 2021

- Aberdeen angus; 34(4): 278-290.
- Additive; 34(2): 95-104.
- Alternative feedstuff; 34(1): 51-62; 34(2): 105-116.
- Alternative feedstuffs; 34(1): 73-81; 34(4): 267-277.
- Alternative production; 34(2): 84-94.
- Anglonubian; 34(1): 40-50.
- Animal performance; 34(2): 105-116.
- Animal production; 34(1): 40-50.
- Antibiotic; 34(1): 51-62.
- Antimicrobial resistance; 34(1): 51-62.
- Antimicrobial; 34(1): 63-72.
- Aquaculture; 34(1): 29-39.
- B-hydroxybutyrate; 34(4): 291-304.
- Bacteria; 34(1): 18-28.
- Bacterial resistance; 34(1): 51-62.
- Barium selenate; 34(2): 145-153.
- Barley; 34(4): 254-266.
- Barr chromatin; 34(2): 126-144.
- Bayesian inference; 34(1): 40-50; 34(3): 166-176.
- Beef; 34(2): 117-125.
- Beef cattle; 34(4): 278-290.
- Beef cuts; 34(2): 117-125.
- Beef quality; 34(2): 117-125.
- Bio-control; 34(2): 154-161.
- Biological intervals; 34(2): 126-144.
- Biotype; 34(4): 278-290.
- Bitch; 34(3): 231-239.
- Blood cells; 34(2): 126-144.
- Blood creatinine; 34(4): 291-304.
- Blood glucose; 34(4): 291-304.
- Blood urea nitrogen; 34(4): 291-304.
- Blood; 34(2): 126-144.
- BLV; 34(3): 177-188.
- Body condition score; 34(4): 291-304.
- Bovine leukemia virus; 34(3): 177-188.
- Bovine meat; 34(2): 117-125.
- Bovine; 34(1): 18-28; 34(2): 145-153; 34(3): 224-230.
- Bovine; 34(4): 278-290.
- Breeders; 34(1): 29-39.
- Breeding value; 34(3): 166-176.
- Brewer's grains; 34(4): 254-266.
- Brewery by-product; 34(4): 254-266.
- Broiler chicken; 34(3): 189-199; 34(2): 84-94.
- Broilers; 34(2): 95-104; 34(3): 189-199.
- Broodfish; 34(1): 29-39.
- Broodstock; 34(1): 29-39.
- Brucellosis; 34(3): 224-230.
- Brycon orbignyanus*; 34(1): 29-39.
- BTB; 34(3): 224-230.
- Buffaloes; 34(3): 166-176.

- Bull; 34(3): 200-211.
- Calcium oxide; 34(1): 51-62.
- Calves; 34(2): 145-153.
- Canine; 34(3): 231-239.
- Canines; 34(2): 126-144.
- Canonical correlation analysis; 34(2): 117-125.
- Carcass; 34(1): 40-50; 34(1): 51-62.
- Carcinoma; 34(3): 231-239.
- Cartilage; 34(3): 212-223.
- Catch-up growth; 34(3): 189-199.
- Cattle breed; 34(4): 278-290.
- Cattle meat; 34(2): 117-125.
- Cattle productivity; 34(1): 5-17.
- Cattle; 34(3): 177-188.
- Cattle; 34(4): 278-290.
- Cell differentiation; 34(3): 212-223.
- Cellular count; 34(2): 126-144.
- Chewing; 34(4): 267-277.
- Chicken; 34(2): 95-104.
- Chondrocytes; 34(3): 212-223.
- Chondrogenesis; 34(3): 212-223.
- Chondrogenic differentiation; 34(3): 212-223.
- Chromatin; 34(2): 126-144.
- Coffee byproducts; 34(1): 51-62.
- Coffee hulls; 34(1): 51-62.
- Collagen; 34(3): 212-223.
- Compensatory growth; 34(3): 189-199.
- Conservation; 34(4): 254-266.
- Corneal ulceration; 34(1): 18-28.
- Corticosteroids; 34(3): 212-223.
- (co)variance; 34(1): 40-50.
- Cows; 34(2): 145-153; 34(3): 177-188.
- Creatine; 34(2): 95-104.
- Cryopreservation; 34(3): 200-211.
- Cryoprotectant; 34(3): 200-211.
- Dairy cattle; 34(1): 18-28; 34(1): 5-17.
- Dairy cows; 34(4): 305-315.
- Dairy industry; 34(4): 241-253.
- Dairy production; 34(1): 5-17.
- Dairy products; 34(4): 241-253.
- Dairy; 34(4): 241-253.
- Digestibility; 34(1): 73-81.
- Digestibility; 34(2): 84-94; 34(2): 105-116.
- Digestibility; 34(4): 254-266.
- Digestibility; 34(4): 305-315.
- Digestive development; 34(2): 84-94.
- Disease surveillance; 34(3): 224-230.
- Dog; 34(2): 126-144.
- Dried brewer's grains feed; 34(4): 254-266.
- Dry matter digestibility; 34(4): 316-323.
- EBL; 34(3): 177-188.
- Echography; 34(3): 231-239.
- Egg yolk; 34(3): 200-211.
- Eggs; 34(2): 154-161.
- Energy; 34(1): 73-81.
- Ensilage; 34(4): 305-315.

- Ensiling; 34(4): 305-315.
- Enterobacteria; 34(1): 51-62.
- Environment; 34(3): 166-176.
- Environmental gradient; 34(3): 166-176.
- Enzootic bovine leukosis; 34(3): 177-188.
- Enzyme; 34(4): 316-323.
- Epidemiology; 34(3): 177-188; 34(3): 224-230.
- Erythroblasts; 34(2): 126-144.
- Erythrogram; 34(2): 126-144.
- Escherichia coli; 34(1): 51-62.
- Ethology; 34(4): 267-277.
- Extender; 34(3): 200-211.
- Eye infection; 34(1): 18-28.
- Feed additive; 34(2): 95-104.
- Feed additives; 34(4): 305-315.
- Feed by-product; 34(1):73-81; 34(4): 254-266.
- Feed efficiency; 34(4): 267-277.
- Feed fines; 34(2): 84-94.
- Feed ingredient; 34(1): 51-62.
- Feed intake; 34(2): 105-116.
- Feed restriction; 34(3): 189-199.
- Feeding; 34(3): 189-199.
- Feedlot; 34(4): 267-277.
- Feedstuffs; 34(4): 316-323.
- Fertilization; 34(1): 29-39.
- Fiber degradation; 34(1): 51-62.
- Fiber; 34(1): 51-62.
- Financial management; 34(1): 5-17.
- Fish; 34(1): 29-39.
- Food safety; 34(2): 154-161.
- Food-borne disease; 34(2): 154-161.
- Food-borne pathogens; 34(2): 154-161.
- Foodstuff; 34(2): 154-161.
- Fruit by-product; 34(2): 105-116.
- Fruit residue; 34(2): 105-116.
- Fungi; 34(4): 254-266.
- Genetic correlation; 34(1): 40-50.
- Genetic diversity; 34(1): 29-39; 34(4): 278-290.
- Genetic evaluation; 34(3): 166-176.
- Genetic parameters; 34(1): 40-50.
- Genetic structure; 34(4): 278-290.
- Genotype by environment interaction; 34(3): 166-176.
- Genotypes; 34(3): 166-176.
- Gibbs sampling; 34(1): 40-50.
- Glucocorticoid; 34(3): 212-223.
- Glutathione peroxidase; 34(2): 145-153.
- Goats' heritability; 34(1): 40-50.
- Granulosa; 34(3): 231-239.
- Growth performance; 34(3): 189-199.
- Guanidinoacetic acid; 34(2): 95-104.
- Hatching; 34(1): 29-39.
- Hematology; 34(2): 126-144.
- Hemogram; 34(2): 126-144.
- Hierarchical reaction; 34(3): 166-176.
- Histology; 34(3): 212-223; 34(3): 231-239; 34(3): 231-239.

- Hyaluronic acid; 34(3): 212-223.
- IBK; 34(1): 18-28.
- Ileal digestibility; 34(4): 316-323.
- Immunity; 34(3): 189-199.
- In situ* degradability; 34(1): 51-62.
- In vitro* assay; 34(1):73-81.
- In vitro* disappearance; 34(4): 316-323.
- Income; 34(1): 5-17.
- Incubation; 34(1): 29-39.
- Information technology; 34(3): 224-230.
- Intake; 34(4): 267-277; 34(4): 305-315.
- Isa Label; 34(2): 84-94.
- Keratoconjunctivitis; 34(1): 18-28.
- Kinship; 34(4): 278-290.
- Lamb; 34(2): 105-116.
- Low-density lipoproteins; 34(3): 200-211.
- Lytic bacteriophage; 34(2): 154-161.
- Mammary gland; 34(4): 241-253.
- Man saltbush; 34(4): 267-277.
- Mastitis etiology; 34(4): 241-253.
- Mayonnaise; 34(2): 154-161.
- Meat color; 34(2): 117-125.
- Meat quality; 34(1): 40-50.
- Meat; 34(2): 117-125.
- Mesenchymal stem cell; 34(3): 212-223.
- Metabolic efficiency; 34(4): 305-315.
- Metabolism; 34(2): 95-104.
- Microbial purines; 34(4): 305-315.
- Microbial resistance; 34(1): 51-62.
- Microsatellite markers; 34(1): 29-39.
- Microsatellites; 34(4): 278-290.
- Milk quality; 34(1): 5-17.
- Milk urea nitrogen; 34(4): 305-315.
- Milk yield; 34(4): 291-304.
- Mineral nutrition; 34(2): 145-153.
- Mitochondrial DNA; 34(4): 278-290.
- Mitochondrial introgression; 34(4): 278-290.
- Moraxella bovis*; 34(1): 18-28.
- Moraxella bovoculi*; 34(1): 18-28.
- Mortality; 34(1): 29-39.
- Multidrug resistance; 34(1): 51-62.
- Multi-resistant organism; 34(1): 51-62.
- Multivariate statistics; 34(2): 117-125.
- Muscle biochemistry; 34(2): 117-125.
- Neoplasia; 34(3): 231-239.
- Nitrogen balance; 34(4): 305-315.
- Non-starch polysaccharides; 34(4): 316-323.
- Nutrient degradability; 34(1): 51-62.
- Nutrient-dense diet; 34(3): 189-199.
- Nutrition; 34(2): 95-104; 34(3): 189-199.
- Nutritional value; 34(4): 254-266.
- Ocular bacteria; 34(1): 18-28.
- Ocular disease; 34(1): 18-28.
- Organ biometry; 34(2): 95-104.
- Organoleptic characteristics; 34(2): 117-125.
- Osteoarthritis; 34(3): 212-223.

- Ovarian cancer; 34(3): 231-239.
- Ovarian tumor; 34(3): 231-239.
- Ovary; 34(3): 231-239.
- Pellet quality; 34(2): 84-94.
- Pelleted feed; 34(2): 84-94.
- Pequi; 34(2): 105-116.
- Performance; 34(2): 84-94.
- Performance; 34(4): 305-315.
- pH; 34(2): 117-125.
- Phenotype; 34(4): 278-290.
- Pig; 34(1):73-81.
- Pigs; 34(4): 316-323.
- Pink eye; 34(1): 18-28.
- Polymorphic markers; 34(4): 278-290.
- Population genetics; 34(4): 278-290.
- Poultry; 34(2): 84-94; 34(2): 95-104; 34(3): 189-199.
- Pregnancy rate; 34(4): 291-304.
- Prevalence; 34(3): 177-188.
- Production costs; 34(1): 5-17.
- Profitability; 34(1): 5-17.
- Protein; 34(1):73-81.
- Public health; 34(1): 51-62; 34(3): 224-230.
- Rank correlations; 34(3): 166-176.
- Raw food; 34(2): 154-161.
- Raw milk; 34(4): 241-253.
- Reaction norm models; 34(3): 166-176.
- Reproductive efficiency; 34(1): 29-39.
- Reproductive system; 34(3): 231-239.
- Restocking; 34(1): 29-39.
- Risk factors; 34(3): 177-188.
- Ruminant; 34(1): 51-62; 34(2): 117-125; 34(3): 177-188.
- Ruminants; 34(4): 254-266.
- Rumination efficiency; 34(4): 267-277.
- Salmonella enterica; 34(2): 154-161.
- Selection index; 34(1): 40-50.
- Selection; 34(1): 40-50.
- Selenium; 34(2): 145-153.
- Semen; 34(3): 200-211.
- Seropositivity; 34(3): 177-188.
- Seroprevalence; 34(3): 177-188.
- Serum biochemistry; 34(2): 84-94.
- Services per conception; 34(4): 291-304.
- Sheep; 34(1): 51-62; 34(2): 105-116; 34(4): 267-277.
- Silage; 34(4): 305-315.
- Slaughter; 34(1): 51-62.
- Somatic cell count; 34(4): 241-253.
- Soy byproduct; 34(1):73-81.
- Soybean milk by-product; 34(1):73-81.
- Spawning; 34(1): 29-39.
- Spray-chilled; 34(1): 51-62.
- Spray-chilling; 34(1): 51-62.
- Storage time; 34(4): 254-266.
- Strip-spawning; 34(1): 29-39.
- Subcutaneous fat thickness; 34(2): 117-125

- Supplementation; 34(2): 145-153.
- Swine; 34(1):73-81.
- Technology adoption; 34(1): 5-17.
- Thyroid hormones; 34(2): 145-153.
- Thyroxine; 34(2): 145-153.
- Total tract digestibility; 34(4): 316-323.
- Trehalose; 34(3): 200-211.
- Triancinolonone acetone; 34(3): 212-223.
- Triiodothyronine; 34(2): 145-153.
- Trypsin inhibitor; 34(1):73-81.
- Tuberculosis; 34(3): 224-230.
- Tumor; 34(3): 231-239.
- Udder; 34(4): 241-253.
- Ultrasonography; 34(1): 40-50; 34(3): 231-239
- Ultrasound; 34(3): 231-239.
- Urea; 34(4): 305-315.
- Vaginal discharge; 34(3): 231-239.
- Veterinary pathology; 34(3): 231-239.
- Virus; 34(3): 177-188.
- Viscosupplementation; 34(3): 212-223.
- Weight gain; 34(2): 105-116.
- Weight gain; 34(4): 267-277.
- Xylanase; 34(4): 316-323.
- Zoonosis; 34(3): 224-230.