

Subject index for Volume 35, 2022

- Acaricide; 35(4): 185-190.
- ADEPPT; 35(4): 233-242.
- Age at first calving; 35(2): 68-81.
- Agriculture; 35(2): 61-67.
- Agriculture; 35(4): 185-190.
- Animal adaptation; 35(4): 217-232.
- Animal feed; 35(1): 36-48.
- Antibiotic; 35(1): 26-35.
- Antibiotic; 35(4): 191-204.
- Antimicrobial resistance; 35(1): 26-35.
- Apparent metabolizable energy; 35(1): 36-48.
- Artificial rearing; 35(3): 174-184.
- Assisted reproduction; 35(2): 118-123.
- Aversive; 35(4): 205-216.
- Bacterial resistance; 35(1): 26-35.
- Bayesian; 35(3): 125-140.
- Beneficial species; 35(4): 185-190.
- Bivariate analysis; 35(3): 125-140.
- Bos indicus*; 35(1): 3-13.
- Bos taurus*; 35(2): 68-81.
- Bos taurus*×*Bos indicus*; 35(2): 68-81.
- Bovine embryo; 35(2): 93-108.
- Bovines; 35(1): 3-13.
- Breaking strength; 35(3): 153-164.
- Broiler chicken; 35(4): 191-204.
- Broiler chickens; 35(1): 36-48.
- Broiler; 35(3): 153-164.
- Brown Swiss×Zebu; 35(2): 68-81.
- Calving interval; 35(2): 68-81.
- Caprine; 35(1): 49-59.
- Carcass yield; 35(1): 49-59.
- Carcass; 35(4): 191-204.
- Carcinogen; 35(2): 61-67.
- Castration; 35(1): 49-59.
- Cattle; 35(1): 3-13.
- Cattle; 35(4): 217-232.
- Chemical composition; 35(1): 36-48.
- Chlamydiaabortus; 35(3): 165-173.
- Chlamydial infection; 35(3): 165-173.
- Chlorpyrifos; 35(2): 61-67.
- Climate change; 35(4): 217-232.
- Cognitive disorders; 35(2): 61-67.
- Colombia; 35(2): 61-67.
- Colombia; 35(2): 82-92.
- Colombia; 35(4): 185-190.
- Colombian corn; 35(1): 36-48.
- Colostrum; 35(3): 165-173.
- Conception rate; 35(2): 68-81.
- Conditioning; 35(4): 205-216.
- Corn kernel quality; 35(1): 36-48.

- Corn; 35(1): 36-48.
- Cows; 35(1): 14-25.
- Cows; 35(1): 3-13.
- Crossbred cows; 35(2): 68-81.
- Crude protein; 35(3): 153-164.
- Desalinated water; 35(3): 174-184.
- Deterrent; 35(4): 205-216.
- Diagnosis; 35(2): 82-92.
- Diagnostic techniques; 35(3): 141-152.
- Diarrhea; 35(4): 233-242.
- Didelphis marsupialis*; 35(4): 205-216.
- Digestibility; 35(1): 14-25.
- Disease detection; 35(3): 141-152.
- Disease surveillance; 35(3): 141-152.
- Dog; 35(2): 82-92.
- Dry matter intake; 35(1): 14-25.
- Dual purpose; 35(2): 68-81.
- Ecotoxicity; 35(4): 185-190.
- Ecotoxicology; 35(4): 185-190.
- Ehrlichia canis*; 35(2): 82-92.
- Elephant grass hay; 35(1): 49-59.
- Embryo culture; 35(2): 118-123.
- Embryo quality; 35(2): 93-108.
- Embryo recovery; 35(2): 93-108.
- Embryo synchronization; 35(2): 118-123.
- Embryo transfer; 35(2): 118-123.
- Embryonic asynchrony; 35(2): 118-123.
- Embryonic development; 35(2): 118-123.
- Endocrine disruptors; 35(2): 61-67.
- Epidemiological surveillance; 35(3): 141-152.
- Equines; 35(2): 109-117.
- Escherichia coli*; 35(1): 26-35.
- Ewes; 35(3): 165-173.
- Fecal score; 35(4): 233-242.
- Feed efficiency; 35(4): 233-242.
- Feedlot; 35(1): 49-59.
- Fertility; 35(2): 68-81.
- Fiber source; 35(1): 49-59.
- Fipronil; 35(4): 185-190.
- First service; 35(2): 68-81.
- Food residues; 35(2): 61-67.
- Forages; 35(4): 217-232.
- Fungal toxins; 35(1): 36-48.
- Gait; 35(3): 153-164.
- Galbanum; 35(4): 191-204.
- Gastritis; 35(2): 109-117.
- Gastrocnemius tendon; 35(3): 153-164.
- Genetic improvement; 35(3): 125-140.
- Genetic parameters; 35(3): 125-140.
- Goat meat; 35(1): 49-59.
- Grain quality; 35(1): 36-48.
- Grazing; 35(4): 217-232.
- Growth performance; 35(4): 191-204.
- Growth rate; 35(4): 233-242.
- Health risks; 35(2): 61-67.
- Helicobacter spp*; 35(2): 109-117.

- Hepatozoon canis*; 35(2): 82-92.
- Heritability; 35(3): 125-140.
- Holstein; 35(1): 14-25.
- Holstein×Zebu; 35(2): 68-81.
- Holstein-Friesian calves; 35(3): 174-184.
- Honeybees; 35(4): 185-190.
- Horse; 35(2): 109-117.
- Human-opossum conflict; 35(4): 205-216.
- Immune response; 35(4): 191-204.
- Import ban; 35(2): 61-67.
- Import policies; 35(2): 61-67.
- Infection; 35(2): 82-92.
- Infection; 35(3): 141-152.
- Influenza A virus; 35(3): 141-152.
- Inheritance; 35(2): 68-81.
- Insecticide; 35(4): 185-190.
- Insecticides; 35(2): 61-67.
- Invertebrates; 35(4): 185-190.
- In vitro* culture; 35(2): 93-108.
- In vitro* embryo production; 35(2): 118-123.
- In vitro* embryo production; 35(2): 93-108.
- In vivo* embryo production; 35(2): 118-123.
- Iron-binding polypeptides; 35(4): 233-242.
- Lactation function shape; 35(1): 3-13.
- Lactation parameters; 35(1): 3-13.
- Lactation; 35(1): 3-13.
- Lactobacillus buchneri*; 35(1): 14-25.
- Lactobacillus plantarum*; 35(1): 14-25.
- Leg abnormality; 35(3): 153-164.
- Linear models; 35(1): 3-13.
- Litter size; 35(3): 125-140.
- Livestock production; 35(2): 61-67.
- Maternal genetic; 35(3): 125-140.
- Medicinal plants; 35(4): 191-204.
- Metabolizable energy; 35(3): 153-164.
- Methane production; 35(4): 217-232.
- Microbial additives; 35(1): 14-25.
- Microbial inoculants; 35(1): 14-25.
- Milk production; 35(1): 3-13.
- Milk yield; 35(1): 14-25.
- Milk yield; 35(1): 3-13.
- Milk; 35(3): 165-173.
- Model comparison; 35(1): 3-13.
- Molecular; 35(3): 141-152.
- Morulae; 35(2): 118-123.
- Mucosa; 35(2): 109-117.
- Multi-resistance; 35(1): 26-35.
- Multi-resistant bacteria; 35(1): 26-35.
- Mycotoxins; 35(1): 36-48.
- Nonlinear models; 35(1): 3-13.
- Non-starch polysaccharides; 35(4): 233-242.
- Nontarget insects; 35(4): 185-190.
- Nursery diet; 35(4): 233-242.
- Nutrient restriction; 35(3): 153-164.
- Nutrients digestibility; 35(3): 174-184.
- Nutritional value; 35(1): 36-48.

- Odors; 35(4): 205-216.
- Opossums; 35(4): 205-216.
- Oral fluids; 35(3): 141-152.
- Organophosphates; 35(2): 61-67.
- Ovine; 35(3): 165-173.
- PCR; 35(2): 109-117.
- Pediococcus pentosaceus; 35(1): 14-25.
- Pesticides; 35(2): 61-67.
- Pig farm; 35(3): 141-152.
- Piglets; 35(4): 233-242.
- Pigs; 35(1): 26-35.
- Polyester mesh culture; 35(2): 93-108.
- Polymerase chain reaction; 35(2): 82-92.
- Post-mortem; 35(2): 109-117.
- Prenatal exposure; 35(2): 61-67.
- Production system; 35(2): 68-81.
- Rabbits; 35(2): 118-123.
- Repellents; 35(4): 205-216.
- Reverse osmosis; 35(3): 174-184.
- Roughage; 35(1): 14-25.
- Roughage; 35(1): 49-59.
- RT-PCR; 35(3): 141-152.
- Rumen fermentation; 35(4): 217-232.
- Ruminants; 35(4): 217-232.
- Salmonella spp.*; 35(1): 26-35.
- Sampling techniques; 35(3): 141-152.
- Sampling; 35(3): 141-152.
- Secretion; 35(3): 165-173.
- Sheep; 35(3): 125-140.
- Sheep; 35(3): 165-173.
- Silvopastoral systems; 35(4): 217-232.
- Single-embryo culture; 35(2): 93-108.
- Small ruminants; 35(1): 49-59.
- Statistical modelling; 35(1): 3-13.
- Stomach; 35(2): 109-117.
- Sugarcane bagasse; 35(1): 49-59.
- Swine; 35(3): 141-152.
- Tartar; 35(2): 109-117.
- Threshold models; 35(3): 125-140.
- Tibia; 35(3): 153-164.
- Tick-borne pathogens; 35(2): 82-92.
- Total dissolved salts; 35(3): 174-184.
- Toxic residues; 35(4): 185-190.
- Toxicity; 35(4): 185-190.
- Tropic; 35(2): 68-81.
- Tropical agriculture; 35(4): 217-232.
- Tropical dairy operations; 35(1): 3-13.
- Ulcer; 35(2): 109-117.
- Univariate analysis; 35(3): 125-140.
- Urban wildlife; 35(4): 205-216.
- US corn; 35(1): 36-48.
- Use of antimicrobials; 35(1): 26-35.
- Vaginal discharge; 35(3): 165-173.
- Valgus/varus angulation; 35(3): 153-164.
- Valgus-varus deformity; 35(3): 153-164.
- Virginiamycin; 35(4): 191-204.

Water intake; 35(3): 174-184.

Water quality; 35(3): 174-184.

Weaning; 35(4): 233-242.

Well-of-the-well culture; 35(2): 93-108.

Wildlife; 35(4): 205-216.

Wildlifemanagement; 35(4): 205-216.

Zeamays; 35(1): 36-48.