










Characterization and antimicrobial resistance of *Moraxella ovis* isolates from clinical cases of contagious ovine keratoconjunctivitis in Mexico

Caracterización y resistencia antimicrobiana de aislamientos de Moraxella ovis de casos clínicos de queratoconjunctivitis contagiosa ovina en México

Caracterização e resistência antimicrobiana de Moraxella ovis isolados de casos clínicos de ceratoconjuntivite contagiosa ovina no México

Giovany Ortiz-Arana¹, Martín Talavera-Rojas¹, Edgardo Soriano-Vargas¹, Erika-Gabriela Palomares-Reséndiz²
Edgar Enríquez-Gómez¹, Celene Salgado-Miranda¹, Jorge Acosta-Dibarrat^{1*}

¹Centro de Investigación y Estudios Avanzados en Salud Animal. Facultad de Medicina Veterinaria y Zootecnia. Universidad Autónoma del Estado de México, México.

²Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias Centro Nacional de Investigación Disciplinaria Salud Animal e Inocuidad Ciudad de México, México.

To cite this article:

Ortiz-Arana G, Talavera-Rojas M, Soriano-Vargas E, Palomares-Reséndiz EG, Enríquez-Gómez E, Salgado-Miranda C, Acosta-Dibarrat J. Characterization and antimicrobial resistance of *Moraxella ovis* isolates from clinical cases of contagious ovine keratoconjunctivitis in Mexico. Rev Colomb Cienc Pecu 2024; 37(1):14–26. <https://doi.org/10.17533/udea.rccp.v37n1a4>

Abstract

Background: Contagious ovine keratoconjunctivitis (OKC) causes blindness in sheep and goats and it is associated with a set of bacterial genera of which some species show antimicrobial resistance. **Objective:** To identify phenotypic-genotypic relationship of antimicrobial resistance from *Moraxella* spp. isolates obtained from clinical cases of contagious ovine keratoconjunctivitis (OKC) in Mexico. **Methods:** A total of 209 samples were obtained from clinical cases of OKC in sheep and 60 *Moraxella ovis* isolates were identified by bacteriological techniques and amplification of *16s rRNA* and *rtxA* genes by PCR. All isolates were evaluated in terms of antimicrobial resistance by the disk diffusion susceptibility test and amplification of resistance genes by PCR. **Results:** We found 14 *Moraxella ovis* isolates with antimicrobial resistance (AMR) and five multiresistant (MDR). The *sul1*, *sul2*, *tetB*, *qnrA*, *qnrB*, *Bla*_{TEM} genes of antimicrobial resistance were amplified, while gene *floR* was not amplified. **Conclusion:** This is the first isolation report of *Moraxella ovis* from ocular lesions in sheep in the State of Mexico, with six antimicrobial resistance genes identified. Results suggest that *Moraxella ovis* plays an important role in the course of the disease and provides a panorama for molecular epidemiological surveillance and bacterial resistance.

Received: March 22, 2023. Accepted: May 1, 2023

*Corresponding author. Centro de Investigación y Estudios Avanzados en Salud Animal, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de México. Phone. +52 (722) 2925548. E-mail: jpacosta00@hotmail.com



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

© 2024 Universidad de Antioquia. Publicado por Universidad de Antioquia, Colombia.

Keywords: antimicrobial resistance; contagious ovine keratoconjunctivitis; epidemiological surveillance; goat; *Moraxella* spp.; multiresistance; resistance genes; sheep.

Resumen

Antecedentes: La queratoconjuntivitis contagiosa ovina (OKC) causa ceguera temporal o permanente en ovinos y caprinos y está asociada a un conjunto de géneros bacterianos, algunos de los cuales muestran resistencia antimicrobiana. **Objetivo:** Identificar la relación fenotípica-genotípica de la resistencia antimicrobiana de aislamientos *Moraxella ovis* obtenidos de casos clínicos de queratoconjuntivitis contagiosa ovina (OKC) en México. **Métodos:** Se obtuvieron un total de 209 muestras de casos clínicos de OKC en ovinos y se identificaron 60 aislamientos de *Moraxella ovis* por técnicas bacteriológicas y amplificación de genes 16s rRNA y *rtxA* por PCR. En todos los aislamientos se evaluó resistencia antimicrobiana mediante prueba de susceptibilidad de difusión en disco y amplificación de genes de resistencia por PCR. **Resultados:** Se encontraron 14 aislamientos de *Moraxella ovis* con resistencia antimicrobiana (AMR) y cinco multirresistentes (MDR). Los genes de resistencia antimicrobiana *sul1*, *sul2*, *tetB*, *qnrA*, *qnrB*, *Bla_{TEM}* fueron amplificados, mientras que el gen *floR* no fue amplificado. **Conclusión:** Este es el primer reporte de aislamiento de *M. ovis* en lesiones oculares en ovinos en el Estado de México, con seis genes de resistencia antimicrobiana identificados. Nuestros resultados sugieren que *Moraxella ovis* juega un papel importante en el curso de la enfermedad y brinda un panorama para su vigilancia epidemiológica molecular y resistencia bacteriana.

Palabras clave: cabra; genes de resistencia; *Moraxella* spp.; multirresistencia; oveja; queratoconjuntivitis contagiosa ovina; resistencia antimicrobiana; vigilancia epidemiológica.

Resumo

Antecedentes: A ceratoconjuntivite contagiosa ovina (OKC) é uma doença infecciosa contagiosa que causa cegueira temporária ou permanente em ovinos e caprinos, esta doença está associada a um conjunto de gêneros bacterianos dos quais alguns deles relataram resistência antimicrobiana. **Objetivo:** O objetivo deste estudo foi identificar a relação fenotípica-genotípica da resistência antimicrobiana de *Moraxella* spp. isolados obtidos de casos clínicos de ceratoconjuntivite contagiosa ovina (OKC) no estado do México. **Métodos:** Um total de 209 amostras foram obtidas de casos clínicos de OKC em ovinos e 60 isolados de *Moraxella ovis* foram identificados por técnicas bacteriológicas e amplificação dos genes *16s rRNA* e *rtxA* por PCR. Todos os isolados foram avaliados quanto à resistência antimicrobiana pelo método de teste de suscetibilidade à difusão em disco e pela amplificação de genes de resistência por PCR respectivamente. **Resultados:** Determinamos 14 isolados de *Moraxella ovis* com resistência antimicrobiana (AMR) e cinco multirresistentes (MDR) e amplificou os genes de resistência antimicrobiana *sul1*, *sul2*, *tetB*, *qnrA*, *qnrB*, *Bla_{TEM}* e não amplificou o gene *floR*. **Conclusão:** É o primeiro relato de isolamento de *Moraxella ovis* em lesões oculares em ovinos no Estado do México e a identificação de seis genes de resistência antimicrobiana. Sugere-se que *Moraxella ovis* desempenha um papel importante no curso da doença e fornece um panorama de interesse em vigilância epidemiológica molecular e resistência bacteriana.

Palavras-chave: cabra; genes de resistência; *Moraxella* spp.; multirresistência; ovelha; ceratoconjuntivite contagiosa ovina; resistência antimicrobiana; vigilância epidemiológica.

Introduction

Contagious ovine keratoconjunctivitis (OKC) is a disease that causes temporary or permanent blindness in sheep and goats. Its clinical signs range from conjunctivitis to corneal ulcers. Several bacterial pathogens have been isolated from OKC, such as *Moraxella*, *Mycoplasma*, and *Chlamydia* (Akerstedt and Hofshagen, 2004; Gupta *et al.*, 2014; Jelocnik *et al.*, 2019). However, three species of the genus *Moraxella* have been associated with infectious keratoconjunctivitis in large and small ruminants, such as *Mor. ovis* (Dagnall, 1994a), *Mor. bovis* (Karthik *et al.*, 2017) and *Mor. bovoculi* (Farias *et al.*, 2015). The differential biochemical tests of *Moraxella* species are phenylalaninedeaminase and gelatinase tests, although they may be inconsistent. Molecular tests are based on the amplification of *16s-23S rRNA* genes *ISR* (Shen *et al.*, 2011; O'Connor *et al.*, 2012; Sosa and Zunino, 2013) and the *rxtA* gene that encodes cytotoxin A (Farias *et al.*, 2015).

Keratoconjunctivitis treatment is based on systemic or local antimicrobials such as tetracyclines, sulfonamides, chloramphenicol, and tulathromycin (Alexander, 2010; Maboni *et al.*, 2015), although several *Moraxella* species have shown resistance to those antimicrobials (Cattray *et al.*, 2007; Loy and Brodersen, 2014; Maboni *et al.*, 2015). Antimicrobial resistance is associated with the presence of genes within the bacterial genome such as *tet* and *otr* that confer resistance to tetracyclines (Martí *et al.*, 2006; Mosquito *et al.*, 2011) *sul* and *dfp* that confer resistance to sulfonamides and trimethoprim (Kern *et al.*, 2002; Ho *et al.*, 2009), *cat*, *cmlA*, and *floR* that confer resistance to phenicols (Schwarz *et al.*, 2004; Chiu *et al.*, 2006), *mph* and *msr* that confer resistance to macrolides and lincosamides (Lüthje and Schwarz, 2006), *Bla_{TEM}*, *Bla_{Rob}* and *Bla_{CARB}* that confer resistance to betalactamases (Bush and Jacoby, 2010; Dallenne *et al.*, 2010) and *qnr*, *gyr*, and *par* that confer resistance to quinolones (Cattoir *et al.*, 2007; Jacoby *et al.*, 2008). Genes such as *tetH*, *sul2*, *floR*, *bla_{Rob-1}*, *srt*, (3')-*Ic*, *mph* and *msr*, have been described in the genome within a pathogenicity island in *Mor. bovoculi* (Dickey *et al.*, 2016) although no other studies have been

reported. Hence, the aim of the present study was to isolate *Moraxella* spp. from ocular lesions in sheep in the State of Mexico (Mexico) and phenotypic-genotypic characterization of the antimicrobial resistance to understand the pathogenesis of these microorganisms in the disease.

Materials and Methods

The experimental protocol was approved by the Review Commission of the Internal Committee for the Care of Laboratory Animals - Teaching, Research, Service and Production of Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma (State of Mexico, Mexico) (CICUAL-DISP FMVZ).

Bacteriological sampling and isolation

Sample size was calculated considering the clinical cases of OKC through the finite population sampling formula (Jaramillo, 2009). A pilot sampling was used for estimating the *p* indicator, obtaining a prevalence of 0.23 (107/454), and estimating 209 samples. Samples (*n*=209) were collected from a total of 845 clinically healthy sheep with ocular lesions (e.g. ulcers, blindness) suggestive of OKC by general physical examination and ophthalmological tests in 15 farms from six municipalities of the State of Mexico between February and June 2017. Specimens were obtained by conjunctival swabs in the lesion area without touching the palpebral edge. The samples were placed in Stuart transport medium (STM, Cat. 1058-A, Dibico. Cuautitlán Izcalli, Mexico), maintained in refrigeration at 4 °C, and processed before 24 h (Akerstedt and Hofshagen, 2004).

Regarding sample isolation, inoculations were performed on 5% ovine blood agar plates (ABS, BLL, Cat BD211037. Becton-Dickinson. CDMX, Mexico) and incubated in aerobic conditions at 37 °C for 24-48 h. Colony growth criteria and classical biochemical tests described in the literature were used for identification of *Moraxella* species involved in keratoconjunctivitis (Angelos *et al.*, 2007; Angelos and Ball, 2007; Shen *et al.*, 2011).

Genotypic identification through the *16s rRNA* and *rtxA* genes of *Moraxella* spp.

For DNA extraction, a heated colony was used in a total volume of 100 µl of sterile distilled water heated to 95 °C for 10 min followed by centrifugation of the cell suspension for 5 min at 9279 G (Eppendorf® Microcentrifuge 5415, Merck KGaA, Darmstadt, Germany) and subsequent DNA collection (Dallenne *et al.*, 2010).

Genes *16s rRNA* and *rtxA* were amplified by a multiplex PCR with a final reaction volume of 25 µl for each one containing: 12.5 µl Master Mix (Gotaq Green Master mix, Cat M7122. Promega Corporation, WI, USA), 1 µl of each primer (*Ovi16S1 F/Ovis1849 R*, *Bviv16S1 F/Bovi1541 R* and *Bovo1915 R*) for the *16s rRNA* gene and for the *rtxA* gene (*MbxA F/MbxA R*, *MbvA F* and *MovA R*), 4 µl of bacterial DNA and 7.5 µl nucleases free water (Nuclease-Free Water Cat. P1195. Promega Corporation, WI, USA). The DNA sequences and PCR product sizes are described in Table 1.

The PCR protocol regarding *16s rRNA* gene was as follows: initial denaturation for 5 min at 95 °C followed by 35 cycles: denaturation for 40 s at 95 °C, alignment for 40 s at 55 °C, extension for 1 min at 72 °C; and a final extension for 7 min at 72 °C. For gene *rtxA* 35: cycles denaturation for 50 s at 95 °C, alignment for 50 s at 65 °C, extension for 1 min at 72 °C, and a final extension for 4 min at 72 °C

carried out in a ThermoCycler (MultiGene™ Mini, TC 020-24, Labnet International Inc. CA, USA). All the amplification products were identified through horizontal electrophoresis in 1% agarose gels stained with 0.5 µg/mL of ethidium bromide and visualized with a UV transilluminator (Mini-Bis 16mm, DNr Bio-Imaging Systems. Neve Yamin Israel; Shen *et al.*, 2011; Farias *et al.*, 2015). A *Staphylococcus aureus* ATCC 25923 strain was used as negative control.

Antimicrobial susceptibility tests

Susceptibility tests were carried out using the disk diffusion method on Mueller Hinton agar (AMH, BD Bioxon. Becton-Dickinson. CDMX, Mexico) supplemented to 5% with ovine blood defibrinated according to the guidelines of the Institute of Clinical and Laboratory standards (CLSI, 2016). The bacterial solution turbidity suspended Mueller-Hinton was adjusted (MH, BBL TM. Becton-Dickinson. CDMX, Mexico) at a scale of 0.5 Mc Farland equivalent to a concentration of 1-2 x10⁸ CFU/mL, the following antimicrobials were used; ampicillin (10 µg), carbencillin (100 µg), cephalothin (30 µg), cefotaxime (30 µg), ciprofloxacin (5 µg), chloramphenicol (30 µg), nitrofurantoin (30 µg) netilmicin (30 µg), gentamicin (10 µg), amikacin (30 µg), sulfamethoxazole/trimethoprim (25 µg), norfloxacin (10 µg), tetracycline (30 µg) and nalidixic acid (30 µg) (BBL™ Sensi-Disc™).

Table 1. Primer designing for *Moraxella* spp. identification using PCR.

Gene	Species	Primers	Sequences 5'- 3'	Frag-ment size	Reference
rtxA	<i>Mor. bovis</i>	MbxAF	GCA AAA CTG GCA ATG ACG A	943 bp	Farias <i>et al.</i> (2015)
		MbxAR	GTG CCA TTG ACC CAA CTA GC		
	<i>Mor. bovoculi</i>	MbvAF	AAT GCT GGT GCT GGT AAC GA	990 bp	
		<i>Mor. ovis</i>	MovAR		
16s rRNA	<i>Mor. ovis</i>	Ovi16S1F	GAA CGA TGA GTA TCC AGC TTG CT	1849 bp	Shen <i>et al.</i> (2011)
		Ovis1849R	CTC TTT ACT TTG GTT AAT TAT TTT GTT GGA		
	<i>Mor. bovoculi</i>	Bovo1915R	TGT ATT GGG TAC AAT CAC CAT GG	1859 bp	
		Bviv16S1F	GAA CGA TGA CTA TCT AGC TTG CTA GAT ATG		
		<i>Mor. bovis</i>	Bovi1541R		

Becton-Dickinson. CDMX, Mexico). The plates were incubated at 37 °C for 18-24 h. Regarding interpretation of the results, the following profiles were established: sensitive (S), intermediate (I), and resistant (R). An *Escherichia coli* ATCC 25922 strain was used as control. As no standardized criteria for the interpretation of sensitivity exist for *Moraxella* spp., breakpoints established for Gram-negative pathogens related with cattle respiratory disease were used (Maboni *et al.*, 2015) (e.g. critical breakpoints against respiratory pathogens of *Pasteurella multocida*, *Moraxella catarrhalis*, *Mannheimia haemolytica*, *Pseudomonas aeruginosa* and *Haemophilus sommus*) (CLSI, 2013; 2016).

Antimicrobial resistance genes

Polymerase chain reaction (PCR) was used to identify antimicrobial resistance genes. Seven primers were used. Sequences and amplification products sizes are described in Table 2. Detection of genes *sul1* and *sul2* (sulfonamides), *Bla_{TEM}* (β -Lactams), *tetB* (tetracyclines), *floR* (florfenicol/chloramphenicol), as well as a multiplex PCR for *qnrA* and *qnrB* genes (quinolones) followed the same reaction conditions previously published (Kern *et al.*, 2002; Chiu *et al.*, 2006; Martí *et al.*, 2006;

Cattoir *et al.*, 2007; Dalenne *et al.*, 2010). Positive controls used were as follows: *Escherichia coli* ATCC 25922 and other *E. coli* isolates from sewage characterized as phenotypically resistant to AM (ampicillin), CB (carbencillin), CF (cephalothin), CL (chloramphenicol), NA (nalidixic acid), TE (tetracycline), NET (netilmicin), NF nitrofurantoin, CPF (ciprofloxacin) and genotypically possessing *qnrA*⁺, *qnrB*⁺ and *Bla_{TEM}*⁺ genes (Talavera-González *et al.*, 2021).

Results

A total of 209/861 examined sheep (24.27% prevalence) in six municipalities of the State of Mexico showed lesions compatible with OKC. With respect to the total number of the studied samples (209), we identified 60 isolates of *Moraxella* spp. by biochemical tests. Through the 16s rRNA gene, 54 strains of *Mor. ovis* were correctly identified from the 60 isolates (90%). The 1541 and 1959 pb amplicons corresponding to *Mor. bovis* and *Mor. bovoculi* not amplified by any isolate, respectively. For the *rtxA* gene, 57 strains of *Mor. ovis* were identified of the 60 isolates (95%) (Table 3) and no isolates amplified a band of 943 pb corresponding to *Mor. bovis*.

Table 2. Antimicrobial resistance used for the identification of *Moraxella* spp.

Resistance	Primers	Sequences 5'-3'	Fragment size	Reference
Sulfonamides	<i>sul1 F</i>	CGG CGT GGG CTA CCT GAA CG	433 bp	Kern <i>et al.</i> (2002)
	<i>sul1 R</i>	GCC GAT CGC GTG AAG TTC CG		
	<i>sul2 F</i>	GCG CTC AAG GCA GAT GGC ATT	293 bp	
	<i>sul2 R</i>	GCG TTT GAT ACC GGC ACC CGT		
Tetracyclines	<i>tet B F</i>	TTG GTT AGG GGC AAG TTT TG	650 bp	Martí <i>et al.</i> (2006)
	<i>tet B R</i>	GTA ATG GGC CAA TAA CAC CG		
Quinolones	<i>qnrA F</i>	AGA GGA TTT CTC ACG CCA GG	580 bp	Cattoir <i>et al.</i> (2007)
	<i>qnrA R</i>	TGC CAG GCA CAG ATC TTG AC		
	<i>qnrB F</i>	GGM ATH GAA ATT CGC CAC TG	264 pb	
	<i>qnrB R</i>	TTT GCY GYY CGC CAG TCG AA		
β -lactamases	<i>MultiTSO-T Bla_{TEM} F</i>	CAT TTC CGT GTC GCC CTT ATT C	800 bp	Dalenne <i>et al.</i> (2010)
	<i>MultiTSO-T Bla_{TEM} R</i>	CGT TCA TCC ATA GTT GCC TGA C		
Florfenicol/ Chloramphenicol	<i>floR F</i>	CTT TGG CTA TAC TGG CGA TG	266 bp	Chiu <i>et al.</i> (2006)
	<i>floR R</i>	GAT CAT TAC AAG CGC GAC AG		

Y=T o C; R=A or G, S=G or C, D=A or G or T, H=A or C or T; M=A o C.

Antimicrobial susceptibility tests

The 60 isolates of *Mor. ovis* were sensitive to gentamicin and norfloxacin 100% (60/60). Antimicrobial resistance of 18.3% (11/60) to nalidixic acid was observed, 11.6% (7/60) to

nitrofurantoin, 10.0% (6/60) to ampicillin, 6.6% (4/60) to chloramphenicol, 3.3% (2/60) to cephalothin, 3.3% (2/60) to tetracycline, and 1.6% (1/60) to cefotaxime, netilmicin, amikacin and sulfamethoxazole/trimetoprim (Table 4).

Table 3. *Moraxella* spp. isolates obtained from clinical ovine keratoconjunctivitis cases in sheep in the State of Mexico.

UPP	Municipality	Animals	Ocular injuries*				Samples	Bacteriology	Molecular		
			C	KC	K	U		Isolates	16S rRNA 1949 pb	RtxA 990 pb	
								<i>Mor. ovis</i>	<i>Mor. ovis</i>	<i>Mor. ovis</i>	
			51	14	2	-	67	29	26	26	
4	Toluca	189	48	7	1	-	56	5	3	5	
3	Xonacatlán	117	23	-	-	-	23	4	4	4	
3	Lerma	80	20	10	1	-	31	10	10	10	
1	Capulhuac	43	15	-	-	-	15	4	4	4	
1	Calimaya	150	8	8	-	1	17	8	7	8	
15	Total	861	16	39	4	1	209*	60	54	57	
							Prevalence	24.27%	28.70%	90%	95%

*Ocular lesions correspond to the number of animals sampled. UPP: Number of Production Units; C: Conjunctivitis; KC: Keratoconjunctivitis; K: Keratitis; U: Corneal ulcers.

Table 4. Susceptibility test of *Moraxella* spp. causing ovine keratoconjunctivitis.

Antimicrobial	<i>Mor. ovis</i> (n= 60) (#) prevalence			Zone diameter
	Resistance	Intermediate	Susceptible	Interpretive criteria (nearest whole mm)
NA	11(18.3)	8(13.3)	41(68.3)	≤22-≥28
NF	7(11.6)	1(1.6)	52(86.6)	≤14-≥17
AM	6(10.0)	7(11.6)	47(78.3)	≤13-≥27
CL	4(6.6)	2(3.3)	54(90.0)	≤25-≥29
CF	2(3.3)	3(5.0)	55(91.6)	≤16-≥20
TE	2(3.3)	1(1.6)	57(95.0)	≤24-≥29
CFX	1(1.6)	1(1.6)	58(96.6)	≤10-≥13
SXT	1(1.6)	6(10)	53(88.3)	≥26
NET	1(1.6)	-	59(98.3)	≤12-15
AK	1(1.6)	-	59(98.3)	≤14-17
CP	.	2(3.3)	58(96.6)	≥21
CB	-	2(3.3)	58(96.6)	≤18-≥25
GE	-	-	60(100)	≤12-≥15
NOF	-	-	60(100)	≤12-≥17

#: Isolates; (#): Percentage. AM: Ampicillin; CB: Carbenicillin; CF: Cephalothin; CFX: Cefotaxime; CP: Ciprofloxacin; CL: Chloramphenicol; NF: Nitrofurantoin; NET: Netilmicin; GE: Gentamicin; AK: Amikacin; STX: Sulfamethoxazole/trimethoprim; NOF: Norfloxacin; TE: Tetracycline; NA: Nalidixic acid (CLSI 2013; 2016).

A total of 68.33% (41/60) isolates showed sensitivity to all the antimicrobials used, and 31.66% (19/60) were resistant to one or more antimicrobials. Furthermore, 23.33% (14/60) isolates were resistant to one or two antimicrobials and 8.33% (5/60) were multi-resistant strains (Table 4).

Phenotypic and genotypic profiles

Mor. ovis presented 36.66% (22/60) and 33.33% (20/60) amplification of *sul1* and *sul2* gene, respectively. Isolates of *Mor. ovis* 46.66% (28/60) amplified gene *Bla_{TEM}*. The gene that confers resistance to tetracyclines, 8.33% (5/60) of *Mor. ovis* isolates amplified a 650 pb band corresponding to *tetB*. *Mor. ovis* isolates (23/60) 38.33% amplified *qnrA* gene and (22/60) 36.66% *qnrB* gene (Table 5).

Discussion

An OKC prevalence of 24.27% (209/861) was obtained. This was lower than reported in the United Kingdom (Dagnall, 1994a; 1994b) 72.38% (97/134). Severity of eye lesions was observed, which could be related to the period taken from the samples to the raising systems and predisposing factors. An increase in OKC cases was reported in Norway during the autumn and winter seasons due to animal management: in summer sheep graze without human intervention, while they are confined in barns during the winter (Akerstedt and Hofshagen, 2004). In Mexico,

sheep breeding systems vary from intensive, semi-intensive to extensive, mixed, and grazing. In the present study, sampling was carried out in the spring-winter period, which is characterized by the presence of flies, dust, direct sunlight as well as other factors involved in the evolution of the disease (Egwu *et al.*, 1989). Dagnall (1994b) reported a prevalence of 28.86% (28/97) for *Mor. ovis* from ovine isolates. Similarly, Akerstedt and Hofshagen, (2004) obtained a prevalence of 28.23% (24/85) in sheep herds with the disease. Similar data were obtained in this work: 27.75% (58/209) of *Mor. ovis* isolates from sheep with OKC.

The first isolation of *Mor. bovoculi* was reported in the United States (Angelos *et al.*, 2007) and later in countries such as Uruguay (Sosa and Zunino, 2013), Argentina, Norway, and Brazil (Libardoni *et al.*, 2012; Farias *et al.*, 2015). The first report of *Mor. bovoculi* in sheep was made by Farias *et al.* (2015) in Brazil. Karthik *et al.* (2017) were the first researchers who identified *Mor. bovis* from ocular injuries in sheep in India. In our study it was not possible to isolate *Mor. bovis* and *Mor. bovoculi* from ocular injuries in sheep.

Shen *et al.* (2011) amplified the 16s *rRNA* gene in 89.5% (51/57) of the isolates, identifying *Mor. bovoculi* (44/51) and *Mor. bovis* (7/51). In this study 90.0% (54/60) of the isolates were amplified and correctly identified as *Mor. ovis*.

Table 5. Phenotypic and genotypic profile of antimicrobial resistance in *Moraxella*.

Phenotypical resistance	Resistance genes	<i>Mor. Ovis</i> (n= 60)			Total	Prevalence of re-sistance genes (%)
		Resistance	Intermediate	Susceptible		
AM	<i>Bla_{TEM}</i>	5(6)	6(7)	17(48)	28(60)	46.66
NA	<i>qnrA</i>	8(11)	1(8)	14(41)	23(60)	38.33
	<i>qnrB</i>	8(11)	2(8)	12(41)	22(60)	36.66
STX	<i>sul1</i>	1(1)	1(1)	20(58)	22(60)	36.66
	<i>sul2</i>	1(1)	1(1)	18(58)	20(60)	33.33
TE	<i>tetB</i>	1(2)	0(1)	4(57)	5(60)	8.33
CL	<i>floR</i>	0(4)	0(2)	0(54)	0(60)	0.00

#: Gene amplification; #: All isolates. AM: Ampicillin; TE: Tetracycline; CL: Chloramphenicol; STX: Sulfamethoxazole/trimethoprim; NA: Nalidixic acid.

Farias *et al.* (2015) amplified the gene *rtxA* in 100% (33/33) of isolates from bovine and ovine with keratoconjunctivitis, identifying *Mor. bovis* (15/33), *Mor. bovoculi* (11/33) and *Mor. ovis* (7/33). In the present study, we amplified the *rtxA* gene 95% (57/60) of all isolates identifying *Mor. ovis*. Regarding the six *Mor. ovis* isolates that did not amplify the 16s rRNA and the three *Mor. ovis* isolates that did not show amplification for *rtxA* gene, they possibly could be related to the variations in the reading frame of the sequences by deletions or absence of repeated sequence regions surrounding the RTX operon which has been reported in non-hemolytic *Mor. bovis* strains (Angelos *et al.*, 2003). In recent studies, Dickey *et al.* (2018) detected a recombination in the nucleotide sequences within the non-coding regions in the *rRNA* and *RTX* genes in strains of *Mor. bovoculi*.

The first study of antimicrobial sensitivity *in vitro* *Mor. ovis* strains performed by Elad *et al.* (1988) reported resistance to penicillin, ampicillin, streptomycin and neomycin. Another study, conducted by Catry *et al.* (2007), reported strains resistant to erythromycin. The most recent work by Maboni *et al.* (2015) showed strains resistant to oxytetracycline and penicillin. All the mentioned researchers described *Mor. ovis* strains sensitive to gentamicin, chloramphenicol, florfenicol, and sulfonamides. In the present study we reported strains of *Mor. ovis* susceptible to gentamicin, as well as resistant strains to ampicillin, chloramphenicol, tetracycline, and sulfamethoxazole/trimethoprim.

Oxytetracycline is usually the first choice of treatment for keratoconjunctivitis (Alexander, 2010); however, *Moraxella* spp. showed resistance over time to this antimicrobial (Maboni *et al.*, 2015). Likewise, florfenicol was reported as an effective therapeutic option in keratoconjunctivitis (Gokce *et al.*, 2002; Angelos *et al.*, 2011). The use of antimicrobials is essential for controlling OKC by *Mor. ovis* to avoid exacerbation of lesions associated with other bacterial infections (Dagnall, 1994b).

To the best of our knowledge, this is the first study to amplify the antimicrobial resistance genes *sul1*, *sul2*, *tetB*, *qnrA* *qnrB* y *Bla_{TEM}* in *Mor. ovis*. A study associated with antimicrobial resistance determinants (ARD) in *Mor. bovoculi* was performed by Dickey *et al.* (2016); they described 10 ARD located on a genomic island greater than 27 kb in the sequences of *Mor. bovoculi* and Mb58069 isolates that were resistant to florfenicol, oxytetracycline, sulfonamides, and showed intermediate resistance to macrolides.

A similar study, conducted by Roberts *et al.* (1991), described tetracycline-resistant strains of *Mor. catarrhalis* that carry the *tetB* gene on its chromosome. The *tetB* gene has the widest range of Gram-negative bacteria, such as *E. coli* (Medina *et al.*, 2011; Mirzaagha *et al.*, 2011), *Acinetobacter baumannii* (Martí *et al.*, 2006), *Actinobacillus actinomycetemcomitans* (Roe *et al.*, 1995), *Haemophilus influenzae* (Robert and Smith, 1980), and *Treponema denticolaum* (Roberts, 1996).

Bacteria carry resistance genes *sul1*, *sul2*, *sul3*, and *Bla_{TEM}* such as *E. coli* (Kern *et al.*, 2002; Infante *et al.*, 2005; Ho *et al.*, 2009; Medina *et al.*, 2011; Gnida *et al.*, 2014; Memariani *et al.*, 2015), *Klebsiella pneumoniae*, *Pseudomona aeruginosa* (Peymani *et al.*, 2017), *Proteus mirabilis* (Feizabadi *et al.*, 2010; Gong *et al.*, 2018), *Salmonella* spp. (Maka *et al.*, 2015), *Stenotrophomonas maltophilia* (Hu *et al.*, 2011). Bacteria carriers of genes *qnrA* and *qnrB* are *K. pneumoniae* (Rodríguez-Martínez *et al.*, 2003), *E. coli* (Wang *et al.*, 2003; Jiang *et al.*, 2008; Aguilar-Montes de Oca *et al.*, 2015), *P. aeruginosa*, *Enterobacter cloacae* (Wu *et al.*, 2007), *Actinobacter baumannii* (Touati *et al.*, 2008), *Salmonella enterica* (Murray *et al.*, 2008), *Enterobacter aerogenes*, *Citrobacter freundii* (Park *et al.*, 2007), *Kluyvera* (Kraychete *et al.*, 2016), among others.

Mor. ovis was identified in the present study by using *16s rRNA* and *RtxA* genes with PCR, confirming PCR as the most sensitive test for diagnosing bacterial agents involved in keratoconjunctivitis. It will be possible to establish new criteria for choosing antimicrobials based

on the phenotypic and genotypic characteristics of antimicrobial resistance, which will also allow for molecular epidemiology surveillance of antimicrobial resistance genes in bacterial populations of *Moraxella* spp.

Declarations

Acknowledgment

To the sheep producers of the State of Mexico for allowing us to take the samples at their farms.

Funding

This work was funded by Universidad Autónoma del Estado de Mexico through the projects "Caracterización, fenotípica, genotípica y resistencia antimicrobiana de *Moraxella* spp., obtenidas de casos clínicos de queratoconjunctivitis ovina" (key: 4629/2019SF) and "Caracterización genotípica de *Mycoplasma* spp., *Chlamydia* spp. y *Moraxella* spp. presentes en casos clínicos de Oueratoconjunctivitis Ovina y su relación con el cuadro clínico de la enfermedad" (key: 6177/2020CIB). The first author received a scholarship from CONACYT Mexico to carry out his doctoral studies.

Conflicts of interest

The authors declare they have no conflicts of interest regarding the work presented in this report.

Author contributions

Acosta-Dibarrat J, Talavera Rojas M, Soriano Vargas E, and Ortiz Arana G designed the experiment. Acosta-Dibarrat J and Ortiz Arana G administered the project. Ortiz Arana G, Palomares-Resendiz EG, Salgado-Miranda C, and Enriquez-Gomez E worked on the aspects involved in the methodology. Ortiz Arana G and Acosta Dibarrat J wrote and prepared the manuscript. All authors provided critical feedback during writing and editing.

Use of artificial intelligence (AI)

No AI or AI-assisted technologies were used during the preparation of this work.

References

- Aguilar-Montes de Oca S, Talavera-Rojas M, Soriano-Vargas E, Barba-León J, Vazquez-Navarrete J. Determination of extended spectrum β -lactamases/AmpC β -lactamases and plasmid-mediated quinolone resistance in *Escherichia coli* isolates obtained from bovine carcasses in Mexico. *Trop Anim Health Prod* 2015; 47:975–981. <https://doi.org/10.1007/s11250-015-0818-3>
- Akerstedt J, Hofshagen M. Bacteriological investigation of infectious keratoconjunctivitis in Norwegian sheep. *Acta Vet Scand* 2004; 45:19–26. <https://doi.org/10.1186/1751-0147-45-19>
- Alexander D. Infectious bovine keratoconjunctivitis: A review of cases in clinical practice. *Vet Clin North Am - Food Anim Pract* 2010; 26(3):487–503. <https://doi.org/10.1016/j.cvfa.2010.09.006>
- Angelos JA, Ball LM. Differentiation of *Moraxella bovoculi* sp. nov. from other coccoid *Moraxellae* by the use of polymerase chain reaction and restriction endonuclease analysis of amplified DNA. *J Vet Diagn Invest* 2007; 19(5):532–534. <https://doi.org/10.1177/104063870701900511>
- Angelos JA, Ball LM, Byrne BA. Minimum inhibitory concentrations of selected antimicrobial agents for *Moraxella bovoculi* associated with infectious bovine keratoconjunctivitis. *J Vet Diagn Invest* 2011; 23(3):552–555. <https://doi.org/10.1177/1040638711040154>
- Angelos JA, Hess JF, George LW. An RTX operon in hemolytic *Moraxella bovis* is absent from nonhemolytic strains. *Vet Microbiol* 2003; 92(4):363–377. [https://doi.org/10.1016/S0378-1135\(02\)00410-8](https://doi.org/10.1016/S0378-1135(02)00410-8)
- Angelos JA, Spinks PO, Ball LM, George LW. *Moraxella bovoculi* sp. nov., isolated from calves with infectious bovine keratoconjunctivitis. *Int J Syst Evol Microbiol* 2007; 57(4): 789–795. <https://doi.org/10.1099/ijs.0.64333-0>

- Bush K, Jacoby GA. Updated functional classification of β -Lactamases. *Antimicrob Agents Chemother* 2010; 54(3):969–976. <https://doi.org/10.1128/AAC.01009-09>
- Catry B, Boyen F, Baele M, Dewulf J, de Kruijff A, Vaneechoutte M, Haesebrouck F, Decostere A. Recovery of *Moraxella ovis* from the bovine respiratory tract and differentiation of *Moraxella* species by tDNA-intergenic spacer PCR. *Vet Microbiol* 2007; 120(3–4):375–380. <https://doi.org/10.1016/j.vetmic.2006.10.037>
- Cattoir V, Poirel L, Rotimi V, Soussy CJ, Nordmann P. Multiplex PCR for detection of plasmid-mediated quinolone resistance qnr genes in ESBL-producing enterobacterial isolates. *J Antimicrob Chemother* 2007; 60(2):394–397. <https://doi.org/10.1093/jac/dkm204>
- Chiu CH, Su LH, Chu CH, Wang MH, Yeh CM, Weill FX, Chu C. Detection of multidrug-resistant *Salmonella enterica* serovar *typhimurium* phage types DT102, DT104, and U302 by multiplex PCR. *J Clin Microbiol* 2006; 44(7):2354–2358. <https://doi.org/10.1128/JCM.00171-06>
- Clinical and Laboratory Standards Institute (CLSI). Antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals. Approved Standard - 4 edition. CLSI document VET01-A4. 2013. Wayne, PA.1-72.
- Clinical and Laboratory Standard Institute (CLSI). Methods for antimicrobial dilution and disk susceptibility testing of infrequently isolated or fastidious bacteria; Approved Guideline- 3 Edition. CLSI document M45. 2016. Wayne, PA. 1-90.
- Dagnall GJR. An investigation of colonization of the conjunctival sac of sheep by bacteria and mycoplasmas. *Epidemiol Infect* 1994a; 112(3):561–567. <https://doi.org/10.1017/S0950268800051268>
- Dagnall GJR. The role of *Branhamella ovis*, *Mycoplasma conjunctivae* and *Chlamydia psittaci* in conjunctivitis of sheep. *Br Vet J* 1994b; 150(1):65–71. [https://doi.org/10.1016/S0007-1935\(05\)80097-1](https://doi.org/10.1016/S0007-1935(05)80097-1)
- Dallenne C, da Costa A, Decré D, Favier C, Arlet G. Development of a set of multiplex PCR assays for the detection of genes encoding important β -lactamases in Enterobacteriaceae. *J Antimicrob Chemother* 2010; 65(3):490–495. <https://doi.org/10.1093/jac/dkp498>
- Dickey AM, Loy JD, Bono JL, Smith TPL, Apley MD, Lubbers BV, Dedonder KD, Capik SF, Larsen RL, White BJ, Blom J, Chitko-McKown CG, Clawson ML. Large genomic differences between *Moraxella bovoculi* isolates acquired from the eyes of cattle with infectious bovine keratoconjunctivitis versus the deep nasopharynx of asymptomatic cattle. *Vet Res* 2016; 47:31(2016). <https://doi.org/10.1186/s13567-016-0316-2>
- Dickey AM, Schuller G, Loy JD, Clawson ML. Whole genome sequencing of *Moraxella bovoculi* reveals high genetic diversity and evidence for interspecies recombination at multiple loci. *PLoS One* 2018; 13(12):e0209113. <https://doi.org/10.1371/journal.pone.0209113>
- Elad D, Yeruham I, Bernstein M. *Moraxella ovis* in cases of infectious bovine keratoconjunctivitis (IBK) in Israel *J Vet Med Ser B* 1988; 35(1–10): 431–434. <https://doi.org/10.1111/j.1439-0450.1988.tb00516.x>
- Egwu GO, Faull WB, Bradbury JM, Clarkson MJ. Ovine infectious keratoconjunctivitis: a microbiological study of clinically unaffected and affected sheep's eyes with special reference to *Mycoplasma conjunctivae*. *Vet Rec* 1989; 125(10):253–256. <https://doi.org/10.1136/vr.125.10.253>
- Farias LDA, Maboni G, Matter LB, Scherer CFC, Libardoni F, de Vargas AC. Phylogenetic analysis and genetic diversity of 3' region of *rtxA* gene from geographically diverse strains of *Moraxella bovis*, *Moraxella bovoculi* and *Moraxella ovis*. *Vet Microbiol* 2015; 178(3–4):283–287. <https://doi.org/10.1016/j.vetmic.2015.05.025>
- Feizabadi MM, Delfani S, Raji N, Manjooi A, Aligholi M, Shahcheraghi F, Parvin M, Yadegarinia D. Distribution of blaTEM, blaSHV, blaCTX-M genes among clinical isolates of *Klebsiella pneumoniae* at Labbafinejad Hospital, Tehran, Iran. *Microb Drug Resist* 2010; 16(1):49–53. <https://doi.org/10.1089/mdr.2009.0096>

- Gnida A, Kunda K, Ziemińska A, Łuczkiwicz A, Felis E, Surmacz-Górska J. Detection of sulfonamide resistance genes via *in situ* PCR-FISH. Polish J Microbiol 2014; 63(2):167–173. <http://www.pjmonline.org/wp-content/uploads/archive/vol6322014167.pdf>
- Gokce HI, Cital M, Genc O, Erdogan HM, Gunes V, Kankavi O. A comparison of the efficacy of florfenicol and oxitetracycline in treatment of naturally occurring infectious bovine keratoconjunctivitis. Ir Vet J 2002; 55(11):573–576.
- Gong J, Zhuang L, Zhang D, Zhang P, Dou X, Wang C. Establishment of a multiplex loop-mediated isothermal amplification method for rapid detection of sulfonamide (sul1, sul2, sul3) in clinical Enterobacteriaceae isolates from poultry. Foodborne Pathog Dis 2018; 15(7):413–419. <https://doi.org/10.1089/fpd.2017.2410>
- Gupta S, Chahota R, Bhardwaj B, Malik P, Verma S, Sharma M. Identification of *Chlamydiae* and *Mycoplasma* species in ruminants with ocular infections. Lett Appl Microbiol 2014; 60(2): 135–139. <https://doi.org/10.1111/lam.12362>
- Ho PL, Wong RC, Chow KH, Que TL. Distribution of integron-associated trimethoprim-sulfamethoxazole resistance determinants among *Escherichia coli* from humans and food-producing animals. Lett Appl Microbiol 2009; 49(5):627–634. <https://doi.org/10.1111/j.1472-765X.2009.02717.x>
- Hu L, Chang X, Ye, Y, Wang Z, Shao Y, Shi W, Li X, Li J. Stenotrophomonas maltophilia resistance to trimethoprim/sulfamethoxazole mediated by acquisition of sul and dfrA genes in a plasmid-mediated class 1 integron. Int J Antimicrob Agents 2011; 37(3):230–234. <https://doi.org/10.1016/j.ijantimicag.2010.10.025>
- Infante B, Grape M, Larsson M, Kristiansson C, Pallecchi L, Rossolini GM, Kronvall G. Acquired sulphonamide resistance genes in faecal *Escherichia coli* from healthy children in Bolivia and Peru. Int J Antimicrob Agents 2005; 25(4):308–312. <https://doi.org/10.1016/j.ijantimicag.2004.12.004>
- Jacoby G, Cattoir V, Hooper D, Martínez-Martínez L, Nordmann P, Pascual A, Poirel L, Wang M. Qnr gene nomenclature. Antimicrob Agents Chemother 2008; 52(7):2297–2299. <https://doi.org/10.1128/AAC.00147-08>
- Jaramillo CJ. Epidemiologia Veterinaria. 1 ed (Mexico) Manual Moderno 2009.
- Jelocnik M, Laurence M, Murdoch FR, Polkinghorne A. Detection of Chlamydiaceae in ocular swabs from Australian pre-export feedlot sheep. Aust Vet J 2019(10); 97: 1–3. <https://doi.org/10.1111/avj.12857>
- Jiang Y, Zhou Z, Qian Y, Wei Z, Yu Y, Hu S, Li L. Plasmid-mediated quinolone resistance determinants qnr and aac(60)-Ib-cr in extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in China. J Antimicrob Chemother 2008; 61(5): 1003–1006. <https://doi.org/10.1093/jac/dkn063>
- Karthik K, Manimaran K, Mahaprabhu R, Shoba K. Isolation of *Moraxella* sp. from cases of keratoconjunctivitis in an organized sheep farm of India. Open J Vet Med 2017; 7(10):138–143. <https://doi.org/10.4236/ojvm.2017.710014>
- Kern MB, Klemmensen T, Frimodt-Møller N, Espersen F. Susceptibility of Danish *Escherichia coli* strains isolated from urinary tract infections and bacteraemia, and distribution of sul genes conferring sulphonamide resistance. J Antimicrob Chemother 2002; 50(4): 513–516. <https://doi.org/10.1093/jac/dkf164>
- Kraychete GB, Botelho LAB, Campana EH, Picão RC, Bonelli RR. Updated multiplex PCR for detection of all six plasmid-mediated qnr gene families. Antimicrob Agents Chemother 2016; 60(12): 7524–7526. <https://doi.org/10.1128/AAC.01447-16>
- Libardoni F, Scherer CFC, Farias L, Vielmo A, Balzan C, Vargas AC. *Moraxella bovoculi* em casos de ceratoconjuntivite infecciosa bovina no Rio Grande do Sul. Pesqui Vet Bras 2012; 32(8):743–746. <https://doi.org/10.1590/S0100-736X2012000800011>

- Loy JD, Brodersen BW. *Moraxella* spp. isolated from field outbreaks of infectious bovine keratoconjunctivitis: a retrospective study of case submissions from 2010 to 2013. *J Vet Diagnostic Investig* 2014; 26(6):761–768. <https://doi.org/10.1177/1040638714551403>
- Lüthje P, Schwarz S. Antimicrobial resistance of coagulase-negative staphylococci from bovine subclinical mastitis with particular reference to macrolide–lincosamide resistance phenotypes and genotypes. *J Antimicrob Chemother* 2006; 57(5):966–969. <https://doi.org/10.1093/jac/dk1061>
- Maboni G, Gressler LT, Espindola JP, Schwab M, Tasca C, Potter L, de Vargas AC. Differences in the antimicrobial susceptibility profiles of *Moraxella bovis*, *M. bovoculi* and *M. ovis*. *Brazilian J Microbiol* 2015; 46:545–549. <https://doi.org/10.1590/S1517-838246220140058>
- Maka L, Mackiw E, Scienzynska H, Modezelewska M, Popowska M. Resistance to sulfonamides and dissemination of sul genes among *Salmonella* spp. isolated from food in Poland. *Foodborne Pathog Dis* 2015; 12(5):383–389. <https://doi.org/10.1089/fpd.2014.1825>
- Martí S, Fernández-Cuenca, F, Pascual Á, Ribera A, Rodríguez-Baño J, Bou G, Cisneros, JM, Pachón J, Martínez-Martínez L, Vila J. Prevalencia de los genes tetA y tetB como mecanismo de resistencia a tetraciclina y minociclina en aislamientos clínicos de *Acinetobacter baumannii*. *Enferm. Infecc Microbiol Clin* 2006; 24(2):77–80. <https://doi.org/10.1157/13085012>
- Medina A, Horcajo P, Jurado S, de la Fuente R, Ruiz-Santana-Quinteira JA, Dominguez-Bernal G, Orden JA. Phenotypic and genotypic characterization of antimicrobial resistance in enterohemorrhagic *Escherichia coli* and atypical enteropathogenic *E. coli* strains from ruminants. *J Vet Diagn Invest* 2011; 23(1):91–95. <https://doi.org/10.1177/104063871102300114>
- Memariani M, Najar-peerayeh S, Salehi TZ, Khalil S. Occurrence of SHV, TEM and CTX-M β -Lactamase genes among enteropathogenic *Escherichia coli* strains isolated from children with diarrhea. *Jundishapur J Microbiol* 2015; 8(4):e15620. [https://doi.org/10.5812/jjm.8\(4\)2015.15620](https://doi.org/10.5812/jjm.8(4)2015.15620)
- Mirzaagha P, Louie M, Sharma R, Yanke LJ, Topp E, McAllister TA. Distribution and characterization of ampicillin- and tetracycline-resistant *Escherichia coli* from feedlot cattle fed subtherapeutic antimicrobials. *BMC Microbiol* 2011; 11:78(2011). <https://doi.org/10.1186/1471-2180-11-78>
- Mosquito S, Ruiz J, Bauer JL, Ochoa TJ. Molecular mechanisms of antibiotic resistance in, associated diarrhea *Rev Peru. Med. Exp. Salud Publica* 2011; 28(4):648–656. <https://doi.org/10.17843/rpmesp.2011.284.430>
- Murray A, Mather H, Coia JE, Brown DJ. Plasmid-mediated quinolone resistance in nalidixic-acid-susceptible strains of *Salmonella enterica* isolated in Scotland. *J Antimicrob Chemother* 2008; 62(5):1153–1155. <https://doi.org/10.1093/jac/dkn340>
- O'Connor AM, Shen HG, Wang C, Opiessnig T. Descriptive epidemiology of *Moraxella bovis*, *Moraxella bovoculi* and *Moraxella ovis* in beef calves with naturally occurring infectious bovine keratoconjunctivitis (Pinkeye). *Vet Microbiol* 2012; 155(2–4):374–380. <https://doi.org/10.1016/j.vetmic.2011.09.011>
- Park Y, Yu JK, Lee S, Oh E, Woo G. Prevalence and diversity of qnr alleles in AmpC-producing *Enterobacter cloacae*, *Enterobacter aerogenes*, *Citrobacter freundii* and *Serratia marcescens*: a multicentre study from Korea. *J Antimicrob Chemother* 2007; 60(4):868–871. <https://doi.org/10.1093/jac/dkm266>
- Peymani A, Naserpour-Farivar T, Zare E, Azarhoosh KH. Distribution of blaTEM, blaSHV, and blaCTX-M genes among ESBL-producing *P. aeruginosa* isolated from Qazvin and Tehran hospitals, Iran. *J Prev Med Hyg* 2017;58(2): 55–160. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5584084/pdf/2421-4248-58-E155.pdf>

- Roberts MC, Smith AL. Molecular Characterization of “Plasmid-Free” Antibiotic resistant *Haemophilus influenzae*. J Bacteriol 1980; 144(1):476–479. <https://doi.org/10.1128/jb.144.1.476-479.1980>
- Roberts MC, Pang Y, Spencer RC, Winstanley TG, Brown BA, Wallace RJ. Tetracycline resistance in *Moraxella (Branhamella) catarrhalis*: Demonstration of two clonal outbreaks by using pulsed-field gel electrophoresis. Antimicrob Agents Chemother 1991; 35(11):2453–2455. <https://doi.org/10.1128/AAC.35.11.2453>
- Roberts MC, Chung WO, Roe DE. Characterization of tetracycline and erythromycin resistance determinants in *Treponema denticola*. Antimicrob Agents Chemother 1996; 40(7):1690–1694. <https://doi.org/10.1128/AAC.40.7.1690>
- Rodríguez-Martínez JM, Pascua A, García I, Martínez-Martínez L. Detection of the plasmid-mediated quinolone resistance determinant qnr among clinical isolates of *Klebsiella pneumoniae* producing AmpC-type β -lactamase. J Antimicrob Chemother 2003; 52:703–706. <https://doi.org/10.1093/jac/dkg388>
- Roe DE, Braham PH, Weinberg A, Roberts MC. Characterization of tetracycline resistance in *Actinobacillus actinomycetemcomitans*. Oral Microbiol Immunol 1995; 10(4):227–232. <https://doi.org/10.1111/j.1399-302X.1995.tb00147.x>
- Schwarz S, Kehrenberg C, Doublet B, Cloeckaert A. Molecular basis of bacterial resistance to chloramphenicol and florfenicol. FEMS Microbiol Rev 2004(5); 28:519–542. <https://doi.org/10.1016/j.femsre.2004.04.001>
- Shen HG, Gould S, Kinyon J, Opriessnig T, O’Connor AM. Development and evaluation of a multiplex real-time PCR assay for the detection and differentiation of *Moraxella bovis*, *Moraxella bovoculi* and *Moraxella ovis* in pure culture isolates and lacrimal swabs collected from conventionally raised cattle. J Appl Microbiol 2011; 111(5):1037–1043. <https://doi.org/10.1111/j.1365-2672.2011.05123.x>
- Sosa V, Zunino P. Diversity of *Moraxella* spp. strains recovered from infectious bovine keratoconjunctivitis cases in Uruguay. J Infect Dev Ctries 2013; 7(11):819–824. <https://doi.org/10.3855/jidc.3458>
- Talavera-González JM, Acosta-Dibarrat J, Reyes-Rodríguez NE; Salgado-Miranda C, Talavera-Rojas M. Prevalence of the qnrB, qnrA and blaTEM genes in temperate bacteriophages of *Escherichia coli* isolated from wastewater and sewer water from slaughterhouses in the State of Mexico. Rev Mex Cienc Pecu 2021; 12(1):298–305. <https://doi.org/10.22319/rmcp.v12i1.5378>
- Touati A, Brasme L, Benallaoua S, Gharout A, Madoux J, De Champs C. First report of qnrB-producing *Enterobacter cloacae* and qnrA-producing *Acinetobacter baumannii* recovered from Algerian hospitals. Diagn Microbiol Infect Dis 2008; 60(3):287–290. <https://doi.org/10.1016/j.diagmicrobio.2007.10.002>
- Wang M, Tran J, Jacoby G. Plasmid-mediated quinolone resistance in clinical isolates of *Escherichia coli* from Shanghai, China. Antimicrob Agents Chemother 2003; 47(7):2242–2248. <https://doi.org/10.1128/AAC.47.7.2242-2248.2003>
- Wu J, Ko W, Tsai S, Yan J. Prevalence of plasmid-mediated quinolone resistance determinants QnrA, QnrB, and QnrS among clinical isolates of *Enterobacter cloacae* in a Taiwanese Hospital. Antimicrob Agents Chemother 2007; 51(4): 1223–1227. <https://doi.org/10.1128/AAC.01195-06>