

Subject index for Volume 36, 2023

- Aelurostrongylus abstrusus*; 36(1):13–21
- Agroindustrial residue; 36(1):44–54
- Alternative forages; 36(1):33–43; 36(2):89–97; 36(2):98–108
- Ammonium formate; 36(2):55–65; 36(2):55–65
- Artificial insemination; 36(3):141–151
- Available phosphorus; 36(2):66–79; 36(3):121–130
- Avermectins; 36(4):210–215
- Backyard ducks; 36(3):131–140
- Bacteria; 36(1):22–32
- Behavior; 36(4):161–171
- Biogas; 36(1):22–32
- Biomarker; 36(4):172–180
- Blood pH; 36(2):55–65
- Bone health; 36(2):66–79; 36(3):121–130
- Bovine; 36(1):22–32
- Broiler; 36(2):66–79; 36(3):121–130; 36(4):181–195
- Broilers; 36(2):55–65
- Brown Swiss; 36(4):161–171
- Buffalo; 36(1):22–32
- Cairina moschata*; 36(3):131–140
- Caprine; 36(1):33–43; 36(2):89–97
- Caprines; 36(2):80–88
- Carcass weight; 36(4):196–209
- Carcass yield; 36(4):196–209
- Cassava; 36(1):44–54
- Cattle farms; 36(3):141–151
- Cattle; 36(4):161–171
- Cellulolytic bacteria; 36(1):22–32
- Chemical composition; 36(1):33–43; 36(1):44–54; 36(2):89–97
- Chickpea; 36(1):33–43; 36(2):89–97
- Coculture; 36(1):22–32
- Conserved forage; 36(1):44–54
- Cooling losses; 36(4):196–209
- Coprophagous beetle; 36(1):3–12
- Corticosterone; 36(3):152–160
- Cow; 36(3):141–151
- Creole; 36(2):80–88
- Crocodile farming; 36(3):152–160
- Crocodylians; 36(3):152–160
- Dairy business; 36(3):141–151
- Dairy calves; 36(4):161–171
- Dairy cattle; 36(3):141–151
- Dairy cow; 36(3):141–151
- Dairy farm; 36(3):141–151
- Dairy goats; 36(2):98–108
- Dairy herd; 36(3):141–151
- Degradability; 36(1):44–54
- Dog; 36(3):109–120; 36(4):172–180
- Drinking water; 36(2):55–65

- Drylands; 36(4):196–209
- Duck farming; 36(3):131–140
- Dung beetle; 36(1):3–12
- Dung insect; 36(1):3–12
- Dung; 36(1):3–12
- Dung-associated invertebrates; 36(1):3–12
- Economic performance; 36(2):98–108; 36(4):196–209
- Energy sources; 36(4):196–209
- Energy; 36(4):196–209
- Estrogen; 36(3):152–160
- Ewe meat; 36(4):196–209
- Ewes; 36(4):196–209
- Exogenous enzymes; 36(2):66–79; 36(3):121–130
- Farm productivity; 36(3):141–151
- Farm size; 36(3):141–151
- Fat; 36(4):196–209
- FCR; 36(3):131–140
- Feed alternatives; 36(4):196–209
- Feed conversion; 36(2):66–79; 36(3):121–130
- Feed efficiency; 36(4):161–171
- Feed; 36(1):33–43; 36(2):89–97
- Feeding cost; 36(4):161–171
- Fermentation characteristics; 36(1):22–32
- Fiber degradation; 36(1):22–32
- Fiber; 36(1):22–32
- Fibrinolytic; 36(3):109–120
- Formic acid; 36(2):55–65
- French-Alpine goats; 36(2):80–88
- Fresh cheese; 36(2):98–108
- Gas production; 36(1):22–32
- Genetic improvement; 36(3):141–151
- Glaucoma; 36(3):109–120
- Goat milk; 36(2):80–88
- Goats; 36(1):33–43
- Goats; 36(2):80–88
- Goats; 36(2):89–97
- Grape marc; 36(1):44–54
- Grazing goats; 36(2):80–88
- Growth performance; 36(2):55–65; 36(2):66–79; 36(3):121–130; 36(3):131–140; 36(4):161–171
- Hay; 36(1):33–43; 36(2):89–97
- Herd size; 36(3):141–151
- Holstein-Friesian; 36(4):161–171
- Immunity; 36(4):181–195; 36(4):210–215
- Immunoglobulin E; 36(4):210–215
- Immunoglobulin G; 36(4):210–215
- Immunoglobulin M; 36(4):210–215
- Infestation; 36(4):172–180
- Insemination; 36(3):141–151
- Interferon- α ; 36(4):210–215
- Interleukin-2; 36(4):210–215
- Interleukin-6; 36(4):210–215
- Interleukin-10; 36(4):210–215
- Intestinal morphology; 36(4):181–195
- Intestinal morphometry; 36(2):55–65

- Intraocular pressure; 36(3):109–120
- In vitro* fermentation; 36(1):22–32
- Ivermectin; 36(1):3–12; 36(4):210–215
- Lactose; 36(2):80–88
- Legumes; 36(1):33–43; 36(2):89–97
- Leopardus tigrinus*; 36(1):13–21
- Licuri; 36(4):196–209
- Little spotted cat; 36(1):13–21
- Little tiger cat; 36(1):13–21
- Loin; 36(4):196–209
- Low phosphorus diet; 36(2):66–79; 36(3):121–130
- Lung parasitism; 36(1):13–21
- Lung; 36(1):13–21
- Lungworm; 36(1):13–21
- Management; 36(3):131–140
- Manihot esculenta*; 36(1):44–54
- Manure beetle; 36(1):3–12
- Methane; 36(1):22–32
- Mexico; 36(1):33–43; 36(2):89–97
- Milk fat; 36(2):80–88
- Milk price; 36(3):141–151
- Milk production; 36(3):141–151
- Milk protein; 36(2):80–88
- Milk quality; 36(2):80–88
- Milk; 36(1):33–43; 36(2):89–97; 36(2):98–108; 36(3):141–151
- Mineral; 36(2):66–79; 36(3):121–130
- Multiparasitism; 36(1):13–21
- Multivariate analysis; 36(3):141–151
- Muscovy ducks; 36(3):131–140
- Nanominerals; 36(4):181–195
- Nanovitamins; 36(4):181–195
- Nematode; 36(1):13–21
- Non-fat solids; 36(2):80–88
- Nutrition; 36(2):66–79; 36(3):121–130
- Ocular disease; 36(3):109–120
- Ocular hypertension; 36(3):109–120
- Oncilla; 36(1):13–21
- On-farm research; 36(2):98–108
- Oxidative stress; 36(4):172–180
- Parasite resistance; 36(1):3–12
- Parasites; 36(1):13–21
- Phytase; 36(2):66–79; 36(3):121–130
- Pneumonia; 36(1):13–21
- Poultry; 36(2):66–79; 36(3):121–130; 36(4):181–195
- Probiotics; 36(3):131–140
- Progesterone; 36(3):152–160
- Propionic acid; 36(2):55–65
- Reproduction; 36(3):141–151
- Reproductive biotechnologies; 36(3):141–151
- Rumen; 36(1):22–32
- Ruminal bacteria; 36(1):22–32
- Saanen; 36(2):80–88
- Salts; 36(2):80–88
- Sarcoptes*; 36(4):172–180

- Sarcoptic mange; 36(4):172–180
- Scabies; 36(4):172–180
- Seasonality; 36(2):80–88
- Selenium; 36(4):181–195
- Sensory analysis; 36(2):98–108
- Sex hormones; 36(3):152–160
- Sheep; 36(4):210–215
- Silage; 36(1):44–54
- Small ruminants; 36(4):196–209
- Small-scale livestock systems; 36(2):98–108
- Stress; 36(3):152–160
- Sunflower; 36(1):33–43; 36(2):89–97
- Sunflower-chickpea hay; 36(2):98–108
- Testosterone; 36(3):152–160
- Tocopherol; 36(4):181–195
- Trace minerals; 36(4):181–195
- Tumor necrosis factor- α ; 36(4):210–215
- Vegetable oil; 36(4):196–209
- Vitamin E; 36(4):181–195
- Vitis vinifera*; 36(1):44–54
- Weaning weight; 36(4):161–171
- Weight gain; 36(2):66–79; 36(3):121–130
- Wildlife; 36(1):13–21
- Yucca; 36(1):44–54
- Zoonosis; 36(4):172–180