

Subject index for Volume 37, 2024

- Abortion; 37(1):52–59
- Acetic acid; 37(1):27–41
- Acute hepatopancreatic necrosis disease; 37(3):185–193
- Additives; 37(1):27–41; 42–51
- Age at first calving; 37(4):233–242
- Aloe; 37(3):185–193
- Animal food product; 37(3):163–184
- Animal; 37(3):163–184
- Antibiotic resistance; 37(3):163–184
- Antibiotics; 37(3):163–184
- Anticoccidial; 37(4):195–207
- Antimicrobial resistance; 37(1):14–26
- Antioxidants; 37(4):223–232
- Antiparasitic drugs; 37(2):101–112
- Artificial insemination; 37(4):223–232; 233–242
- Bacillus licheniformis*; 37(3):185–193
- Backgrounding systems; 37(1):42–51
- Bacterial resistance; 37(3):163–184
- Basil; 37(3):185–193
- Beef cattle; 37(4):233–242
- Beef quality; 37(1):3–13
- Biometric measurements; 37(4):243–250
- Body condition; 37(2):88–100
- Body size; 37(3):152–162
- Body weight; 37(4):243–250
- Bovine; 37(2):101–112; 113–121; 37(4):233–242
- Breaking strength; 37(2):61–72
- Broiler; 37(2):61–72; 37(4):195–207
- Bubalus bubalis*; 37(4):243–250
- Buffalo; 37(4):243–250
- Buffaloes; 37(3):135–151
- Calicophoron microbothrioides*; 37(2):113–121
- Calves; 37(3):135–151
- Carcass traits; 37(1):3–13; 37(4):195–207
- Carcass yield; 37(4):195–207
- Cattle; 37(1):3–13; 37(2):88–100; 113–121; 37(3):123–134; 135–151
- Chemical composition; 37(1):27–41
- Coinfection; 37(4):208–222
- Colombia; 37(3):135–151; 163–184
- Color parameters; 37(1):3–13
- Common opossum; 37(3):152–162
- Condemnation; 37(3):135–151
- Contagious ovine keratoconjunctivitis; 37(1):14–26
- Cow; 37(2):73–87; 88–100
- Cow-calf operations; 37(1):42–51
- Coxiella burnetii*; 37(1):52–59
- Criollo lechero tropical; 37(3):123–134
- Crossbred; 37(2):88–100
- Dairy cattle; 37(2):101–112

- Diagnosis; 37(1):52–59; 37(2):113–121
- Didelphis marsupialis*; 37(3):152–162; 37(4):208–222
- Diet; 37(1):27–41
- Distomatosis; 37(3):135–151
- DNA markers; 37(3):123–134
- Drone; 37(4):243–250
- Drug-resistant bacteria; 37(3):163–184
- Dry matter; 37(1):27–41
- Dual-purpose; 37(2):88–100
- Effective population size; 37(3):123–134
- Eggs; 37(2):113–121
- Ensilage; 37(1):27–41
- Ensilaged forages; 37(1):42–51
- Enterobacteriaceae; 37(3):163–184
- Epidemiological surveillance; 37(1):14–26
- Epidemiology; 37(1):52–59; 37(2):113–121
- Erythrina americana*; 37(4):223–232
- Expected progeny differences; 37(4):233–242
- Farm; 37(2):101–112
- Fasciola hepatica*; 37(2):101–112; 37(3):135–151
- Fascioliasis; 37(2):101–112
- Fasciolosis; 37(3):135–151
- Feed restriction; 37(4):195–207
- Feeding phases; 37(4):195–207
- Fermentative profile; 37(1):27–41
- Forage conservation; 37(1):27–41
- Forage; 37(1):27–41
- Fresh semen; 37(4):223–232
- Gait score; 37(2):61–72
- Garlic; 37(3):185–193
- Gas production; 37(1):27–41
- Gastrocnemius; 37(2):61–72
- Genetic diversity; 37(3):123–134
- Ginger; 37(3):185–193
- Goat; 37(1):14–26
- Grape seed extract; 37(2):61–72
- Green micro-algae; 37(1):42–51
- Growth performance; 37(4):195–207
- Growth; 37(4):243–250
- Heces; 37(2):113–121
- Helminthiasis; 37(4):208–222
- Holstein Friesian; 37(1):3–13; 37(2):73–87
- Human-opossum conflict; 37(3):152–162
- Inbreeding; 37(3):123–134
- Infection frequency; 37(3):135–151
- Ionophores; 37(4):195–207
- Lactation; 37(2):88–100
- Leg health; 37(2):61–72
- Litopenaeus vannamei*; 37(3):185–193
- Litter size; 37(3):152–162
- Live weight; 37(4):243–250
- Livestock; 37(1):52–59
- Livestock 4.0; 37(4):243–250
- Low-protein diets; 37(1):42–51
- Marsupial; 37(3):152–162

- Marsupialia; 37(4):208–222
- Mathematical models; 37(4):243–250
- Methanogenesis; 37(1):42–51
- Mexico; 37(1):52–59
- Milk composition; 37(2):73–87; 88–100
- Milk yield; 37(2):73–87; 88–100
- Moraxella* spp.; 37(1):14–26
- Multiresistance; 37(1):14–26
- Non-casing components; 37(1):3–13
- Non-genetic factors; 37(2):73–87
- Nutrient restriction; 37(2):61–72
- Opossum; 37(4):208–222
- Quality; 37(2):88–100
- Paramphistomosis; 37(2):113–121
- Parasite load; 37(2):113–121
- Parasites; 37(4):208–222
- Parasitic diseases; 37(4):208–222
- Parasitology; 37(2):101–112
- Parasitosis; 37(3):135–151
- Pathogen resistance; 37(3):163–184
- Performance; 37(2):61–72
- Phenotypic correlation; 37(2):73–87
- Population structure; 37(3):123–134
- Precision livestock farming; 37(4):243–250
- Price formation; 37(4):233–242
- Protected areas; 37(4):208–222
- Protein supplementation; 37(1):42–51
- Q fever; 37(1):52–59
- Rams; 37(4):223–232
- Refrigerated semen; 37(4):223–232
- Reproductive failure; 37(1):52–59
- Resistance genes; 37(1):14–26
- Resistant bacteria; 37(3):163–184
- Romosinuano; 37(3):123–134
- Rumen fluke; 37(2):113–121
- Ruminants; 37(1):52–59; 27–41; 37(2):113–121
- Salmonella*; 37(3):163–184
- Semen market; 37(4):233–242
- Semen price; 37(4):233–242
- Sexual dimorphism; 37(3):152–162
- Sheep; 37(1):14–26; 37(4):223–232
- Shrimp; 37(3):185–193
- Skin color; 37(4):195–207
- Skin pigmentation; 37(4):195–207
- Slaughter age; 37(1):3–13
- Slaughter characteristics; 37(1):3–13
- Slaughterhouse; 37(3):135–151
- Somatic cell count; 37(2):73–87
- Spermatozoa; 37(4):223–232
- Staphylococcus aureus*; 37(3):163–184
- Super-bacteria; 37(3):163–184
- Tendon; 37(2):61–72
- Tibia; 37(2):61–72
- Trematode; 37(2):101–112; 113–121
- Types of muscle; 37(1):3–13
- Unmanned aerial vehicle; 37(4):243–250

Urban parks; 37(4):208–222

Urban wildlife; 37(3):152–162

Vaginal swabs; 37(1):52–59

Vibrio parahaemolyticus; 37(3):185–193

Vitamin E; 37(2):61–72

Weaning weight; 37(4):233–242

Weight prediction; 37(4):243–250

Wildlife; 37(4):208–222