

CYTOTOXIC LABDANE DITERPENOIDS ISOLATED FROM THE HEXANE FRACTION OF THE *Croton stipuliformis* STEM BARK

DITERPENOIDES LABDANOS CITOTÓXICOS AISLADOS DE LA FRACCIÓN
HEXÁNICA DE LA CORTEZA DE *Croton stipuliformis*

Martha S. FRANCO¹, Claudia P. CORDERO², Sandra J. MORANTES²,
Fabio ARISTIZABAL², Coralia OSORIO^{1*}

Received: 02 September 2010 Accepted: 18 July 2011

ABSTRACT

The composition of the cytotoxic hexane fraction of the *Croton stipuliformis* bark was studied by means of a gas chromatography coupled to a mass spectrometry; sesquiterpenes and sterols were characterized as the main constituents. Along with these compounds, three labdane and *seco*-labdane diterpenoids, 8(17),12*E*,14-labdatrien-18-oic acid [3], 12*E*-3,4-*seco*-labda-4(18),8(17),12,14-tetraen-3-oic acid [2], and its methyl ester [1] were isolated from this fraction through preparative HPLC, and their structures were elucidated by HR-FABMS, 1D and 2D NMR analyses. Additionally, the cytotoxic activity of these three compounds against human tumor cell lines HEp-2, HT-29, MKN-45, MCF-7, and HeLa was assessed. The three compounds showed a non-specific and moderate cytotoxicity against the above-mentioned cell lines.

Keywords: *Croton stipuliformis*, Euphorbiaceae, labdane, MTT assay, cytotoxicity.

RESUMEN

La composición de la fracción hexánica de la corteza de *Croton stipuliformis* se estudió por cromatografía de gases acoplada a espectrometría de masas; se caracterizaron los sesquiterpenos y esteroides como los principales constituyentes. Adicionalmente a estos compuestos, tres diterpenoides labdanos y *seco*-labdanos, el ácido 8(17),12*E*,14-labdatrien-18-oico [3], el ácido 12*E*-3,4-*seco*-labda-4(18),8(17),12,14-tetraen-3-oico [2] y su ester metílico [1] se aislaron de esta fracción por HPLC preparativa y sus estructuras se elucidaron con base en los análisis por HRFABMS, espectroscopía de RMN mono y bidimensional. Adicionalmente, se determinó la actividad citotóxica de estos tres compuestos frente a las líneas celulares de tumores humanos HEp-2, HT-29, MKN-45, MCF-7 y HeLa. Todos los compuestos mostraron una citotoxicidad moderada y no específica frente a las líneas celulares mencionadas anteriormente.

Palabras Clave: *Croton stipuliformis*, Euphorbiaceae, labdano, ensayo MTT, citotoxicidad.

¹ Departamento de Química. Universidad Nacional de Colombia. AA 14490 Bogotá, Colombia.

² Departamento de Farmacia. Universidad Nacional de Colombia. AA 14490 Bogotá, Colombia.

* Corresponding author: cosorior@unal.edu.co

INTRODUCTION

Euphorbiaceae is a big family of about 8,000 species that usually grow in tropical regions. In Colombia, this family is comprised of 78 genera and 390 species, from which 80 species belong to the *Croton* genus (1). *Croton stipuliformis* (Euphorbiaceae) is a species native to Colombia and it grows under mild climates between 500 and 1800 meters above sea level in the Andes area at the west-central region of Colombia. It is a tree that reaches a height of 25–30 m, and it is commonly known as “guacamayo,” “candelero” or “mopo.” Usually, it is confused with *C. cupreatus*, a phenotypically close species (2). In the previously mentioned region, this native plant is recognized because its bark produces a yellow-orange latex called “Sangre de drago,” which is used for different pharmacological purposes, such as ulcer treatment, pain, anaemia, and cancer (3).

The latex exuded by several *Croton* species has shown cytotoxic activity against various human tumor cell lines (3, 4). Several classes of diterpenoids have been identified as responsible for this activity; among which, labdanes, cembranoids, traquilobanes, pimaranes, *ent*-kauranes, and clerodanes, are the most remarkable.

As part of our current studies on the bioactive compounds found in Colombian native plants (5–7), the aim of this research was to perform a bioguided fractionation of the hexane fraction from the bark of *Croton stipuliformis*, in order to isolate and characterize the compounds responsible for its cytotoxic activity.

MATERIALS AND METHODS

General

^1H - and ^{13}C NMR (400 and 100 MHz, respectively) spectra were acquired on a Bruker Biospin[®] 400 spectrometer (Karlsruhe, Germany). NMR spectra were recorded in CDCl_3 , and referenced to a TMS signal. HRFAB-MS spectra were recorded in a glycerol matrix in the positive ion mode with a Micromass AutoSpec[®]-Q spectrometer (Manchester, UK). GC-MS and direct inlet-MS (EIMS, 70 eV) were carried out on a Shimadzu[®] GCMS QP5050 instrument. The analytical HPLC was performed on a Merck-Hitachi[®] system equipped with a L-6200 A pump, a L-4500 diode array detector and a D-6000 A unit (Darmstadt, Germany); while for the preparative HPLC, a L-4250 UV-Vis detec-

tor was used. Silica gel (0.063–0.200 mm, Merck) was used for column chromatography, and the TLC was performed on silica gel GF₂₅₄ plates (0.10 mm, Merck[®]). Optical rotations were measured with a Polartronic E, Schmidt & Haensch[®] polarimeter (Berlin, Germany). The solvents were purchased from Merck[®] (Darmstadt, Germany).

Plant material

The *C. stipuliformis* stem bark was collected at Chinchiná, Caldas, Colombia. A voucher specimen (COL 512797) was identified by Dr. J. Murillo and deposited at the Instituto de Ciencias Naturales, Universidad Nacional de Colombia.

Plant material extraction

The dried *C. stipuliformis* stem bark (3.2 kg) was extracted with methanol (1.5 L, two times) at 60°C during 4 h period of time. The methanol was evaporated in vacuo to obtain a crude extract (207 g). A part of the methanolic extract (115 g) was suspended in water (500 mL) and extracted with dichloromethane (150 mL x 3); the two phases were separated and after their concentration, organic (CSO, 39.6 g) and aqueous (CSA, 52.9 g) fractions were obtained. A portion of the organic fraction (12.4 g) was extracted with hexane (60 mL x 3) to obtain the CSH fraction (5.2 g).

Fractionation and purification of cytotoxic compounds

The bioactive hexane fraction (695 mg) was subjected to open silica gel column chromatography (70 g), sequentially eluting it with 200 mL each of hexane:AcOEt in ratios of 10:0, 7:3, 5:5, 3:7 and 0:10 to afford four pooled fractions as follows: CSHa (15 mg), CSHb (150 mg), CSHc (182 mg), and CSHd (180 mg). The fractions were pooled together based on their similar R_f values on thin layer chromatography (TLC), which was developed with the mobile phase system of hexane:AcOEt (1:1); then, they were sprayed using an anisaldehyde:sulfuric acid solution (1%: 2% v/v) in EtOH, followed by heating at 110°C. All of the fractions were analyzed through GC-MS, and CSHb and CSHc fractions were also analyzed by HPLC-DAD on a Synergi[™] 4 μ MAX-RP 80A column (4.0 μm , 250 mm x 4.60 mm i.d., Phenomenex[®]), using a methanol:acetonitrile gradient as mobile phase.

CSHb and CSHc fractions were further subjected to preparative HPLC using a Gemini™ 5 μ C18 column (5 μ m, 250mm x 10mm i.d., Phenomenex®), with methanol:acetonitrile (9:1) as mobile phase at a flow rate of 5 mL/min. Thus, the CSHb fraction only afforded compound **1** (105 mg), while the CSHc fraction yielded compounds **1** (90 mg), **2** (28 mg), and **3** (24 mg).

12E-3,4-*seco*-labda-4(18),8(17),12,14-tetraen-3-oic acid methyl ester (**1**) was obtained as a yellow oil. $[\alpha]_D^{25} = +13.0^\circ$ (CH₂Cl₂; *c* 0.04). EIMS *m/z* {rel. int.}: 316 [M⁺] {10}, 301 [M-CH₃]⁺ {7}, 285 [M-CH₃O]⁺ {5}, 273 [M-C₃H₇]⁺ {45}, 259 {7}, 247 {10}, 229 [273-CO₂]⁺ {40}, 219 {10}, 201 [229-CH₂CH₂]⁺ {25}, 187 {20}, 173 {40}, 161 {50}, 147 {35}, 145 {42}, 133 {48}, 119 {70}, 105 {60}, 93 {55}, 91 {75}, 81 {80}, 79 {100}, 77 {52}, 67 {58}, 55 {90}, 53 {52}. HRFABMS found 317.2481 [M+H]⁺, C₂₁H₃₃O₂ requires 317.2474. ¹H NMR (400 MHz, CDCl₃): δ 0.75 (3H, s, H₃-20), 1.53-1.61 (1H, m, H-6a), 1.72 (2H, overlapped, H-6b, H-11a), 1.72 (3H, s, H₃-19), 1.74 (1H, overlapped, H-1a), 1.76 (3H, s, H₃-16), 1.77 (1H, overlapped, H-1b), 1.88 (1H, br t, *J* = 6.2 Hz, H-9), 1.99 (1H, dt, *J* = 12.8, 4.8 Hz, H-7a), 2.19 (1H, overlapped, H-2a), 2.24 (1H, dd, *J* = 12.0, 3.2 Hz, H-5), 2.25 (1H, overlapped, H-11b), 2.34 (1H, ddd, *J* = 12.8, 4.0, 2.4 Hz, H-7b), 2.44 (1H, ddd, *J* = 13.8, 11.6, 5.6 Hz, H-2b), 3.64 (3H, s, OCH₃), 4.48 (1H, br s, H-17a), 4.68 (1H, br s, H-18a), 4.85 (1H, br s, H-18b), 4.86 (1H, d, *J* = 1.8 Hz, H-17b), 4.88 (1H, d, *J* = 10.8 Hz, H-15a), 5.04 (1H, d, *J* = 17.4 Hz, H-15b), 5.34 (1H, t, *J* = 6.0 Hz, H-12), 6.31 (1H, dd, *J* = 17.4, 10.7, H-14); ¹³C NMR (100 MHz, CDCl₃): δ 12.1 (C-16), 17.6 (C-20), 23.6 (C-11), 23.7 (C-19), 28.1 (C-2), 30.2 (C-6), 32.7 (C-1), 37.7 (C-7), 41.4 (C-10), 49.3 (C-9), 51.0 (C-5), 51.8 (OCH₃), 108.7 (C-17), 110.4 (C-15), 113.9 (C-18), 133.2 (C-12), 133.9 (C-13), 141.6 (C-14), 147.2 (C-4), 147.5 (C-8), 174.4 (C-3).

12E-3,4-*seco*-labda-4(18),8(17),12,14-tetraen-3-oic acid (**2**) was obtained as a colorless oil. $[\alpha]_D^{25} = -55.6^\circ$ (CH₂Cl₂; *c* 0.06). EIMS *m/z* {rel. int.}: 302 [M⁺] {10}, 287 [M-CH₃]⁺ {8}, 273 [M-29]⁺ {5}, 259 [M-C₃H₇]⁺ {15}, 247 {3}, 245 {8}, 229 [M-COOH-C₂H₄]⁺ {30}, 219 {10}, 203 {20}, 201 {20}, 187 {20}, 173 {23}, 161 {50}, 147 {40}, 133 {45}, 131 {22}, 119 {70}, 107 {67}, 105 {70}, 93 {73}, 91 {75}, 81 {80}, 79 {100}, 77 {49}, 69 {39}, 67 {39}, 65 {25}, 55 {70}, 53 {48}. HRFABMS found 301.1233 [M-H]⁺, C₂₀H₂₉O₂ requires 301.1229. ¹H NMR (400 MHz,

CDCl₃): δ 0.75 (3H, s, H₃-20), 1.58 (1H, m, H-6a), 1.71 (1H, overlapped, H6b), 1.72 (1H, overlapped, H-11a), 1.72 (3H, s, H₃-19), 1.74 (1H, overlapped, H-1a), 1.75 (3H, br s, H₃-16), 1.76 (1H, overlapped, H-1b), 1.87 (1H, br t, *J* = 6.8 Hz, H-9), 1.98 (1H, dt, *J* = 12.8, 5.2 Hz, H-7a), 2.20-2.24 (1H, m, H-11b), 2.21 (1H, dd, *J* = 12.4, 3.6 Hz, H-5), 2.25 (1H, ddd, *J* = 15.2, 12.4, 5.6 Hz, H-2a), 2.34 (1H, ddd, *J* = 12.8, 4.0, 2.4 Hz, H-7b), 2.46 (1H, ddd, *J* = 15.2, 12.0, 5.2 Hz, H-2b), 4.47 (1H, br s, H-17a), 4.68 (1H, br s, H-18a), 4.85 (1H, br s, H-18b), 4.86 (1H, d, *J* = 1.8 Hz, H-17b), 4.88 (1H, d, *J* = 10.4 Hz, H-15a), 5.04 (1H, d, *J* = 17.6 Hz, H-15b), 5.33 (1H, t, *J* = 6.4 Hz, H-12), 6.30 (1H, dd, *J* = 17.2, 11.2, H-14); ¹³C NMR (100 MHz, CDCl₃): δ 12.0 (C-16), 17.5 (C-20), 23.4 (C-11), 23.5 (C-19), 28.0 (C-2), 30.0 (C-6), 32.3 (C-1), 37.5 (C-7), 41.2 (C-10), 49.2 (C-9), 50.8 (C-5), 108.6 (C-17), 110.3 (C-15), 113.8 (C-18), 132.8 (C-12), 133.9 (C-13), 141.4 (C-14), 147.0 (C-8), 147-2 (C-4), 180.1 (C-3).

8(17),12E,14-labdatrien-18-oic acid (**3**) was obtained as a white solid. $[\alpha]_D^{25} = -15.4^\circ$ (CH₂Cl₂; *c* 0.03). EIMS *m/z* {rel. int.}: 302 [M⁺] {25}, 287 [M-CH₃]⁺ {40}, 273 [M-29]⁺ {10}, 257 [M-COOH]⁺ {10}, 246 {38}, 241 {22}, 231 {27}, 221 {8}, 201 {20}, 185 {15}, 175 {70}, 161 {20}, 147 {48}, 135 {53}, 134 {45}, 121 {52}, 119 {70}, 107 {55}, 105 {54}, 93 {75}, 91 {80}, 81 {90}, 79 {100}, 77 {40}, 67 {43}, 55 {52}, 53 {45}. The HRFABMS gave 301.2149 [M-H]⁺ as result, which is compatible with the molecular formula C₂₀H₂₉O₂. ¹H and ¹³C-NMR data were in agreement with those previously published by Smith *et al.*, 2007 (8).

GCMS analyses

HRGC-EIMS analyses were carried out on a RTX-5 column (30 m x 0.25 mm i.d., 0.25 μ m film thickness). The column oven was maintained at 60°C for 4 min and then programmed to increase the temperature from 60 to 300°C at 4°C/min. The final temperature was held for 15 min, and the injector temperature was maintained at 300°C. Helium was used as carrier gas at 1.0 mL/min; injection volume was 1 μ L in split mode (1:10). MS data were recorded in a mass range of 30 - 500 U, with an electron energy of 70 eV, and they were processed by a Class 5000 v 2.2 MS-Workstation® software. The CSHa fraction was also analyzed on a DB-Wax column (30 m x 0.25 mm i.d., 0.25 μ m film thickness). The injector port temperature was maintained at 230°C and the

split ratio was set to 1:10. The oven temperature was initially fixed at 50°C for 4 min, then it was increased at 4°C/min until reaching 200°C, and then it was maintained at that level for 10 min. Helium was the carrier gas used at 1.0 mL/min.

Kovats retention indices on the two previously described different columns were calculated and compared with the indices found in the literature values (9). The constituents of the CSHa and CSHd fractions were identified by matching their mass spectra with those from known databases (10-12).

Biological evaluation

A bioassay of the cytotoxic activity against human tumor cell culture *in vitro* was performed by means of the MTT (3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide) colorimetric assay (13, 14). Cancer cell lines of human larynx (HEp-2), colon (HT-29), gastric (MKN-45), breast (MCF-7) and cervical cells (HeLa) were obtained from the American Type Culture Collection (ATCC®, Rockville, MD). The cells were cultured in MEM (Minimum Essential Medium, SIGMA®) supplemented with 5% fetal bovine serum (Vitacell, ATCC, VA) and gentamycin (50 µg/mL).

For the experiments, the cells were plated in 100 µL of culture medium in 96-well; after 24 h, the fractions and pure compounds (1 - 3) were added to each well at different concentrations (50, 5, and 0.5 µg/mL in DMSO), and then they were incubated for 48 h in a total of 200 µL of medium volume. Each treatment was evaluated in triplicate. At the end of the treatment, the medium in each well was replaced with fresh medium (100 µL) containing 0.25 mg/mL of MTT. Four hours later, the formazan product of the MTT reduction was dissolved in 100 µL of DMSO, and absorbance was read at 570 nm on a BIORAD 550 spectrophotometer. The percentages of cell survival related to the growth control wells (wells containing only cells and medium) were calculated, and the LC₅₀ (concentration that reduces the exposed sample survival rate to 50%) was determined. Doxorubi-

cin hydrochloride was used as a positive control substance; for this purpose, four serial dilutions were prepared at a maximum concentration of 10 µM (5.8 µg/mL).

Statistical analysis

Data were expressed as the mean values ± the standard error (SE). GraphPad-Prism® 4.0 was used to compare LC₅₀ values by means of the ANOVA analysis, followed by the Tukey Test, with a 95% of confidence.

Computational methods

All computations were carried out with the GAMESS program package (version 2006, Gordon Research Group, Iowa State University, USA).

RESULTS AND DISCUSSION

A preliminary screening of various doses of crude methanol extract of *C. stipuliformis* indicated its activity against five human cancer cell lines. Following the partitioning of the MeOH extract, the cytotoxic constituents were identified in the organic fraction (CSO) with LC₅₀ values below or near to 100 mg/mL (15); while the aqueous extract exhibited no significant inhibition of growth in the population of the tested cancer cells.

The GC-MS analysis of the *C. stipuliformis* bark hexane fraction revealed that CSH contained diterpenoid compounds (75.0%), sterols (16.0%), aliphatic esters (5.4%), sesquiterpenes (1.8%), aliphatic alcohols (1.6%), and other types of compounds (0.2%). Further fractionation of hexane fraction was needed due to its complexity. It afforded four fractions (i.e. CSHa-d), which were also analyzed through GC-MS, as it can be seen in figure 1. This analysis allowed determining that the composition between fractions was different as a result of an efficient separation; thus, the first CSHa fraction contained mono- and sesquiterpenoids; CSHb and CSHc showed the presence of diterpenoids; and CSHd consisted mainly of triterpenes and sterols.




Figure 1. Total ion chromatogram (GC-MS) of a) CSHa, b) CSHb, c) CSHc, d) CSHd fractions from *C. stipuliformis* bark obtained using a RTX-5 column (CSHa chromatogram was obtained on a DB-Wax column). Peak numbers correspond to the compound numbers in table 1.

Table 1. Chemical constituents from CSHa, CSHb, CSHc, and CSHd fractions determined through GC-MS.

No.	Compound	RI		Relative composition (%) ^a	No.	Compound	RI		Relative composition (%) ^a
		(DB-Wax)	(RTX-5)				(DB-Wax)	(RTX-5)	
CSHa					CSHa				
a1	α -Pinene	1007	961	1.19	a36	Cadalene	2203	-	0.14
a2	β -Ocimene	1102	-	0.23	a37	Methyl hexadecanoate	2212	1930	8.86
a3	Butanol	1145	-	18.39	a38	β -Eudesmol	2218	-	0.64
a4	43,55,67,82,108,139,169,199 ^b	1203	1799	0.14	a39	Methyl hydrojasmonate	2221	-	0.60
a5	α -Copaene	1484	1384	2.25	a40	(<i>Z,E</i>)-Farnesol	2270	-	0.38
a6	β -Cariophyllene	1589	1428	10.48	a41	Farnesilaldehyde	2278	-	0.48
a7	β -Ciclocitral	1594	-	0.58	a42	41,56,136,93,77,107,67,153 ^b	2282	-	0.52
a8	Allo-aromadendrene	1634	-	0.15	a43	Methyl heptadecanoate	2308	-	0.34
a9	α -Terpineol	1646	-	0.39	a44	Octadecanaldehyde	2318	-	0.43
a10	α -Humulene	1657	-	0.94	a45	(<i>E,E</i>)-Farnesol	2326	-	0.33
a11	<i>cis</i> -Carveol	1672	-	0.50	a46	Isoamyl anisate	2336	-	0.38
a12	γ -Muurolene	1680	1486	3.81	a47	41,79,55,69,91,105,119,216 ^b	2349	2160	0.52
a13	Ionene	1690	-	0.34	a48	β -Sesquiphellandrene	2355	-	0.69
a14	Germacrene D	1693	1486	1.72	a49	Methyl octadecanoate	2413	2074	3.82
a15	9-Aristolene	1697	-	0.61	a50	41, 55, 69, 74, 85, 97, 111, 264 ^b	2434	-	2.79
a16	β -Sclinenene	1707	-	1.18	a51	Isoamyl anthranilate	2442	-	0.75
a17	α -Amorfene	1716	1525	0.94	a52	Heneicosane	2493	2090	7.56
a18	Aristolene	1717	-	0.59	a53	41,79,55,68,96,108,292,149 ^b	>2500	2111	5.49
a19	δ -Cadinene	1752	1533	7.72	a54	Compound 2	>2500	2301	0.56
a20	β -Sesquiphellandrene	1761	-	0.29	a55	Compound 1	>2500	2247	6.73
a21	α -Cubebene	1773	-	0.16	a56	α -Ionol	>2500	-	0.08
a22	α -Cadinene	1784	-	0.26	CSHb				
a23	Fonolenol	1829	-	0.25	a55	Compound 1	-	2247	71.31
a24	Cuparene	1841	-	0.15	CSHc				
a25	α -Calarene	1882	-	0.14	a55	Compound 1	-	2247	50.30
a26	β -Calarene	1932	-	0.21	a54	Compound 2	-	2310	15.32
a27	Cariophyllene oxide	1968	1599	0.26	c1	Corticosterone	-	2416	5.24
a28	Methyl tetradecanoate	1996	-	0.28	c2	Compound 3	-	2422	13.01
a29	<i>E</i> -Nerolidol	2018	-	0.37	CSHd				
a30	Elemol	2079	-	0.75	d1	Ergosta-5-en-3-ol	-	2855	11.24
a31	Zingiberenol	2112	-	0.70	d2	55,83,69,159,255,271,300,412 ^b	-	2871	12.16
a32	Acorenone	2146	-	0.31	d3	β -Amirene	-	2890	17.70
a33	δ -Cadinol	2159	-	0.28	d4	β -Sitosterol	-	2901	34.94
a34	ϵ -Muurolene	2167	-	0.40	d5	Ergosta-4-en-3-one	-	2960	5.17
a35	T-Muurolol	2176	-	0.54					

^a Composition calculated based on the peak area % of each fraction.^b The eight most intense peaks of MS.

- Not detected.

The careful analysis of mass spectra and retention indexes of CSHa constituents allowed the identification of 51 compounds in this fraction, which identity is presented in table 1. The major constituents were identified as follows: the sesquiterpenes β -cariophyllene (10.48%), δ -cadinene (7.72%), γ -muurolene (3.81%), and α -copaene (2.25%); the diterpene **1** (6.73%); aliphatic esters methyl hexadecanoate (8.86%) and methyl octadecanoate (3.82%); butanol (18.39%), and heneicosane (7.56%). This high terpenoid content is comparable with those found for other *Croton* species, such as *C. zambesicus* (16) and *C. sellowii* (17).

The CSHd fraction was mainly constituted by sterols, among which β -sitosterol (34.94%), β -amirene (17.70%), ergosta-5-en-3-ol (11.24%), and ergosta-4-en-3-one (5.17%) were the major constituents. It is important to point out that most of the identified sterols have Δ^5 -nuclei, which is common in the *Croton* species (4).

The strongest cytotoxic activities were detected for CSHb and CSHc, which were subsequently fractionated by means of preparative HPLC in order to obtain pure compounds **1-3**. The major constituent of the CSHb fraction (**1**) has a formula of $C_{21}H_{32}O_2$, as it was established through the HRFAB-

MS analyses. The ^{13}C NMR spectra confirmed that compound **1** possesses a carboxyl group (δ_{C} 174.4). Also, the ^1H NMR spectra suggested that compound **1** possesses a $-\text{CH}=\text{C}(\text{CH}_3)-\text{CH}=\text{CH}_2$ group, which is typical for a labda-12(*E*),14-diene skeleton (18, 19), along with the presence of a cyclohexane ring, which was determined by of COSY and HMBC correlations. An exomethylene group was shown by the ^1H and ^{13}C NMR signals at δ_{H} 4.48 (H-17a, *brs*) and 4.86 (H-17b, *d*, $J = 0.8$), and δ_{C} 108.7 (C-17). The ^1H NMR spectrum also showed two additional singlet due to the presence of two methyl groups at δ_{H} 0.75 (H₃-20) and 1.72 (H₃-19); the latter was attached to an olefinic

group, which was attached to the cyclohexane ring. Therefore, the carboxyl group must be confined to an aliphatic chain attached to the cyclohexane system, which could be probed by the correlations between the carboxylic carbon at δ_{C} 174.4 (C-3) and methylene protons at δ_{H} 2.19 (H-2a) and 2.44 (H-2b) observed in the HMBC spectrum. These correlations revealed that the structure of compound **1** resembles a 3,4-*seco*-labda-12(*E*),14-diene skeleton due to the ring A opening; thus, it was elucidated as the methyl ester of (+)-12*E*-3,4-*seco*-labda-4(18),8(17),12,14-tetraen-3-oic acid (maravuic acid methyl ester). The chemical structure of this compound is presented in figure 2.




Figure 2. Labdane and *seco*-labdane diterpenes isolated from the *Croton stipuliformis* bark.

As it can be seen in figure 3, the relative stereochemistry of the substituents at the cyclohexane ring was proposed based on the analysis of the NOESY spectrum. The ^1H NMR spectrum of compound **1** clearly showed an axial configuration (δ 2.24, dd, $W_{1/2} = 21$ Hz) for H-5. Irradiation of methyl protons at δ_{H} 0.75 (H₃-20) enhanced the methylene proton signal at δ_{H} 1.72 (H-6_{ax} and H-11a), 2.25 (H-11_b) and 4.68 (H-18_a). The irradiation of H-9 (δ_{H} 1.88) enhanced the proton signals of H-5 (δ_{H} 2.24) and H-7_{ax} (δ_{H} 1.99), thus revealing a *trans*-1,3-diaxial configuration of H-5, H-7_{ax}, and H-9 protons. Further NOESY correlation between H-5 and H₃-19 (δ_{H} 1.72) was also found. Thus, it was possible to establish that the cyclohexane ring has a chair conformation with the

isopropenyl group at C-5 and the side-chain at C-9 in a β -equatorial orientation. (*E*)-Geometry of the C-12 double bond was evident from the upfield chemical shift of the methyl group at C-13 (δ_{H} 1.76, δ_{C} 12.1), and it was confirmed by the correlations between H-12 (δ_{H} 5.34) and H-14 (δ_{H} 6.31), as well as, the correlations between H₃-16 (δ_{H} 1.76) and H-11b (δ_{H} 2.25) in the NOESY spectrum. Finally, the correlation between the olefinic proton H-17b (δ_{H} 4.86) and H-7b (δ_{H} 2.34) suggests an equatorial orientation for this proton. The compound **1** structure optimization, which was done using the GAMESS computer program (20), is presented in figure 3. It is notable that compound **1** can be considered as an artifact of the MeOH reflux because the acid was also isolated.




Figure 3. Key NOESY correlations of compound **1**.

¹H and ¹³C NMR spectral data for compound **2** resembles those of compound **1**, except for the presence of a methoxy group (δ_{H} 3.64, δ_{C} 51.8). Therefore, compound **2** was identified as the 12*E*-3,4-*seco*-labda-4(18),8(17),12,14-tetraen-3-oic acid or Maravuic acid, a *seco*-labdane diterpene previously isolated from *Croton stipuliformis* leaves (21) and *Croton matourensis* bark (22).

The molecular formula of compound **3** was deduce as C₂₀H₃₀O₂ based on the HRFAB-MS and the EIMS [M⁺] (*m/z* 302). The unequivocal assignment of compound **3** was established through

the analysis of data from the HMQC, HMBC, and COSY experiments. The spectral data of the decalin ring system with 8(17) exomethylene of this compound was in agreement with those of the 12,15-epoxy-8(17),13-labdadienoic acid isolated from *Chunninghamia konishii* (23). Thus, compound **3** was identified as 8(17),12*E*,14-labdatrien-18-oic acid (communic acid), which has been previously reported in *Xylopija langsdorffiana* (24), *Isodon lophanthoides* (25) and *Pinus resinosa* (26), among others. The cardiovascular effects of compound **3** as hypotensor have been reported before (27); however, this is the first study about its cytotoxic activity.

The cytotoxicity of *Croton stipuliformis* bark methanol extract against human tumor cell lines were found in the organic fraction (CSO), as well as in the hexane fraction (CSH); as it was expected, crude methanol extract showed lower inhibitory effects compared to the subsequent fractions. The pure compounds (**1-3**) were evaluated for cytotoxicity against human larynx adenocarcinoma (HEp-2), human colon adenocarcinoma (HT-29), human gastric carcinoma (MKN-45), human breast carcinoma (MCF-7), and human cervix carcinoma (HeLa); and doxorubicin was used as positive control. The results for the afore mentioned evaluations are presented in table 2. All three diterpenes showed non-specific moderate cytotoxicities against those cell human tumor lines.

Table 2. Cytotoxicity data for methanolic extract, fractions, and compounds **1-3**^a.

Treatment	Cell lines ^b				
	HEp-2	HT-29	MKN-45	MCF-7	HeLa
Methanol extract	98.9 ± 10.0	108.8 ± 14.14	103.5 ± 1.3	90.3 ± 4.9	105.4 ± 1.8
CSO fraction	36.3 ± 16.9	56.5 ± 4.5	19.7 ± 1.4	40.3 ± 8.7	58.0 ± 3.5
CSH fraction	22.5 ± 3.4	17.2 ± 2.4	14.9 ± 3.0	16.6 ± 0.6	34.3 ± 1.4
Compound 1	47.2 ± 13.4	24.2 ± 4.9	22.6 ± 5.5	24.8 ± 4.9	19.1 ± 4.0
Compound 2	15.4 ± 0.4	16.6 ± 0.6	14.4 ± 1.2	16.0 ± 2.3	12.1 ± 0.9
Compound 3	25.0 ± 2.2	26.5 ± 1.4	-	-	26.5 ± 1.2
Doxorubicin HCl	0.08 ± 0.04	0.51 ± 0.18	0.17 ± 0.10	0.08 ± 0.02	0.95 ± 0.46

^a Results are expressed as the calculated lethal concentration 50 LC₅₀ (μg/mL). Data represent mean values ± SD (*n* = 3) from three independent studies.

^b HEp-2, human larynx adenocarcinoma; HT-29, human colon adenocarcinoma; MKN-45, human gastric carcinoma; MCF-7, human breast carcinoma; HeLa, human cervix carcinoma.

- Not determined.

Compound **1** showed a weak activity against human larynx adenocarcinoma (47.2 µg/mL) in comparison to the other human tumour lines tested, which can be interpreted as a slight selectivity towards the growth of cancer cell lines MCF-7, HeLa, HT-29 and MKN-45. Analogue compound **2** showed higher cytotoxic activity values than those of compounds **1** and **3**. It is very interesting to see that compounds **2** and **3** were more active than compound **1**; therefore, it is possible to establish that the presence of the methyl ester group diminished the cytotoxicity of compound **1**. This behaviour was also found for the benzyl ester of compound **2**, which was synthesized as part of this work (data not shown), and allowed us to confirm the significance of acid moiety in cytotoxicity.

It is important to point out that the cytotoxic activity of the three diterpenes isolated from *Croton stipuliformis* is higher than other known diterpenes, usually found in the *Croton* genus, such as crotonin and dehydrocrotonin (28-30).

CONCLUSIONS

The hexane fraction of *Croton stipuliformis* presented a moderate cytotoxic activity against different *in vitro* human cancer cell lines. Moreover, the cytotoxicity of diterpenes **1-3** isolated from the afore mentioned species was demonstrated for the first time in this study.

ACKNOWLEDGMENTS

Financial support from DINAIN-Universidad Nacional de Colombia is gratefully acknowledged. The authors are grateful to Dr. Ricardo Acuña from CENICAFE (Colombia) for providing us with the sample for this study. The authors also would like to thank Prof. Dr. Francisco José Heredia from Universidad de Sevilla for his support on the recording of HRMS experiments.

REFERENCES

- Murillo JA. Las Euphorbiaceae de Colombia. Biota Colombiana. 2004 Dec; 5 (2): 183-200.
- Murillo JA. Composición y distribución del género *Croton* (Euphorbiaceae) en Colombia, con cuatro especies nuevas. Caldasia. 1999 Sep; 21 (2): 141-166.
- Jones K. Review of Sangre de Drago (*Croton lechleri*)-a south american tree sap in the treatment of diarrhea, inflammation, insect bites, viral infections, and wounds: traditional uses to clinical research. J Altern Complem Med. 2003 Dec; 9 (6): 877-896.
- Salatino A, Faria Salatino ML, Negri G. Traditional uses, chemistry and pharmacology of *Croton* species (Euphorbiaceae). J Braz Chem Soc. 2007 Jan-Feb; 18 (1): 11-33.
- Sagawa T, Takaishi Y, Fujimoto Y, Duque C, Osorio C, Ramos F, et al. Cyclobutane dimers from the Colombian medicinal plant *Achyrocline bogotensis*. J Nat Prod. 2005 Mar; 68 (4): 502-505.
- Ramos F, Takaishi Y, Kawazoe K, Osorio C, Duque C, Acuña R, et al. Immunosuppressive diacetylenes, ceramides and cerebrosides from *Hydrocotyle leucocephala*. Phytochemistry. 2006 Jun; 67 (11): 1143-1150.
- Cordero CP, Morantes SJ, Páez A, Rincón J, Aristizábal FA. Cytotoxicity of withanolides isolated from *Acinetus arborescens*. Fitoterapia. 2009 Sep; 80 (6): 364-368.
- Smith ECJ, Williamson EM, Wareham N, Kaatz GW, Gibbons S. Antibacterials and modulators of bacterial resistance from the immature cones of *Chamaecyparis lawsoniana*. Phytochemistry. 2007 Jan; 68 (2): 210-217.
- Kondjoyan N, Berdagué JL. A compilation of relative retention indices for the analysis of aromatic compounds. Theix, France: Laboratoire Flavcur; 1996.
- McLafferty FW, Stauffer D B. The Wiley/NBS Registry of Mass Spectral Data, 5th ed. New York, United States: Wiley; 1991.
- Joulain D, König WA. The Atlas of Spectral Data of Sesquiterpene Hydrocarbons. Hamburg, Germany: Verlag; 1998.
- Adams RP. Identification of essential oil components by gas chromatography/Quadrupole Mass Spectroscopy. Carol Stream, IL, United States: Allured Publishing Corporation, 2001.
- Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assay. J Immunol Methods. 1983 Dec; 65 (1-2): 55-63.
- Cordero C, Aristizábal F. Evaluación preliminar *in vitro* de citotoxicidad de extractos vegetales, empleando métodos colorimétricos. Rev Col Biotecon. 2002 Jan-Feb; 4 (1): 100-106.
- Teicher, B., Totowa, N. J. Eds. Anticancer drug development guide: preclinical screening, clinical trials and approval. New York, USA: Humana Press, 1997. Boyd M. The NCI *in vitro* anticancer drug discovery screen: concept, implementation and operation 1985-1995; p. 23-42.
- Fekam Boyom F, Keumedjio F, Jazet Dongmo PM, Ngadjui BT, Amvam Zollo PH, Menut C, et al. Essential oils from *Croton zambesicus* Muell. Arg. growing in Cameroon. Flavour Frag J. 2002 May-Jun; 17 (3): 215-217.
- Conserva EM, Palmeira JSF, Moura F, Alves V, Oliveira FM, Bento ES, et al. Neutral components from hexane extracts of *Croton sellowii*. Flavour Frag J. 2004 Jan-Feb; 19 (1): 69-71.
- Roengsumran S, Petsom A, Sommit D, Vilaivan T. Labdane diterpenoids from *Croton oblongifolius*. Phytochemistry. 1999 Feb; 50 (3): 449-453.
- Noma M, Suzuki F, Gamou K, Kawashima N. Two labdane diterpenoids from *Nicotiana raimondii*. Phytochemistry. 1982 Feb; 21 (2): 395-397.
- Schmidt MW, Baldrige KK, Boatz JA, Elbert ST, Gordon MS, Jensen JH, et al. General atomic and molecular electronic structure system. J Comput Chem. 1993 Nov; 14 (11): 1347-1363.
- Ramos FA, Takaishi Y, Kashiwada Y, Osorio C, Duque C, Acuña R, et al. Ent-3,4-seco-labdane and ent-labdane diterpenoids from *Croton stipuliformis* (Euphorbiaceae). Phytochemistry. 2008 Sep; 69 (12): 2406-2410.
- Schneider C, Breitmaier E, Bayma J, de Franca LF, Kneifel H, Krebs HC. Maravuiic acid, a new *seco*-labdane diterpene from *Croton matourensis*. Liebigs Ann. 1995 Mar; (4): 709-710.
- Li YC, Kuo YC. Labdane-type diterpenoids from the wood of *Chunninghamia konishii*. Chem Pharm Bull. 2002 Apr; 50 (4): 498-500.
- Ribeiro LA, Tavares JF, De Andrade N, Da Silva M, Da Silva B. The (8)17,12E,14-labdatrien-18-oic acid (labdane 302), a labdane-type diterpene isolated from *Xylopia langsdorffiana* St.

- Hil. & Tul. (Annonaceae), relaxes the guinea pig trachea. *Rev Bras Farmacogn.* 2007 Jun; 17 (2): 197-203.
25. Jiang B, Lu ZQ, Zhang HJ, Zhao QS, Sun HD. Diterpenoids from *Isodon lophanthoides*. *Fitoterapia.* 2000 Aug; 71 (4): 360-364.
 26. Zinkel DF, Clarke WB. Resin acids of *Pinus resinosa* needles. *Phytochemistry.* 1985 Jun; 24 (6): 1267-1271.
 27. Pereira de Oliveira A, Furado F, Sobral da Silva M, Tavares J, Mafra RA, Araujo DA, *et al.* Calcium channel blockade as a target for the cardiovascular effects induced by the 8(17),12E,14-labdatrien-18-*oic* acid (labdane-302). *Vasc Pharmacol.* 2006 May; 44 (5): 338-344.
 28. Rodríguez JA, Haun M. Cytotoxicity of *trans*-dehydrocrotonin from *Croton cajucara* on V79 cells and rat hepatocytes. *Planta Med.* 1999 Aug; 65 (6): 522-526.
 29. Haun M, Conte Anazetti M, Silva Melo P, Durán N. Comparative cytotoxicity of dimethylamide-crotonin in the promyelocytic leukemia cell line (HL60) and human peripheral blood mononuclear cell. *Toxicology.* 2003 Jun 30; 188 (2-3): 261-274.
 30. Grynberg NF, Echevarria A, Lima JE, Pamplona SSR, Pinto AC, Maciel MAM. Anti-tumour activity of two 19-nor-clerodane diterpenes, *trans*-dehydrocrotonin and *trans*-crotonin, from *Croton cajucara*. *Planta Med.* 1999 Dec; 65 (8): 687-689.

CIDUA

Centro de Documentación e Información de Medicamentos,
Alimentos, Cosméticos y Productos Naturales
Teléfono: 2195455

Correo electrónico: cidua@farmacia.udea.edu.co; cidua1@gmail.com

ACTIVIDADES

VIERNES DEL CIDUA: “Un espacio de los estudiantes, para los estudiantes”: un grupo humano se reúne para establecer lazos de comunicación entorno a un conocimiento y a la vez se adquiere formación sobre un tema específico.

CONSULTA TELEFÓNICA: Servicio dirigido a la comunidad en general, para orientar e informar acerca del uso y consumo de medicamentos o dudas referentes a las temáticas en las cuales se especializa el Centro. El usuario plantea a los auxiliares la inquietud; esta solicitud se remite a profesionales especializados en dichas temáticas quienes dan respuesta de manera ágil y oportuna a la inquietud planteada por el usuario entre tres o cinco días hábiles. El servicio se ofrece de lunes a viernes durante el horario de atención (8:00 am. a 6:00 pm.)

CONSULTA EN SALA: Acceso y consulta a los diferentes materiales bibliográficos del Centro. Este servicio es utilizado en su mayoría por los estudiantes de las áreas de formación de la Facultad de Química Farmacéutica quienes tienen la posibilidad de disponer del espacio para consultar sus propios materiales y para solicitar otras referencias bibliográficas que pueden ampliar sus campos de estudio.

SERVICIO ORIENTACION AL USUARIO: Asesoría personalizada que se brinda al usuario en la búsqueda y selección de la información de su interés y en la utilización de los recursos y servicios que ofrece el Centro de Documentación. El fondo documental del CIDUA es cerrado, por tanto existen unas estrategias de búsqueda (Opac) y auxiliares que le colaborarán para ayudar a la recuperación de la información y así prestar un servicio oportuno y satisfacer las necesidades informáticas del usuario.

CIRCULACIÓN Y PRÉSTAMO: El usuario podrá retirar en calidad de préstamo y por un tiempo determinado material bibliográfico disponible en el Centro de Documentación. Este servicio se presta especialmente a la comunidad académica de la Universidad de Antioquia y para los usuarios de instituciones con las cuales el Sistema de Bibliotecas tenga convenios de préstamo interbibliotecario.

PRÉSTAMO INTERBLIOTECARIO O INTERUNIVERSITARIO: Programa de cooperación bibliotecaria, a través de convenios para acceder a documentos de otros fondos documentales que la unidad que tramita la petición no posee. Este servicio atiende las peticiones de sus usuarios internos a quien les da una carta de autorización para que tramiten ante la biblioteca que tiene el material de su interés (temáticas comunes).

SERVICIO DE ALERTA: Permite informar al usuario acerca de las novedades bibliográficas adquiridas por el centro de Información y Documentación, a través de un boletín electrónico que estará montado en la página del CIDUA para consulta de toda la comunidad académica y se enviara oportunamente al correo electrónico de los docentes de la Facultad de Química Farmacéutica.

BIBLIOGRAFIA ESPECIALIZADA: Consiste en realizar una búsqueda bibliográfica de un tema específico de acuerdo al interés del usuario. Esta búsqueda se realiza en los diferentes materiales con los cuales cuenta el CIDUA y se presta a cualquier usuario que solicite el servicio. La bibliografía realizada es enviada al correo electrónico o puede entregarse al usuario personalmente.