

# CONJUGATED LINOLEIC ACID, FATTY ACID PROFILE AND PROCESS PROPERTIES IN KUMIS - FERMENTED MILK CONSUMED IN COLOMBIA

ACIDO LINOLEICO CONJUGADO, PERFIL DE ÁCIDOS GRASOS Y PROPIEDADES  
DEL PROCESO EN KUMIS - BEBIDA FERMENTADA CONSUMIDA EN COLOMBIA

Julián A. OSORIO<sup>1,2</sup>, Carolina RAMÍREZ<sup>2</sup>, Carlos F. NOVOA<sup>2</sup>, Luis-Felipe GUTIÉRREZ<sup>2\*</sup>

Received: 01 December 2010 Accepted: 14 June 2011

## ABSTRACT

In this study, we reported the concentration of conjugated linoleic acid of the main commercial kumis consumed and distributed in Colombia, as well as the concentration of conjugated linoleic acid of an artisanal kumis elaborated with two different types of milk (skim liquid and powder reconstituted). Conjugated linoleic acid (C18:2c9t11) contents, expressed as mg of rumenic acid/g fat, ranged from  $7.63 \pm 0.96$  to  $22.62 \pm 3.85$ . Also, the main fatty acids of kumis samples were identified and quantified. pH value ranged between  $3.84 \pm 0.02$  and  $4.28 \pm 0.01$ , and titratable acidity ranged between  $0.69 \pm 0.01$  and  $0.94 \pm 0.02\%$  of lactic acid. Consistence and flux indices presented values between  $2.01 \pm 0.05$  and  $7.08 \pm 0.39$  (Pa·s<sup>n</sup>) and from 0.43 to 0.26, respectively. These results indicate that kumis is a food product that could be used for supplying important amounts of conjugated linoleic acid in the human diet.

**Keywords:** Conjugated linoleic acid, fermented milk, kumis, fatty acids, rheological properties.

## RESUMEN

En este estudio reportamos las concentraciones de ácido linoleico conjugado en algunos de los kumis comerciales de mayor distribución y consumo en Colombia, así como de kumis artesanal elaborado con dos tipos diferentes de leche (líquida semidescremada y leche en polvo reconstituida). Los contenidos de ácido linoleico conjugado (C18:2c9t11), expresadas en mg de ácido ruménico/g grasa, variaron de  $7,63 \pm 0,96$  a  $22,62 \pm 3,85$ . Los principales ácidos grasos de las muestras de kumis fueron también identificados y cuantificados. Los valores de pH y la acidez titulable variaron entre  $3,84 \pm 0,02$  y  $4,28 \pm 0,01$ , y entre  $0,69 \pm 0,01$  y  $0,94 \pm 0,02\%$  ácido láctico, respectivamente. Los índices de consistencia y de flujo presentaron valores entre  $2,01 \pm 0,05$  y  $7,08 \pm 0,39$  (Pa·s<sup>n</sup>), y de 0,43 a 0,26, respectivamente. Estos resultados indican que el kumis es un alimento que podría ser utilizado para aportar cantidades importantes de ácido linoleico conjugado en la dieta humana.

**Palabras clave:** ácido linoleico conjugado, leche fermentada, kumis, ácidos grasos, propiedades reológicas.

<sup>1</sup> Departamento de Ingeniería Agrícola y de Alimentos. Facultad de Ciencias Agropecuarias. Universidad Nacional de Colombia, sede Medellín. A.A. 568. Medellín, Colombia.

<sup>2</sup> Instituto de Ciencia y Tecnología de Alimentos (ICTA). Grupo de investigación en aseguramiento de la calidad de alimentos y desarrollo de nuevos productos. Universidad Nacional de Colombia, sede Bogotá, Carrera 30 No. 45-03, Edificio 500A.

\* Corresponding author: lfgutierrez@unal.edu.co

## INTRODUCTION

Conjugated linoleic acid (CLA) is a generic term used to describe the isomers of linoleic acid, which is an essential fatty acid found in appreciable amounts in food products derived from ruminant animals, especially in the lipid fraction of milk and meat (1). Most dairy products contain different CLA isomers, of which 85% to 95% consist of rumenic acid (*cis*-9, *trans*-11, octadecadienoic acid, C18:2*c9t11*) in amounts ranging from 6 to 16 mg/g of total fat (2). The metabolic pathway proposed for the formation of CLA isomers includes the isomerization and biohydrogenation of unsaturated fatty acids by rumen bacteria (*Butyrivibrio fibrisolvens*), and the desaturation of vaccenic acid (C18:1*t11*) by  $\Delta^9$ -desaturase in the mammary gland (3). According to Ledoux *et al.*, 2005 (4), the incomplete biohydrogenation of linoleic acid in the rumen provides the surplus of CLA found in milk fat.

Biologically active CLA isomers (*cis*-9, *trans*-11, octadecadienoic acid: C18:2*c9t11* and *cis*-10, *trans*-12, and octadecadienoic acid: C18:2*c10t12*) have shown to have beneficial effects on human health, and can be considered as therapeutic nutrients with protective effects against various common diseases such as obesity, atherosclerosis, chronic inflammatory diseases and cancer (5-8). According to the published reports, the recommended CLA intake in order to achieve beneficial effects on health varies between 0.7 and 6.8 g CLA/day (9-11).

In the '90s decade, it was hypothesized that CLA concentration could increase during milk fermentation (12-13). Recent literature suggests that the CLA content in yogurts could be increased by using certain lactic acid bacteria. Different strains of *Lactobacillus acidophilus*, *Lactobacillus rhamnosus*, *Lactobacillus plantarum*, *Bifidobacterium bifidum*, *Pro-*

*pionibacterium freudenreichii* ssp. *shermanii*, *Lactococcus lactis*, and *Lactococcus casei* have been reported to enhance the CLA concentration during milk fermentation (14-18). According to Shantha *et al.*, 1995 (13), typical values of CLA in fermented milks range from 3.41 to 9.12mg/g of fat.


Kumis, a traditional fermented milk product widely consumed in Colombia, is produced at industrial and artisanal scales by fermenting whole or semi-skimmed milk with mesophilic cultures (*Lactococcus lactis* subsp. *cremoris* and *Lactococcus lactis* subsp. *lactis*) during 20 - 24 hours, at temperatures ranging from 26 to 28°C, until reaching a pH value varying between 4.00 and 4.25. The available data of the CLA concentration in milk and dairy products in Colombia are quite scarce; only two studies have been reported for milk products from the Sabana de Bogotá region (19-20). To the best of our knowledge, data of CLA content in kumis have not been still reported in Colombia. In this work, we report for the first time the CLA concentration and the fatty acid profiles of the most widely distributed and consumed kumis product brands in Colombia, in order to set a basis for further research focused on developing a CLA-enriched kumis. Some physicochemical properties such as the pH value, titratable acidity, and rheological parameters are also reported.

## MATERIALS AND METHODS

The main characteristics of the kumis samples analyzed in this work are presented in table 1. Three samples of each brand from the same batch and date of production were obtained from local markets, and once arrived at the laboratory they were codified and stored at 6 - 8°C. Figure 1 shows the process followed for preparing samples of artisanal kumis.

**Table 1.** Main characteristics of the analyzed kumis samples.

Sample code	Type	Characteristics
A	Commercial	Hygienized milk, sucrose, and lactic cultures
B	Commercial	Pasteurized milk, sucrose, powder milk and lactic cultures
C	Commercial	Fluid milk, sucrose, powder milk, lactic cultures ( <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and ssp. <i>lactis</i> , <i>Streptococcus thermophilus</i> )
D	Commercial	Partial skim milk, sucrose, lactic protein, lactic cultures ( <i>Lc. lactis</i> , <i>Lc cremoris</i> , <i>L. casei</i> )
E	Commercial	Hygienized milk, sucrose and lactic cultures
F	Commercial	Hygienized milk, sucrose and lactic cultures
G	Commercial	Hygienized milk, sucrose, lactic cultures ( <i>Streptococcus lactis</i> or <i>cremoris</i> )
I	Commercial	Hygienized milk, sucrose and lactic cultures.
EC1	Artisanal	Partial skim milk, sucrose, <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and spp. <i>lactis</i> (Choozit™ MA11 LYO 25 DCU®. Danisco)
EC2	Artisanal	Partial skim milk, sucrose, <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and spp. <i>lactis</i> (Chr Hansen R-708®)
CC1	Artisanal	Water, powder milk, sucrose <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and spp. <i>lactis</i> (Choozit™ MA11 LYO 25 DCU®. Danisco)
CC2	Artisanal	Water, powder milk, sucrose, <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and spp. <i>lactis</i> (Chr Hansen R-708®)
IC1	Artisanal	Water, powder milk, <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and spp. <i>lactis</i> (Choozit™ MA11 LYO 25 DCU®. Danisco)
IC2	Artisanal	Water, powder milk, sucrose, <i>Lactococcus lactis</i> ssp. <i>cremoris</i> and spp. <i>lactis</i> (Chr Hansen R-708®)



**Figure 1.** Procedure for the preparation of artisanal kumis samples.

### Extraction and quantification of the fat content

The fat from the kumis samples was extracted following the methodology described by Folch *et al.*, 1957 (21) with some modifications. In a typical extraction, samples (~6 g) were mixed with chloroform-methanol 2:1, and then centrifuged at 5000 rpm during 10 min. The mixture was transferred to a separatory funnel and then, 7 mL KCl (0.88% w/v) were added. After a vigorous agitation, the organic phase was separated, dried with anhydrous sodium sulfate, and after removing the solvent by vacuum evaporation, the obtained fat samples were collected, evaporated under nitrogen, weighed, and stored in sealed amber glass vials at -20°C until the analysis was performed.

### Fatty acid analysis and quantification

Fat samples were converted into their methyl esters (FAME) by derivatization in alkaline media, using sodium methoxide 0.5 M as indicated by Christie *et al.*, 2007 (22). FAMES were analyzed through gas chromatography (GC), using an Agilent® 7890A gas chromatograph (Agilent, USA). The oven temperature was programmed as follows: from 60°C (isothermal for 1 min) to 190°C at 20°C/min, and an isothermal period of 12.5 min at 190°C,

for a total analysis time of 19 min. The injector and detector temperatures were set at 250°C. Helium was used as carrier gas at a flow rate of 2.0 mL/min. The separation of FAMES was performed on a BPX-70 capillary column (60 m×0.25 mm i.d.×0.25 μm film thickness; SGE, Melbourne, Australia). Fatty acids were identified by comparing their retention times with those of the FAME standards (C4-C20 and CLA FAMES), which were purchased from Sigma Aldrich (Sigma Aldrich, USA), under the same conditions. Peaks were integrated using Agilent ChemStation® software. The quantification of the identified fatty acids was carried out following the internal standard method with calibration plots for each analyzed fatty acid.

### Physicochemical characteristics

pH measurements of kumis samples were determined using a Orion® 420A+ pH-meter with automatic compensation of temperature a 25°C. Before the analysis, the pH-meter was calibrated using reference pH 4.0 and pH 7.0 buffer solutions, and the kumis samples were gently stirred. Titratable acidity was evaluated through titration with NaOH 0.1 M using phenolphthalein as indicator.

### Rheological measurements

Rheological measurements were carried out at constant temperature (10 ± 0.5°C) using a rotational rheometer (Haake ROTOVISCO RV 20) with concentric cylinders, and a SV II sensor system. Shear rates ranging from 1.17 to 117.12 s<sup>-1</sup> were applied. The shear rate (s<sup>-1</sup>) vs. shear stress (Pa) data were fitted to the following power law model:

$$\tau = K\dot{\gamma}^n \quad \text{Equation 1.}$$

where:

$\tau$ : Shear stress (Pa);  $K$ : Consistency index (Pa·s<sup>n</sup>);  $\dot{\gamma}$ : Shear rate (s<sup>-1</sup>);  $n$ : Flow index.

### Experimental design and statistical analysis

A complete randomized experimental design was used to analyse the effects of the product, on CLA concentration, fatty acids profile, consistence index ( $K$ ), flux index ( $n$ ), pH and titratable acidity. All assays were made in triplicate. The obtained data were analyzed using the following model (23):

$$Y_{ij} = \mu + \alpha_i + \varepsilon_{ij} \quad \text{Equation 2.}$$

where:

$Y_{ij}$ : Variable value;  $\mu$ : Variable general mean;  $\alpha$ : Effect of product (i=14);  $\epsilon_{ij}$ : Experimental error.

The analysis of variance was used to identify differences in fatty acid concentration, consistence index (K), flux index (n), pH, and titratable acidity. The Shapiro Wilk test was used to verify the normality of the experimental error. The Tukey test was employed for the mean separation and correlation. A statistical analysis was carried out using the statistical analysis system software (SAS®).

### Index of atherogenicity

The index of atherogenicity was calculated following the method described by Ulbricht and Southgate, 1991 (24), by means of the next equation:

$$IA = \frac{C12:0 + 4C14:0 + C16:0}{\sum \text{Unsaturated fatty acids}} \quad \text{Equation 3.}$$

### Estimation of daily CLA intake in Colombia

The average of CLA intake in Colombia was evaluated from the data published by the National Survey of Nutritional Situation in Colombia 2005 (25), which provides the food intake (g/day) for different food products. We estimated the fat (%w/w) and CLA contents (mg/g fat) of each product from their typical available data (26-27).

## RESULTS AND DISCUSSION

### CLA concentrations and fatty acid profiles

Table 2 presents the results of CLA concentrations in the kumis samples, which ranged from  $7.63 \pm 0.96$  to  $22.62 \pm 3.85$  mg/g fat, which are amounts equivalent to 1.13% and 2.81% of fatty acid methylesters (FAMES), respectively.

**Table 2.** CLA concentrations and index of atherogenicity (IA) in kumis samples<sup>a,b</sup>.


Sample	CLA (mg/g fat)	CLA (g FA/100 FAME)	Portion (g)	CLA intake (mg/portion)	IA
A	18.29 ± 1.44 <sup>adf</sup>	2.81 ± 0.20	200	77.55 ± 6.11	2.1
B	12.84 ± 1.39 <sup>cf</sup>	1.86 ± 0.20	150	47.96 ± 5.19	2.3
C	15.47 ± 4.92 <sup>aef</sup>	2.96 ± 0.70	150	43.16 ± 13.73	2.1
D	22.62 ± 3.85 <sup>af</sup>	2.90 ± 0.55	150	41.39 ± 7.04	2.2
E	13.13 ± 1.74 <sup>acf</sup>	1.69 ± 0.25	210	59.56 ± 7.89	2.5
F	7.63 ± 0.96 <sup>bc</sup>	1.13 ± 0.14	200	34.18 ± 4.30	2.5
G	10.59 ± 1.01 <sup>bce</sup>	1.37 ± 0.14	150	41.62 ± 3.97	2.3
I	9.17 ± 0.70 <sup>bc</sup>	1.31 ± 0.10	150	35.07 ± 2.68	2.6
EC1	20.83 ± 3.93 <sup>a</sup>	2.71 ± 0.56	230	66.11 ± 12.47	2.6
EC2	19.66 ± 1.41 <sup>af</sup>	2.68 ± 0.20	230	61.50 ± 4.41	2.7
CC1	16.63 ± 3.16 <sup>aef</sup>	2.58 ± 0.45	230	93.71 ± 17.81	2.4
CC2	11.77 ± 2.21 <sup>bd</sup>	1.31 ± 0.31	230	88.25 ± 16.57	2.9
IC1	10.56 ± 2.51 <sup>b</sup>	1.32 ± 0.36	230	85.01 ± 20.20	1.9
IC2	12.06 ± 1.98 <sup>bd</sup>	1.78 ± 0.28	230	121.77 ± 19.99	1.5

<sup>a</sup> Values in the same column followed by different letters are significantly different,  $p \leq 0.05$ .

<sup>b</sup> Values are means ± standard deviation of triplicate determinations.

As it can be seen in table 2, the CLA concentration values are slightly higher than those reported in literature for fermented dairy beverages, which range from 0.35 to 3.95 FAMES (%) (28-30). These results could be explained because pasture is the main component of the animal diet in Colombia, and various studies have concluded that the pasture feeding can enhance the CLA concentration in milk (3). Due to the fact that the production process of commercial kumis samples was unknown, it was not possible to identify the factors explaining the variation among their CLA concentrations. However, data for artisanal kumis indicate that kumis samples elaborated with liquid skim milk presented higher CLA concentration values than those produced with reconstituted whole powder milk. Differences between milk constituents, as well as the effects of the process for obtaining powder milk, could explain these results. No significant differences ( $p > 0.05$ ) were found between the CLA content of kumis elaborated with different commercial lactic cultures. Our results are in agreement with those reported by Boylston and Beitz, 2002 (28) for yogurt produced from milk of cows fed with soy oil and conjugated linoleic acid, and with the ones published by Rico *et al.*, 2007 (20) for milk fat from the Sabana de Bogotá region.

Figure 2 shows the concentrations of short chain saturated, medium chain (atherogenic), and long chain unsaturated and C18:0 fatty acids in kumis samples. As it can be observed in figure 2a, butyric acid (C4:0) was the predominant fatty acid in the group of short chain saturated fatty acids, varying between 18 and 69 mg/g of fat. Short chain fatty acids presented values varying between 9% and 15% of the total fatty acids. According to Parodi, 2004 (5), this group of fatty acids has no effects on blood cholesterol levels. Moreover, from this group of fatty acids, it is possible to highlight the antitumoral properties and the synergistic capacity in the treatment of hypercholesterolemia reported for butyric acid (29, 31-32). However, caproic (C6:0), caprylic (C8:0) and capric (C10:0) fatty acids have been reported to possess antibacterial and antiviral properties, presenting activity against HIV (33).



**Figure 2.** Concentration of (a) short chain saturated, (b) medium chain (atherogenic) and (c) long chain unsaturated and C18:0 fatty acids in kumis samples.

Concentrations of medium chain (atherogenic) fatty acids are presented in figure 2b. As it can be observed, palmitic acid (C16:0) was the most important fatty acid of this group, ranging from 98 to 280 mg/g of fat. These values are in agreement with those recently published by Collomb *et al.*, 2002 (34) and Simionato *et al.*, 2010 (35), who reported values for palmitic acid between 208 and 276 mg/g of fat. Myristic (C14:0) and lauric (C12:0) acids were found

in amounts ranging between 39 and 81 mg/g of fat, and from 14 to 22 mg/g of fat, respectively.

Saturated and unsaturated C18 fatty acids found in kumis samples are showed in figure 2c. Oleic (C18:1c) and stearic (C18:0) acids presented the highest values of this group. C18:1c ranged from 90 to 230 mg/g of fat, whereas C18:0 ranged between 64 and 163 mg/g of fat. Similar values have been reported in the literature for these fatty acids (34-35). According to Parthasarathy *et al.*, 1990 (36), the stearic acid has no effects on the increase of serum cholesterol. This compound represents approximately 12% of the fatty acids in milk fat, and like oleic acid, which ranges between 15 and 23%, it is effective for reducing plasma cholesterol (36). Moreover, contrary to the hypercholesterolemic effects of atherogenic fatty acids, several studies indicate that long-chain saturated fatty acids produce a slow decrease of arterial occlusions and of platelet aggregation. Also, the presence of

fatty acids such as linoleic and  $\alpha$ -linolenic, having recognized beneficial effects on cardiovascular health, contribute to the biological potentialities of the kumis fat (37).

### Physicochemical characteristics

Table 3 shows the physicochemical properties of the analyzed kumis samples. pH values and titratable acidity were in agreement with the reported data for fermented dairy beverages. pH values ranged between 3.84 and 4.28, lactic acid varied from 0.69% to 0.94%. The fat content of samples oscillated between 1.22 and 4.39% w/w, whereas the interval of values of the consistence and flow index were 2.0 and 7.0 Pa·s<sup>n</sup>, and 0.26 and 0.43, respectively. Due to the low formation of exopolysaccharides for the starter culture used in milk fermentation, the consistence index of kumis samples was lower than those reported for other fermented milks, such as the yogurt (38 - 40).

**Table 3.** Physicochemical characteristics of kumis samples<sup>a,b</sup>.

Sample	pH	Lactic acid (%)	Fat (%)	K (Pa·s <sup>n</sup> )	n
A	4.18 ± 0.02 <sup>b</sup>	0.69 ± 0.01 <sup>h</sup>	2.12 ± 0.12 <sup>def</sup>	3.08 ± 0.61 <sup>cd</sup>	0.36 ± 0.03
B	4.08 ± 0.02 <sup>de</sup>	0.74 ± 0.01 <sup>f</sup>	2.49 ± 0.15 <sup>cd</sup>	2.23 ± 0.54 <sup>c</sup>	0.40 ± 0.04
C	4.11 ± 0.01 <sup>cd</sup>	0.74 ± 0.01 <sup>f</sup>	1.86 ± 0.20 <sup>defg</sup>	2.48 ± 0.86 <sup>c</sup>	0.37 ± 0.07
D	4.02 ± 0.04 <sup>f</sup>	0.80 ± 0.01 <sup>d</sup>	1.22 ± 0.30 <sup>g</sup>	2.34 ± 0.99 <sup>c</sup>	0.38 ± 0.06
E	4.21 ± 0.01 <sup>b</sup>	0.72 ± 0.01 <sup>g</sup>	2.16 ± 0.25 <sup>de</sup>	3.21 ± 1.11 <sup>cd</sup>	0.37 ± 0.05
F	4.04 ± 0.02 <sup>cf</sup>	0.78 ± 0.01 <sup>e</sup>	2.24 ± 0.15 <sup>d</sup>	2.01 ± 0.05 <sup>c</sup>	0.42 ± 0.01
G	4.01 ± 0.00 <sup>f</sup>	0.75 ± 0.01 <sup>f</sup>	2.62 ± 0.06 <sup>cd</sup>	4.56 ± 0.35 <sup>cd</sup>	0.29 ± 0.01
H	3.84 ± 0.02 <sup>g</sup>	0.81 ± 0.00 <sup>cd</sup>	2.55 ± 0.26 <sup>cd</sup>	2.47 ± 0.18	0.39 ± 0.01
EC1	4.17 ± 0.02 <sup>b</sup>	0.81 ± 0.00 <sup>c</sup>	1.38 ± 0.06 <sup>efg</sup>	6.35 ± 0.41 <sup>ab</sup>	0.26 ± 0.01
EC2	4.20 ± 0.01 <sup>b</sup>	0.83 ± 0.01 <sup>b</sup>	1.36 ± 0.21 <sup>fg</sup>	7.08 ± 0.39 <sup>a</sup>	0.27 ± 0.01
CC1	4.19 ± 0.01 <sup>b</sup>	0.94 ± 0.01 <sup>a</sup>	2.45 ± 0.49 <sup>d</sup>	5.91 ± 0.36 <sup>abc</sup>	0.43 ± 0.24
CC2	4.16 ± 0.01 <sup>bc</sup>	0.94 ± 0.01 <sup>a</sup>	3.26 ± 0.14 <sup>bc</sup>	5.45 ± 0.33 <sup>abc</sup>	0.29 ± 0.01
IC1	4.28 ± 0.01 <sup>a</sup>	0.82 ± 0.01 <sup>bc</sup>	3.50 ± 0.46 <sup>b</sup>	4.42 ± 0.22 <sup>cd</sup>	0.32 ± 0.01
IC2	4.26 ± 0.01 <sup>a</sup>	0.81 ± 0.01 <sup>cd</sup>	4.39 ± 0.33 <sup>a</sup>	4.98 ± 0.67 <sup>bc</sup>	0.30 ± 0.02

<sup>a</sup> Values in the same column followed by different letters are significantly different,  $p \leq 0.05$ .

<sup>b</sup> Values are means ± standard deviation of triplicate determinations.

### Index of atherogenicity

The index of atherogenicity (IA) of kumis samples is presented in table 2. This index allows comparing food products and diets, and it is related to the polyunsaturated/saturated fatty acids ratio. This index indicates the contribution of dietary factors that promote or protect against the development of

coronary heart disease (CHD) subsequent to the onset of atherosclerosis. The equation proposed by Ulbricht and Southgate, 1991 (41) for calculating the atherogenic index, indicates that C12:0, C14:0, and C16:0 FA are atherogenic, and that n-3, n-6, and monounsaturated FA are antiatherogenic. Thus, the higher the IA, the more atherogenic dietary components are (41).

The IA of the kumis samples analyzed in this study ranged between 1.5 and 2.9. The Atherogenicity indices ranging between 1.35 and 2.80 were recently published by Huang *et al.*, 2008 (42) for milk obtained from cows with soy oil supplemented diets, conjugated linoleic acid, or both. The lowest IA were obtained when supplementing with soy oil and CLA, and the highest value corresponded to milk sample control. Gagliostro *et al.*, 2007 (30) reported IA of 2.32 in milk and yogurt with CLA concentrations of 1.04% and 1.09%, respectively, whereas for CLA-enriched milk products and yogurt, the IA ranged between 0.74 and 0.80 for CLA concentrations of 4.14% and 3.95%, respectively. Our results are in accordance with the available data, and the low IA observed in kumis samples indicates that kumis is a food product that could provide protection against coronary heart diseases.

#### Estimation of daily CLA intake in Colombia

Aiming to establish the contribution of CLA intake of kumis in the Colombian diet, it was necessary to estimate the average intake of CLA in Colombia because there was no available data. This estimation was made on the basis of the National Survey of Nutritional Situation in Colombia 2005 (25) as indicated above, and the results are presented in table 4.

From the data presented in table 2, the contribution of the kumis samples to the daily CLA intake, could reach values as higher than 47%, considering the average daily CLA intake to be 254.5 mg/day. These results suggest that at least one portion of kumis could be added to the daily diet, aiming to enhance the daily CLA intake.

**Table 4.** Estimated daily intake of CLA in Colombia.

Age (years)	2 to 3	4 to 8	9 to 13	14 to 18	19 to 50	51 to 63	Average
CLA intake (mg/day)	293.5	281.1	274.5	263.2	262.2	195.1	254.5

However, when comparing the estimated daily CLA intake with the available data (Germany: 360-440; Switzerland: 160 ± 60; European Community: 380; Canada: 95 ± 41, and United States: 151 - 212 mg CLA/day) (27, 43-48), the daily CLA intake in Colombia could be higher than in Canada, similar to those reported for United States and Switzerland, and lower than the one registered for Europe.

## CONCLUSIONS

The concentration of CLA in commercial samples of kumis has been evaluated for the first time in Colombia in this study. Results demonstrated that the CLA content of the analyzed kumis samples was slightly higher than the ones previously reported for various dairy products. The calculated values of CLA intake for each kumis portion indicated that this dairy product could provide up to 47% of the estimated daily CLA intake of 254.5 mg/day. This fact suggests that, in order to increase the daily CLA intake in the Colombian population, at least one kumis portion should be added to the daily diet. The index of atherogenicity of kumis samples were low, indicating that kumis consumption could contribute to the protection against coronary heart diseases. The estimated daily CLA intake in Colombia, also calculated for the first time, indicates that CLA consumption in Colombia is similar to that of United States, but lower than the one registered for Europe.

## ACKNOWLEDGMENTS

This research was funded by the Ministerio de Agricultura y Desarrollo Rural of the República de Colombia and Instituto de Ciencia y Tecnología of the Universidad Nacional de Colombia Sede Bogotá. Additional support was provided by DANISCO® Colombia Ltda., and Productos Naturales de la Sabana S.A. ALQUERIA®. Martha S. Franco and Jairo Moreno are acknowledged for their technical assistance and collaboration.

## REFERENCES

- Garcialopez S, Echeverria E, Tsui I, Balch B. Changes in the content of conjugated linoleic-acid (CLA) in processed cheese during processing. *Food Res Int.* 1994; 27 (1): 61-64.
- Parodi PW. Conjugated octadecadienoic acids of milk-fat. *J Dairy Sci.* 1977 Oct; 60 (10): 1550-1553.
- Collomb M, Schmid A, Sieber R, Wechsler D, Ryhanen EL. Conjugated linoleic acids in milk fat: Variation and physiological effects. *Int Dairy J.* 2006 Nov; 16 (11): 1347-1361.
- Ledoux M, Chardigny JM, Darbois M, Soustre Y, Sebedio JL, Laloux L. Fatty acid composition of French butters, with special emphasis on conjugated linoleic acid (CLA) isomers. *J Food Compos Anal.* 2005 Aug; 18 (5): 409-425.
- Parodi PW. Milk fat in human nutrition. *Aust J Dairy Technol.* 2004 Apr; 59 (1): 3-59.
- Terpstra AH. Effect of conjugated linoleic acid on body composition and plasma lipids in humans: an overview of the literature. *Am J Clin Nutr.* 2004 Mar; 79 (3): 352-361.

7. Khanal RC. Potential health benefits of conjugated linoleic acid (CLA): A review. *Asian Austral J Anim.* 2004 Sep; 17 (9): 1315-1328.
8. Roche HM, Noone E, Nugent A, Gibney MJ. Conjugated linoleic acid: a novel therapeutic nutrient?. *Nutr Res Rev.* 2001 Jun 23; 14 (1): 173-187.
9. Thom E, Wadstein J, Gudmundsen O. Conjugated linoleic acid reduces body fat in healthy exercising humans. *J Int Med Res.* 2001 Sep-Oct; 29 (5): 392-396.
10. Belury MA. Dietary conjugated linoleic acid in health: Physiological effects and mechanisms of action. *Annu Rev Nutr.* 2002 Apr 4; 22: 505-531.
11. Moloney F, Yeow TP, Mullen A, Nolan JJ, Roche HM. Conjugated linoleic acid supplementation, insulin sensitivity, and lipoprotein metabolism in patients with type 2 diabetes mellitus. *Am J Clin Nutr.* 2004 Oct; 80 (4): 887-895.
12. Aneja RP, Murthi NT. Conjugated linoleic acid contents of Indian curds and ghee. *Indian J Dairy Sci.* 1990; 43 (1): 231-238.
13. Shantha NC, Ram LN, O'Leary J, Hicks CL, Decker EA. Conjugated linoleic-acid concentrations in dairy-products as affected by processing and storage. *J Food Sci.* 1995 Jul-Aug; 60 (4): 695-697.
14. Sieber R, Collomb M, Aeschlimann A, Jelen P, Eyer H. Impact of microbial cultures on conjugated linoleic acid in dairy products - A review. *Int Dairy J.* 2004 Jan; 14 (1): 1-15.
15. Kim YJ, Liu RH. Increase of conjugated linoleic acid content in milk by fermentation with lactic acid bacteria. *J Food Sci.* 2002 Jun-Jul; 67 (5): 1731-1737.
16. Lin TY. Influence of lactic cultures, linoleic acid and fructooligosaccharides on conjugated linoleic acid concentration in non-fat set yogurt. *Aust J Dairy Technol.* 2003 Apr; 58 (1): 11-14.
17. Xu S, Boylston TD, Glatz BA. Conjugated linoleic acid content and organoleptic attributes of fermented milk products produced with probiotic bacteria. *J Agr Food Chem.* 2005 Nov 16; 53 (23): 9064-9072.
18. Yadav H, Jain S, Sinha PR. Production of free fatty acids and conjugated linoleic acid in probiotic dahi containing *Lactobacillus acidophilus* and *Lactobacillus casei* during fermentation and storage. *Int Dairy J.* 2007 Aug; 17 (8): 1006-1010.
19. Pabón M, Carulla J, 136-145. Compuestos lipídicos benéficos para la salud humana asociados a la nutrición animal. *Rev Colomb Cienc Pecu.* 2008; 21 (1): 136-145.
20. Rico JM, Moreno B, Pabón ML, Carulla JE. Composición de la grasa láctea de la sabana de Bogotá con énfasis en ácido ruménico - CLA cis-9, trans-11. *Rev Colomb Cienc Pecu.* 2007; 20 (1): 30-39.
21. Folch J, Lees M, Sloane GH. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem.* 1957; 226 (1): 497-509.
22. Christie WW, Dobson G, Adlof RO. A practical guide to the isolation, analysis and identification of conjugated linoleic acid. *Lipids.* 2007 Dec; 42 (12): 1073-1084.
23. Montgomery DC. Design and analysis of experiments. 7th ed. Hoboken, NJ, United States: Wiley; 2009.
24. Ulbricht TLV, Southgate DAT. Coronary heart-disease - 7 Dietary factors. *Lancet.* 1991 Oct 19; 338 (8773): 985-992.
25. Instituto Colombiano de Bienestar Familiar. Encuesta Nacional de la Situación Nutricional en Colombia 2005 [Internet]. (Bogotá, Colombia): ICBF; 2006 [cited 2010 November 8th]; Available from: [www.icbf.gov.co/icbf/directorio/.../IENSINLIBROCOMPLETO.pdf](http://www.icbf.gov.co/icbf/directorio/.../IENSINLIBROCOMPLETO.pdf)
26. Instituto Colombiano de Bienestar Familiar. Tabla de composición de alimentos colombianos [Internet]. (Bogotá, Colombia): ICBF. [cited 2010 september 1st]; Available from: [http://alimentoscolombianos.icbf.gov.co/alimentos\\_colombianos/consulta\\_alimento.asp](http://alimentoscolombianos.icbf.gov.co/alimentos_colombianos/consulta_alimento.asp)
27. Fritsche J, Steinhart H. Amounts of conjugated linoleic acid (CLA) in German foods and evaluation of daily intake. *Z Lebensm Unters F A.* 1998; 206 (2): 77-82.
28. Boylston TD, Beitz DC. Conjugated linoleic acid and fatty acid composition of yogurt produced from milk of cows fed soy oil and conjugated linoleic acid. *J Food Sci.* 2002 Jun-Jul; 67 (5): 1973-1978.
29. Maier S, Reich E, Martin R, Bachem M, Altug V, Hautmann RE, et al. Tributyrin induces differentiation, growth arrest and apoptosis in androgen-sensitive and androgen-resistant human prostate cancer cell lines. *Int J Cancer.* 2000 Oct 15; 88 (2): 245-251.
30. Gagliostro GA, Rodríguez A, Pellegrini P, Museo G, Gatti P, Garciarena D. Effect of yogurt-making technology on the composition of fatty acids. *Rev Arg Prod Anim.* 2007; 27 (1): 352-354.
31. Coradini D, Biffi A, Costa A, Pellizzaro C, Pirronello E, Di-Fronzo G. Effect of sodium butyrate on human breast cancer cell lines. *Cell Proliferat.* 1997 Mar-Apr; 30 (3-4): 149-159.
32. Menzel T, Schaubert J, Kreth F, Kudlich T, Melcher R, Gostner A, et al. Butyrate and aspirin in combination have an enhanced effect on apoptosis in human colorectal cancer cells. *Eur J Cancer Prev.* 2002 Jun; 11 (3): 271-281.
33. Thormar H, Isaacs CE, Kim KS, Brown HR. Inactivation of visna virus and other enveloped viruses by free fatty-acids and monoglycerides. *Ann Ny Acad Sci.* 1994; 724: 465-471.
34. Collomb M, Butikofer U, Sieber R, Jeangros B, Bosset JO. Correlation between fatty acids in cows' milk fat produced in the lowlands, mountains and highlands of Switzerland and botanical composition of the fodder. *Int Dairy J.* 2002; 12 (8): 661-666.
35. Simionato JI, Garcia JC, dos Santos GT, Oliveira CC, Visentainer JV, de Souza NE. Validation of the determination of fatty acids in milk by gas chromatography. *J Brazil Chem Soc.* 2010; 21 (3): 520-524.
36. Parthasarathy S, Khoo JC, Miller E, Barnett J, Witztum JL, Steinberg D. Low-density-lipoprotein rich in oleic-acid is protected against oxidative modification - implications for dietary prevention of atherosclerosis. *P Natl Acad Sci USA.* 1990 May; 87 (10): 3894-3898.
37. Fontecha J, Recio I, Pilosof AMR, editors. Funcionalidad de los componentes lácteos. Juárez M, Fontecha J. España: CEE Limencop, S.L.; 2009. Componentes bioactivos de la grasa láctea; p. 251-273.
38. Hassan AN, Frank JF, Schmidt KA, Shalab SI. Rheological properties of yogurt made with encapsulated nonropy lactic cultures. *J Dairy Sci.* 1996 Dec; 79 (12): 2091-2097.
39. Decourcelle N, Lubbers S, Vallet N, Rondeau P, Guichard E. Effect of thickeners and sweeteners on the release of blended aroma compounds in fat-free stirred yoghurt during shear conditions. *Int Dairy J.* 2004 Sep; 14 (9): 783-789.
40. Purwandari U, Shah NP, Vasiljevic T. Effects of exopolysaccharide-producing strains of *Streptococcus thermophilus* on technological and rheological properties of set-type yoghurt. *Int Dairy J.* 2007 Nov; 17 (11): 1344-1352.
41. Ulbricht TLV, Southgate DAT. Coronary heart disease: seven dietary factors. *The Lancet.* 1991 Oct 19; 338 (8773): 985-992.
42. Huang Y, Schoonmaker JP, Bradford BJ, Beitz DC. Response of milk fatty acid composition to dietary supplementation of soy oil, conjugated linoleic acid, or both. *J Dairy Sci.* 2008 Jan; 91 (1): 260-270.
43. Fremann D, Linseisen J, Wolfram G. Dietary conjugated linoleic acid (CLA) intake assessment and possible biomarkers of CLA intake in young women. *Public Health Nutr.* 2002 Feb; 5 (1): 73-80.
44. Jahreis G, Fritsche J, Mockel P, Schone F, Moller U, Steinhart H. The potential anticarcinogenic conjugated linoleic acid, cis-9,trans-11 C18: 2, in milk of different species: Cow, goat, ewe, sow, mare, woman. *Nutr Res.* 1999 Oct; 19 (10): 1541-1549.
45. Jiang J, Wolk A, Vessby B. Relation between the intake of milk fat and the occurrence of conjugated linoleic acid in human adipose tissue. *Am J Clin Nutr.* 1999 Jul; 70 (1): 21-27.



- 46. Wolff RL, Precht D. Reassessment of the contribution of bovine milk fats to the trans-18: 1 isomeric acid consumption by European populations. Additional data for rumenic (cis-9, trans-11 18: 2) acid. *Lipids*. 2002 Dec; 37 (12): 1149-1150.
- 47. Ens JG, Ma DWL, Cole KS, Field CJ, Clandinin MT. An assessment of c9,t11 linoleic acid intake in a small group of young Canadians. *Nutr Res*. 2001 Jul; 21 (7): 955-960.
- 48. Ritzenthaler KL, McGuire MK, Falen R, Shultz TD, Dasgupta N, McGuire MA. Estimation of conjugated linoleic acid intake by written dietary assessment methodologies underestimates actual intake evaluated by food duplicate methodology. *J Nutr*. 2001 May 1; 131 (5): 1548-1554.



## WHAT'S NEW IN Vitae ?

The Scientific and Editorial Board is delighted to announce you that Vitae has been accepted for addition in Scopus.

Scopus is the largest abstract and citation database of research literature and quality web sources covering nearly 18,000 titles from more than 5,000 publishers!



---

**SINCE 1991:**








**We've improved and expanded in the last two years:**

**NEW**













**Impact Factor 2010 = 0.205**

*Thank you very much for your valuable advice and support.*

AN'ANASHA

NASHA'O