

# INHIBITION OF QUORUM SENSING BY COMPOUNDS FROM TWO *EUNICEA* SPECIES AND SYNTHETIC SATURATED ALKYLGLYCEROLS

## INHIBICIÓN DEL QUORUM SENSING POR COMPUESTOS DE DOS ESPECIES DE *EUNICEA* Y ALQUILGLICEROLES SATURADOS SINTÉTICOS

Clara BARRAGÁN A., M.Sc.<sup>1</sup>, Edelberto SILVA G., M.Sc.<sup>2</sup>,  
Bárbara MORENO M. Esp. P.N.<sup>1</sup>, Humberto MAYORGA W., PhD.<sup>1\*</sup>

Received: January 21 of 2017. Approved: August 27 of 2018

### ABSTRACT

**Background:** the emergence of bacterial resistance has led to a search for new natural products as alternatives starting points to prevent and control diseases caused by microorganisms. Among the potential candidates were bioactive compounds from octocorals of the genus *Eunicea* due to their chemical structures and their wide range of biological activities. **Objective:** the purpose of this study was to evaluate the quorum sensing inhibition (QSI) and antibacterial activity of compounds previously isolated from two *Eunicea* species and synthesized alkylglycerols (AKG). **Methods:** the QSI of three nonpolar compounds and a mixture of AKGs from *Eunicea* were evaluated by a microtiter plate assay using *Chromobacterium violaceum* (ATCC 31532). Four naturally occurring, saturated, and enantiomerically pure AKGs, all of which were derived from the chiral precursor (*R*)-solketal, were synthesized from alkyl chains of 12, 14, 16 and 18 carbons, and their structures were spectroscopically verified by NMR, ESI-MS and optical rotation data. Their QSI by the disc diffusion assay and minimum inhibitory concentrations (MIC) against 14 clinical bacterial isolates in microtiter plates were determined. **Results:** cembradiene **1**, the AKG mixture and AKG (2*S*)-3-*O*-dodecyl-1,2-propanediol **4** inhibit QS at the same concentration as kojic acid (10 µg/well or 20 µg/disc, respectively). In this study, the bioactive compounds **1**, stearyl oleate **2**, acylglycerol **3** and AKGs **4** and (2*S*)-3-*O*-tetradecyl-1,2-propanediol **5** showed *in vitro* IQS activity for the first time. Additionally, **4** and **5** displayed *in vitro* antibacterial activity against *Listeria innocua* and *Staphylococcus aureus* (MIC = 32 µg/mL for both **4** and **5**), *Enterococcus faecalis* (128 µg/mL and 64 µg/mL respectively), *Micrococcus luteus* (128 µg/mL for both) and *Brevibacillus brevis* (*Bacillus brevis*) (512 µg/mL and 64 µg/mL respectively). **Conclusion:** results suggest that natural compounds **1**, **2**, **3**, **4** and **5** showed QSI, also **4** and **5** have antibacterial activity and they are an interesting alternative to continue researching their effect against pathogenic microorganisms.

**Keywords:** Natural products, alkylglycerols, synthesis, quorum sensing inhibition, antibacterial.

<sup>1</sup> Departamento de Química, Grupo de Productos Naturales Vegetales Bioactivos y Química Ecológica, Facultad de Ciencias, Universidad Nacional de Colombia. Bogotá D.C., Colombia.

<sup>2</sup> Departamento de Farmacia, Grupo Productos Naturales Vegetales Bioactivos y Química Ecológica, Facultad de Ciencias, Universidad Nacional de Colombia. Bogotá D.C., Colombia.

\* Corresponding author: [hmayorgaw@unal.edu.co](mailto:hmayorgaw@unal.edu.co)

## RESUMEN

**Antecedentes:** la aparición de resistencia bacteriana ha llevado a la búsqueda de nuevos productos naturales como puntos de partida alternativos para prevenir y controlar las enfermedades causadas por microorganismos. Entre los posibles candidatos se encontraban los compuestos bioactivos de octocorales del género *Eunicea* debido a sus estructuras químicas y su gama amplia de actividades biológicas. **Objetivos:** el propósito de este estudio fue evaluar la inhibición del *quorum sensing* (IQS) y la actividad antibacteriana de compuestos aislados previamente de dos especies de *Eunicea* y de alquilglicerol (AQG) sintetizados. **Métodos:** la IQS de tres compuestos no polares y de una mezcla de AQGs de *Eunicea* se evaluaron mediante un ensayo de placa de microtitulación usando *Chromobacterium violaceum* (ATCC 31532). Cuatro AQG naturales, saturados y enantioméricamente puros, todos los cuales se derivaron del precursor quirral (*R*)-solketal, se sintetizaron a partir de cadenas alquílicas de 12, 14, 16 y 18 carbonos, y sus estructuras se verificaron espectroscópicamente mediante RMN, ESI-MS y datos de rotación óptica. Se determinó su IQS mediante el ensayo de difusión de disco y las concentraciones inhibitorias mínimas (CIM) en placas de microtitulación frente a 14 aislamientos bacterianos clínicos. **Resultados:** cembradieno **1**, la mezcla de AQGs y AQG (2*S*)-3-*O*-dodecil-1,2-propanodiol **4** inhiben el QS a la misma concentración que el ácido kójico (10 µg/pocillo o 20 µg/disco, respectivamente). En este estudio, los compuestos bioactivos **1**, estearil oleato **2**, acilglicerol **3** y AQGs **4** y (2*S*)-3-*O*-tetradecil-1,2-propanodiol **5** mostraron actividad de IQS *in vitro* por primera vez. Además, **4** y **5** presentaron actividad antibacteriana *in vitro* contra *Listeria innocua* y *Staphylococcus aureus* (CIM = 32 µg/mL para **4** y **5**), *Enterococcus faecalis* (128 µg/mL y 64 µg/mL respectivamente), *Micrococcus luteus* (128 µg/mL para ambos) y *Brevibacillus brevis* (*Bacillus brevis*) (512 µg/mL y 64 µg/mL respectivamente). **Conclusiones:** los resultados sugieren que los compuestos naturales **1**, **2**, **3**, **4** y **5** mostraron IQS, también **4** y **5** tenían actividad antibacteriana y son una alternativa interesante para continuar investigando su efecto contra microorganismos patógenos.

**Palabras clave:** Productos naturales, alquilglicerol, síntesis, inhibición del *quorum sensing*, antibacteriana.

## INTRODUCTION

The emergence and rapid spread of resistant bacterial strains and the overuse of antibiotics is a significant public health problem, and the development of other strategies requires searching for new natural products that are active against pathogenic microorganisms. In recent years, several known or newly discovered bioactive molecules have been evaluated against microorganisms (1-3). In the same way, quorum sensing inhibitors are potential leads for the design of new anti-pathogenic drugs (4). A process called quorum sensing (QS) is a form of population density-dependent cell-cell communication and gene regulation that allows bacteria to coordinate settlement, virulence, luminescence, metabolite production and biofilm maturation (5, 6). The interruption of QS may prevent the development of bacterial biofilms, and in this sense, it constitutes an opportunity to attenuate the pathogenicity of bacteria resistant to available antibiotics (7). In addition, promising QSI compounds have been shown to make biofilms more susceptible to antimicrobial treatments, and this indicates that a combination treatment of

both QSI and antibiotics may prove useful against resistant bacteria (8).


In previous research with sponges and octocorals from the Colombian Caribbean Sea, some biologically active compounds were identified (9-11). Two cembradiene diterpenoids and other nonpolar compounds were isolated from two octocoral species of the genus *Eunicea* collected in Santa Marta Bay (*Eunicea* sp.), and they afforded only modest activity against a variety of isolated marine bacteria from immersed fouled surfaces (10,12). However, an isolated alkyglycerol (AKG), the **7** (Figure 1), showed remarkable effectiveness against biofilm formation by three of these marine bacteria, namely, *Ochrobactrum pseudogrignonense*, *Alteromonas macleodii* and *Vibrio harveyi*, as well as against a known biofilm-forming bacterium, *Staphylococcus aureus* ATCC 25923 (12), but other AKGs could not be purified from these *Eunicea* species due to their low natural abundances.

Alkyglycerols are ether lipids naturally found in a variety of organisms such as microorganisms, fish, and invertebrates such as corals and sponges, milk, bone marrow, mammalian blood cells

including human blood cells and to a lesser extent in higher plants. AKGs have multiple biological and therapeutic properties, such as the stimulation of hematopoiesis, immunological defenses, anti-tumor activity, and anti-metastasis activity as well as improving sperm quality and vaccination efficiency (13, 14). Natural AKGs were found as complex mixtures of compounds that vary by length and the degree of unsaturation of the alkyl chain; however, the absolute configuration at their asymmetric carbon is always *S* (15). Although biological testing has been done on AKGs isolated from natural sources, the separation of individual components is tedious (16), and therefore, some of the biological assays have been performed with synthesized AKGs (14). As an example, in the

antibacterial activity studies, the synthetic and racemic AKG dodecylglycerol was the most potent against *Enterococcus faecium*, *Streptococcus mutans* and *S. aureus* (17, 18).

Continuing our studies on bioactive compounds against microorganisms, the aim of this work was to assess the QSI using *C. violaceum* of three compounds, a mixture of AKGs previously isolated from two *Eunicea* species, of four synthesized natural enantiomers of saturated AKGs (Figure 1) and evaluate the antibacterial activity of the AKGs against clinical isolates of bacteria, some of them known for their pathogenic capacity. The results exhibited that some compounds from those two octocorals may have potential for the control of bacterial infections.



**Figure 1.** Chemical structures of the natural compounds and synthetic alkylglycerols (AKG) included in this study: a cembradiene diterpenoid, (+)-(1*S*, 2*R*, 11*S*, 12*R*, 15*S*, 3*E*, 7*E*)-11-acetoxy-2,12-oxa-15,17-epoxy-cembra-3,7-diene **1**; a wax ester, stearyl oleate **2**; an acylglycerol, 1,3-dihexadecanoyl-2-(9*Z*-octadecenoyl)-glycerol **3**; the alkylglycerols (AKGs), (2*S*)-3-*O*-dodecyl-1,2-propanediol **4**; (2*S*)-3-*O*-tetradecyl-1,2-propanediol **5**; (2*S*)-3-*O*-hexadecyl-1,2-propanediol **6**; and (2*S*)-3-*O*-octadecyl-1,2-propanediol **7**.

## MATERIALS AND METHODS

### Reagents and equipment


The reagents (*R*)-(-)-2,3-*O*-isopropylidene-glycerol ((*R*)-solketal), tetra-*n*-butyl ammonium bromide (*n*-Bu<sub>4</sub>NBr), sodium bis-(2-methoxyethoxy)aluminum hydride (Red-Al), kojic acid (KA) and alkyl bromides were purchased from Alfa Aesar (USA); methanesulfonyl chloride (MsCl), *p*-toluenesulfonic acid (*p*-TsOH), *p*-hydroxybenzaldehyde (*p*-HB) and methyl octadecanoate were from Merck (Germany). All reagents were of analytical grade, and solvents were dried and distilled prior to use. TLC was performed on silica gel 60 F<sub>254</sub> (Merck) plates and visualized by UV light and 0.5 % (p/v) ceric ammonium sulfate in a 10 % sulfuric acid solution. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on an Avance 400 spectrometer (Bruker, USA) or an Avance 300, using CDCl<sub>3</sub> and TMS as internal standards. For liquid chromatography-electrospray ionization mass spectrometry (LC-ESI-MS), an LCMS-2010-ESI system (Shimadzu, Japan) was used in the positive ion mode with a RP-18 column (50 mm x 4.6 mm id x 5 μm, Waters, USA). Ten microliters of the fraction dissolved at 1mg/mL in chloroform:methanol:water:ammonium acetate 0.5:9.0:0.5:0.004 (v/v/v/v) was injected and eluted with a gradient of 9:1:0.004 methanol:water:ammonium acetate-10:0.004 methanol:ammonium acetate for 10 min at a flux of 1.5 mL/min. Additionally, 5 μL samples of pure compounds were eluted with 9:1 methanol:water-methanol. Optical rotations were measured on an ADP440 (Bellingham + Stanley, USA), and a Spectronic 20 Genesys spectrophotometer (Thermo Scientific, USA) for measuring the optical density at 600 nm was used for the evaluation of the bacterial growth.

### Analysis of compounds and a fraction from two octocorals of the genus *Eunicea*

In our earlier works, organic compounds **1**, **2**, and **3** (Figure 1) and a nonpolar fraction of two octocoral species of the genus *Eunicea* from Santa Marta Bay, Colombia, were obtained by bio-guided isolation using an antibacterial test against several marine bacteria and they were previously assigned (**10**, **12**). In this work, the natural samples were used for QSI assays, and the nonpolar fraction was subjected to further analysis by <sup>1</sup>H-NMR, <sup>13</sup>C-NMR and LC-ESI-MS.

### Synthesis of saturated alkylglycerols 4-7


Natural saturated AKGs **4-6** were synthesized to obtain enough material for their bioassays, following the procedures described in the literature (**15**, **16**, **19**). To the chiral precursor (*R*)-solketal (0.60 g, 4.54 mmol), the corresponding alkyl bromide (1.13 g, 4.54 mmol) and *n*-Bu<sub>4</sub>NBr (0.29 g, 0.91 mmol), KOH (0.32 g, 5.68 mmol) was added, and the resulting mixture was stirred for 15 hours at 40°C. The reaction was quenched by the addition of distilled water (3 mL), extracted with ethyl ether (3 x 3 mL) and washed with H<sub>2</sub>O until the pH was neutral. The organic phase was concentrated under vacuum using a rotary evaporator, and a semisolid enriched in the alkyl isopropylidene-glycerol was obtained (Scheme 1). This intermediate was deprotected in 7 mL of THF:water (15:5) with *p*-TsOH (55.4 mg, 0.30 mmol) and refluxed for 15 h. After extraction with chloroform (3x3 mL) and washing with H<sub>2</sub>O, the organic phase was dried with anhydrous MgSO<sub>4</sub> and concentrated under vacuum. The obtained solid was recrystallized from hexane at 0°C to afford pure AKGs **4**, **5** and **6**. Each step in the synthesis was monitored by TLC.



**Scheme 1:** Synthesis of alkylglycerols **4-6**.

The synthesis of **7** was performed according to methods published by some authors (16, 19, 20). Red-Al (0.83 g, 4.15 mmol) was added to methyl octadecanoate (0.62 g, 2.08 mmol) dissolved in anhydrous THF (5 mL) at 0°C and stirred for 5 h at 24°C. NaOH (10% in water) was then added at 10°C, and the organic layer was washed with water, dried and concentrated. Octadecanol was subsequently mesylated in dichloromethane with anhydrous

pyridine (0.3 mL, 4.53 mmol) at 0°C by the addition of MsCl (0.3 mL, 4.53 mmol) and stirring for 24 h at 25°C (Scheme 2). After quenching the reaction with distilled water, the organic phase was extracted with 1 N HCl, washed with water, dried, and concentrated. Finally, the newly formed mesylate was converted to AKG **7** as described above, using (*R*)-solketal (0.4 g, 2.92 mmol), *n*-Bu<sub>4</sub>NBr (0.2 g, 0.58 mmol), and KOH (0.3 g, 5.31 mmol).



**Scheme 2:** Synthesis of alkylglycerol **7**.

### Quorum sensing inhibition assays

The QSI abilities of compounds **1**, **2**, and **3** as well as the fraction from *Eunicea* sp. were evaluated *in vitro* in 96-well microtiter plates (TPP, Switzerland) (6), using *C. violaceum* (ATCC 31532) as the biosensor (4). Pre-inoculum was cultured overnight in tryptic soy broth (TSB, Merck), and concentration was adjusted to an optical density of 0.2-0.3 at 600 nm. Then, 100  $\mu$ L was added to each well, followed by 100  $\mu$ L of natural samples dissolved in dimethylsulphoxide (DMSO) at final concentrations of 5, 10, 20 and 40  $\mu$ g, respectively, and the plate was incubated for 24 hours at 28 °C. Thus, the QSI activity was established by the appearance of a colorless and opaque well but without affecting bacterial growth, and it was evaluated as the minimum quantity in  $\mu$ g per well of sample required to inhibit violacein pigment. The sample solvent (DMSO), as solvent without antimicrobial activity it served as the negative control and KA was used as the positive control. The QSI potential of synthesized AKGs was assessed using the standard disk diffusion method (21, 22), with *C. violaceum* (ATCC 31532) and Luria-Bertani

broth (LB, Merck) as growth medium. Sterile filter paper discs 5 mm in diameter (Whatman, USA) were impregnated with 10, 20, 30 and 50  $\mu$ g of each AKG dissolved in dichloromethane (DCM) and dried at room temperature for 20 minutes. The inoculum was adjusted to 10<sup>6</sup> CFU/mL (0.5 McFarland) in LB agar in Petri dishes. The discs were placed on the agar, the plates were incubated for 24-48 h at 27°C and the inhibition halo was measured. This assay also measures the minimum amount of each compound that inhibits pigment production (violacein) around the disc and indicates that the QS of *C. violaceum* was disrupted but does not interfere with bacterial growth (22). As positive controls, KA (5), and *p*-HB (28), were used at the same concentrations and treated as described above per disc, and DCM as inactive solvent to antimicrobial activity was used as a negative control.

### Evaluation of the antibacterial activity

To further investigate the antibacterial properties, the synthesized AKGs were selected since all natural compounds were isolated from these octocorals in low quantities. Their MICs were determined

*in vitro* in microtiter plates against 14 clinical bacterial isolates (23-25). Each AKG was dissolved in dimethylformamide (DMF) (1 mg/mL) to final concentrations of 1, 2, 4, 8, 16, 32, 64, 128, 256 and 512  $\mu\text{g/mL}$  in wells. Bacterial strains were grown with Mueller-Hinton agar (MHA, Merck). The inoculum (100  $\mu\text{L}$ ,  $10^4$  CFU/mL) in each well was mixed with the AKG in MH broth to 200  $\mu\text{L}$ , and the plates were incubated at 37°C for 18-24 h. An inoculated well with DMF in culture medium and blank well containing only assay medium were used as growth controls. The MIC of 64  $\mu\text{g/mL}$  of racemic dodecylglycerol against *S. aureus* (18) was used as a standard reference data point. Among the clinical isolates, the gram-positive bacteria *Listeria innocua*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Brevibacillus brevis*, and *Micrococcus luteus* as well as the gram-negative bacteria *Escherichia coli*, *Salmonella enteritidis*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Enterobacter agglomerans*, *Klebsiella oxytoca*, *Acinetobacter baumannii*, and *Proteus mirabilis* were provided by the Hospital of Neiva

(Huila), Hospital of Tunal, Hospital of Engativá and Universidad del Bosque (Bogotá).


### Statistical analysis

Bioassay determinations were replicated three times and reported as the mean values using the standard error and analysis of variance ANOVA's Tukey test. A difference was considered statistically significant when  $p < 0.05$  (26).

## RESULTS

### Analysis of compounds and a fraction from two octocorals of the genus *Eunicea*

In this work, natural compounds and a nonpolar fraction were reserved for QSI assays. The nonpolar fraction was found by NMR to be a mixture of AKGs, and by LC-MS-ESI, it was found to have the abundant ions  $[M + \text{Na}]^+$ ,  $[M + \text{H}]^+$ , and  $[M + \text{NH}_4]^+$  (Figure 2).



**Figure 2.** TIC (total ion chromatogram) obtained by LC-MS-ESI of a fraction from *Eunicea* sp., showing the separation and spectra of the major identified AKGs 6 and 7.

### Synthesis of saturated alkylglycerols 4-7

Chemical structures of the synthesized AKGs were confirmed by NMR and LC-ESI-MS spectra along with the specific optical rotations as follows.

**Compound 4.** Yield 50% as a bright white solid, (560 mg, 2.15 mmol). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>/TMS): δ 3.88-3.82 (m, 1H, H-2), 3.70 (dd, *J* = 11.4, 3.8 Hz, 1H, H-1a), 3.63 (dd, *J* = 11.4, 5.4 Hz, 1H, H-1b), 3.52 (dd, *J* = 9.8, 4.0 Hz, 1H, H-3a), 3.48 (dd, *J* = 9.8, 5.9 Hz, 1H, H-3b), 3.45 (2x dt, *J* = 9.1, 6.7 Hz, 2H, H-1'), 2.56 (s, 2H, CH<sub>2</sub>OH x 2), 1.56 (br. quint, *J* = 6.8 Hz, 2H, H-2'), 1.35-1.16 (m, 18H, CH<sub>2</sub> x 9), 0.87 (t, *J* = 6.8 Hz, 3H, H-12') ppm. <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 72.60 (C-3), 72.00 (C-1'), 70.64 (C-2), 64.40 (C-1), 32.05 (C-10'), 29.79, 29.76, 29.74, 29.72, 29.59, 29.48 (7C, C-2' and C-4'-C-9'), 26.21 (C-3'), 22.82 (C-11') 14.24 (C-12'). LC-MS-ESI *m/z* 261.05 [M + H]<sup>+</sup> (34%), 283.05 [M + Na]<sup>+</sup> (100%). [α]<sub>D</sub><sup>20</sup> +2.2 (c 4.3, CHCl<sub>3</sub>).

**Compound 5.** Yield 52% as a white solid (385 mg, 1.33 mmol). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>/TMS): δ 3.89-3.83 (m, 1H, H-2), 3.71 (dd, *J* = 11.4, 3.8 Hz, 1H, H-1a), 3.63 (dd, *J* = 11.4, 5.3 Hz, 1H, H-1b), 3.52 (dd, *J* = 9.7, 4.0 Hz, 1H, H-3a), 3.48 (dd, *J* = 9.7, 6.1 Hz, 1H, H-3b), 3.45 (2x dt, *J* = 9.3, 6.7 Hz, 2H, H-1'), 2.80 (br. s, 1H, CHOH), 2.40 (br. s, 1H, CH<sub>2</sub>OH), 1.57 (br. quint, *J* = 6.8, 2H, H-2'), 1.25 (br. s, 22H, CH<sub>2</sub> x 11), 0.87 (t, *J* = 6.8 Hz, 3H, H-14'). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 72.62 (C-3), 72.00 (C-1'), 70.60 (C-2), 64.40 (C-1), 32.06 (C-12'), 29.83, 29.81, 29.80, 29.75, 29.72, 29.71, 29.59, 29.50 (9C, C-2' and C-4'-C-11'), 26.21 (C-3'), 22.83 (C-13'), 14.26 (C-14'). LC-MS-ESI *m/z* 289.10 [M + H]<sup>+</sup> (40%), 311.10 [M + Na]<sup>+</sup> (100%). [α]<sub>D</sub><sup>20</sup> +2.1 (c 4.1, CHCl<sub>3</sub>).

**Compound 6.** Yield 27% as a white greasy solid (307 mg, 0.97 mmol). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>/TMS): δ 3.89-3.81 (m, 1H, H-2), 3.70 (dd, *J* = 11.4, 3.8 Hz, 1H, H-1a), 3.64 (dd, *J* = 11.4, 5.4 Hz, 1H, H-1b), 3.55-3.49 (m, 1H, H-3a), 3.51-3.46 (m, 1H, H-3b), 3.47-3.40 (m, 2H, H-1'), 2.20 (br. s, 2H, CHOH x 2), 1.59-1.52 (m, 2H, H-2'), 1.25 (br. s, 26H, CH<sub>2</sub> x 13), 0.87 (t, *J* = 6.7 Hz, 3H, H-16'). <sup>13</sup>C NMR and <sup>13</sup>C NMR APT (75 MHz, CDCl<sub>3</sub>) δ 72.57 (C-3), 71.98 (C-1'), 70.66 (C-2), 64.38 (C-1), 32.05 (C-14'), 29.82, 29.79, 29.74, 29.72, 29.60, 29.49 (11C, C-2' and C-4'-C13'), 26.21 (C-3'), 22.81 (C-15'), 14.23 (C-16'). LC-MS-ESI *m/z* 317.20 [M + H]<sup>+</sup> (20%), 339.15 [M + Na]<sup>+</sup> (100%). [α]<sub>D</sub><sup>20</sup> +2.5 (c 3.7, CHCl<sub>3</sub>).

**Compound 7.** Yield 25% as an amorphous solid (180 mg, 0.52 mmol). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>/TMS) δ 3.90-3.83 (m, 1H, H-2), 3.71 (dd, *J* = 11.5, 3.7 Hz, 1H, H-1a), 3.64 (dd, *J* = 11.5, 5.4 Hz, 1H, H-1b), 3.53 (dd, *J* = 9.9, 4.3 Hz, 1H, H-3a), 3.49 (dd, *J* = 9.9, 6.4 Hz, 1H, H-3b), 3.46 (2x dt, *J* = 9.5, 6.8 Hz, 2H, H-1'), 2.85 (br. s, 1H, CHOH), 2.05 (br. s, 1H, CHOH), 1.57 (br. quint, *J* = 6.8 Hz, 2H, H-2'), 1.25 (br. s, 30H, CH<sub>2</sub> x 15), 0.88 (t, *J* = 6.8 Hz, 3H, H-18'). <sup>13</sup>C NMR and <sup>13</sup>C NMR APT (100 MHz, CDCl<sub>3</sub>) δ 72.58 (C-3), 72.00 (C-1'), 70.66 (C-2), 64.37 (C-1), 32.07 (C-16'), 29.85, 29.82, 29.77, 29.75, 29.74, 29.71, 29.62, 29.51 (13C, C-2' and C-4'-C-15'), 26.22 (C-3'), 22.84 (C-17'), 14.27 (C-18'). LC-MS-ESI *m/z* 345.25 [M + H]<sup>+</sup> (31%), 367.25 [M + Na]<sup>+</sup> (100%). [α]<sub>D</sub><sup>20</sup> +2.7 (c 4.0, CHCl<sub>3</sub>).

### Quorum sensing inhibition assays

Cembradiene **1** and the mixture of AKGs presented inhibition at the same concentration as KA, the control, while **2** and **3** were less active (Table 1). It was observed that neither AKGs affected the growth of *C. violaceum*.


**Table 1.** Quorum sensing inhibition of an AKG mixture and compounds **1**, **2**, and **3** isolated from *Eumicea* sp. in *C. violaceum* (ATCC 31532) by the microtiter plate assay.

Compounds	Quorum sensing inhibition of natural samples					
	1 <sup>a</sup>	2 <sup>a</sup>	3 <sup>a</sup>	AKG mixture <sup>a</sup>	KA <sup>b</sup>	DMSO <sup>c</sup>
values in μg/well <sup>a</sup>	10	20	40	10	10	--

<sup>a</sup>Minimum quantity in μg of compound per well required to inhibit violacein pigment. <sup>b</sup>KA was used as a positive control. <sup>c</sup>Dimethylsulphoxide (DMSO) was used as a negative control. --, No inhibition was observed in the assay conditions. Data represent the mean from three independent experiments. \*The ANOVA showed a statistically significant difference between the controls and treated groups (p < 0.05).

Moreover, the results of QSI showed that AKG **4** exhibited the same minimum inhibition halo of 14 ± 1 mm at 20 μg/disc as KA, while AKG **5** and the other control, p-HB gave a halo of 12 ± 3 mm

at 50 μg/disc (Figure 3, Table 2). The *C. violaceum* growth confirmed that only they inhibited violacein production.



**Figure 3.** Inhibition halo (in mm) of QSI by the disc diffusion assay of synthesized alkylglycerols 4-7 in *C. violaceum* (ATCC 31532). KA and p-HB were positive controls. DCM was a negative control. Bars indicate the standard deviation of three replicates. Statistically significant differences ( $p < 0.05$ , ANOVA) between the treatment and the negative control (without AKG) are marked by asterisks.

**Table 2.** Quorum sensing inhibitory activity of synthesized alkylglycerols 4-7 against the biosensor strain *C. violaceum* (ATCC 31532) by the disc diffusion assay.

Compounds	Quorum sensing inhibition of AKGs						
	4*	5*	6	7	KA	p-HB <sup>b</sup>	DCM <sup>c</sup>
values in $\mu\text{g}/\text{disc}^a$	20	50	--	--	20	50	--

<sup>a</sup>Minimum quantity in  $\mu\text{g}$  of compound per disc required to inhibit violacein pigment. <sup>b</sup>KA and p-HB were used as a positive control. <sup>c</sup>Dichloromethane (DCM) was used as a negative control. --, No inhibition zone observed in the assay conditions. Only the synthesized AKGs 4-7 were selected for this assay since all natural 1, 2, 3 compounds were isolated from these two octocorals in insufficient amount. Data are the mean of three separate experiments. \*The ANOVA showed a statistically significant difference relative to controls ( $p < 0.05$ ).

### Evaluation of the antibacterial activity

The results show that AKGs 4 and 5 were the most active against five of the gram-positive bacterial strains (Table 3), while against the gram-negative strains *S. enteritidis*, *K. pneumonia*,

*E. agglomerans*, *K. oxytoca*, and *A. baumannii*, the AKGs were inactive. However, AKG 4 was active against *P. aeruginosa* and *P. mirabilis* at the highest concentration assayed.

**Table 3.** Growth inhibitory activities of synthesized alkylglycerols (AKGs) 4-7 against clinical bacterial isolates by the microtiter plate assay.

Clinical isolates	Minimum inhibitory concentrations (MICs) of AKGs ( $\mu\text{g}/\text{mL}$ )			
	4*	5*	6*	7*
Gram-positives				
<i>L. innocua</i>	32	32	512	512
<i>S. aureus</i>	32 <sup>a</sup>	32	512	512
<i>E. faecalis</i>	128	64	512	512
<i>B. brevis</i>	512	64	512	512
<i>M. luteus</i>	128	128	64	512
Gram-negatives				
<i>E. coli</i>	--	--	512	<sup>b</sup>
<i>P. aeruginosa</i>	512	--	--	512
<i>S. marcescens</i>	--	--	32	--
<i>P. mirabilis</i>	512	512	--	<sup>b</sup>

<sup>a</sup>Racemic dodecylglycerol has an MIC of 64  $\mu\text{g}/\text{mL}$  against *S. aureus* (17), and this was used as a reference value. A treatment with DMF in culture medium without compound was used as bacterial growth control, while the treatment containing only assay medium served as blank control. --, Not detected under the assay conditions. Data represent the mean of three different experiments. \*The ANOVA revealed significant differences ( $p < 0.05$ ) between the tests and the growth control of each of these bacteria. <sup>b</sup>No significant difference was observed between treated and the bacterial growth control.



## DISCUSSION

In this work, natural compounds **1**, **2**, and **3** as well as the AKG mixture were subjected to QSI assays. The LC-MS-ESI of the AKG mixture had ions corresponding with molecular masses of 316 g/mol and 344 g/mol, respectively, and thus, **6** and **7** were identified in a 1:2 ratio as the major AKGs, along with other probable saturated AKGs with lower masses (Figure 2). Recently, only **7** had been purified (12), and unfortunately, we were unable to investigate the effects of all compounds from the two octocorals in the QSI and antibacterial assays because some of them were available in low quantities. Particularly, alkylglycerols (AKGs) are found in low concentrations in marine sponges and corals and at trace levels in mammal, including humans (16). Previous studies were developed with the most abundant AKGs or their mixtures, and recent advances in AKG synthesis have been crucial for the exploration of their biological activities (13, 14). In this work, the most common natural enantiomers of saturated AKGs **4-7** (13), were synthesized (Figure 1). One of the most suitable and well-established routes for their preparation is from a chiral precursor with an alkyl halide or sulfonate (15, 16, 19, 20). Thus, yields of the AKGs were satisfactory, ranging from 25 % to 52 %, as no further chromatographic purifications were performed, and sufficient material was obtained for the biological tests. Their chemical structures were confirmed by spectroscopic data with those previously published. Mass spectrometry analysis of compound **4** by LC-MS-ESI showed the followed two peaks:  $m/z$  261.05 (34 %) corresponding to the ion  $[M + H]^+$  and  $m/z$  283.05 (100 %) corresponding to the ion  $[M + Na]^+$ , which are consistent with a molecular mass of 260.05 g/mol and formula of  $C_{15}H_{32}O_3$ . The positive optical rotation,  $[\alpha]^{20}_D +2.2$  (c 4.3,  $CHCl_3$ ), is typical of the *S* configuration of the natural AKGs (16), confirming its optically purity, and the NMR data were similar to those obtained for AKGs **6** and **7**, allowing **4** to be assigned as (2*S*)-3-*O*-dodecyl-1,2-propanediol, a natural AKG previously synthesized (13). Compound **5** had the formula  $C_{17}H_{36}O_3$ , by LC-MS-ESI. Its optical rotation and NMR spectra were consistent with those of **4**, and it was confirmed to be (2*S*)-3-*O*-tetradecyl-1,2-propanediol, a natural and previously synthesized AKG (13). Compound **6** was  $C_{19}H_{40}O_3$ , its optical

rotation and NMR data are similar to those of the AKG synthesized (15), and thus, it was named (2*S*)-3-*O*-hexadecyl-1,2-propanediol, a natural compound known as chimyl alcohol. Compound **7** had the formula  $C_{21}H_{44}O_3$ , and its optical rotation and NMR data are in agreement with those of the natural batyl alcohol (12) previously synthesized (15); therefore, it was assigned as (2*S*)-3-*O*-octadecyl-1,2-propanediol.

In relation to the natural samples tested for QSI by the microtiter plate assay, compounds **1**, **2**, and **3** as well as the mixture of AKGs display activity. **1** and the AKG mixture presented the largest inhibition at the same concentration as the positive control, KA (10  $\mu$ g/well), while **2** and **3** were less active. Compound **1** was first isolated from a specimen of *Eunicea* collected near Providencia Island and displayed antiplasmodial activity against *Plasmodium falciparum* (27). In our recent research, we isolated **1** again from *Eunicea* sp. collected in Santa Marta Bay, and its absolute configuration was determined, but it showed weak antimicrobial activity against marine bacteria (10). In this work, **1** displayed significant QSI and was as active as KA, the positive control used and a known inhibitor of QS systems (5). Other *Eunicea* and *Pseudoplexaura* cembranoid diterpenes have shown prominent QSI activity (11). Surprisingly, the nonpolar fraction was as active as KA or as **1**, and since this fraction is a mixture of alkylglycerols, it indicated that natural AKGs could be interesting QS inhibitors, which was another reason that motivated us to synthesize some of its most likely constituents, four of the major natural saturated alkylglycerols (13). Compound **2** was previously biosynthesized and it has been detected in the eyelids of humans and mammals. It was isolated from *Eunicea* sp. and showed biofilm inhibition up to 91 % against *S. aureus* ATCC 25923 (12). **3** only has been isolated from *Eunicea* sp., and it was effective against biofilm formation by *V. harveyi* (Phy-2A) and *P. aeruginosa* (ATCC 27853), with a percentage higher than 25 % (12). In this work, compounds **2** and **3** showed QSI at 20 and 40  $\mu$ g/well, respectively, but they were less active than KA.

When synthetic AKGs were evaluated by the disc diffusion assay, **4** was the most active, exhibiting the same QSI as KA (20  $\mu$ g/disc), followed by **5**, which had the same QSI as p-HB at 50  $\mu$ g/disc, all of them without affecting the microbial growth of *C. violaceum* (ATCC 31532). Meanwhile, **6** and

**7** were practically inactive to the concentrations assayed. AKGs **4** and **5** could be responsible for the QSI activity determined in the AKG mixture from *Eunicea* sp. In addition to KA, p-HB was selected as a control because it demonstrated QSI against *C. violaceum* (ATCC 31532) (28), and it is a natural component of vanilla extract, which is also a QS inhibitor (29). Other nonpolar and natural compounds that can interfere with QS include, furanones, chlorinated metabolites (called honaucins), unsaturated lactones (nocapyrones), furanosesquiterpenoid (felixinin), sesquiterpene alcohol (farnesol), unsaturated fatty acids (pitinoic acid and *cis*-9-octadecenoic acid), and a cyclopropyl fatty acid (lyngbyoic acid) (30, 31). In the current work, the bioactive natural and known compounds, **1**, **2**, **3**, **4** and **5** displayed *in vitro* QSI for the first time.

The antibacterial potential of the synthesized AKGs was evaluated against clinical isolates of bacteria, and the MICs were determined in a microtiter plate assay. In this way, **4** and **5** showed *in vitro* antibacterial activity against five of the gram-positive strains: *L. innocua* and *S. aureus* (MICs of 32  $\mu\text{g}/\text{mL}$ ), with less effectiveness against *E. faecalis*, *B. brevis* and *M. luteus*. The AKGs, **4** and **5**, were much less active against gram-negative clinical isolates, as shown in Table 3. These AKGs showed bacteriostatic effects on the clinical isolates because evaluation of the viability by seeding in fresh culture medium led to the recovery of each bacterial species (data not shown). The results from this study are consistent with previous observations for synthesized and racemic AKGs with chains of 8, 10, 12, 14 and 16 carbons, and among them, racemic dodecylglycerol was an effective antibacterial against the gram-positive strains of *S. aureus*, *E. faecium*, and *S. mutans*. It also showed low activity against the gram-negative bacteria *P. aeruginosa* and *K. pneumonia* (17, 18). However, in the present work, a MIC value of 32  $\mu\text{g}/\text{mL}$  was found for the pure enantiomer **4** against *S. aureus*, which was better than that of racemic dodecylglycerol with the published value of 64  $\mu\text{g}/\text{mL}$  (17). Nevertheless, previously **7** showed low activity against the gram-negative marine bacteria *Pseudoalteromonas piscida*, *Ruegeria* sp., *Vibrio alginolyticus*, *V. furnissii* and *V. harveyi* (32). We have reported that **7** isolated from *Eunicea* sp. had moderate activity against the gram-negative marine bacteria *O. pseudogrignonense* (4-4DEP) and *A. macleodii* (29-C), the reference

strain *P. aeruginosa* ATCC 27853, and even the gram-positive *S. aureus* ATCC 25923 (12). Many other natural lipids show antimicrobial activity (14, 17, 30), but the considerable resistance of gram-negative clinical isolates to AKGs could be attributed to their outer membrane, which creates a protective barrier against hydrophobic compounds such as lipids (33). Among the strains that were susceptible to AKGs, *L. innocua*, which is widely distributed in nature, is able to overcome extreme conditions of pH, temperature, and salinity, and although it is not considered pathogenic, it has caused some infections in humans requiring treatments with prolonged doses of antibiotics (34). Its powerful ability to grow in certain foods has been suggested to be an expression of QS (35). *S. aureus* is a widespread pathogen and known food contaminant that has developed antibiotic resistance and can cause a variety of QS-regulated infectious diseases (36,37). *E. faecalis* is a multiantibiotic resistant pathogen that resides in the human gastrointestinal tract, but studies have proven the potential for developing QS antagonists that could help control this pathogen (38). *M. luteus* has been identified from several sources, such as human tissues, and although it is not considered pathogenic, is it characterized by withstanding various environmental conditions (39). *B. brevis* is found in soils, air, and water, and it exhibits moderate resistance (40).

Present research provides additional information. It constitutes the first report of the QSI activity of natural compounds **1**, **2**, **3**, **4** and **5**. Additionally, the optically pure AKGs **4** and **5** showed more potent antimicrobial activity against *L. innocua* and *S. aureus* with a MIC of 32  $\mu\text{g}/\text{mL}$ , and they were active against *E. faecalis*, *B. brevis*, and *M. luteus*.

The main limitation of this research was that the natural compounds were available in low quantities; however, these compounds may be an alternative for new studies against pathogenic microorganisms, and the results motivated us to produce other natural AKGs because of their relatively simple structures and short chemical synthesis. Also, these advances open the possibilities to obtain by synthesis that class of compounds, specifically the alkylglycerols, since they have already found some applications in several industrial and pharmaceutical domains and could be considered as promising compounds either from an economic or an ecological point of view (13, 14).

## CONCLUSIONS

This study illustrated, for the first time, the importance of bioactive natural compounds **1**, **2**, **3**, and AKGs **4** and **5** as QS inhibitors in *C. violaceum* (ATCC 31532). The naturally occurring enantiomers of four AKGs (**4-7**) were synthesized. Cembradiene **1** together with the AKG mixture from *Eunicea* and AKG **4** inhibit QS at the same concentration as KA. In addition, **4** and **5** displayed specific antibacterial activity against gram-positive clinical isolates in the following order: *L. innocua* and *S. aureus* (MIC = 32 µg/mL for both **4** and **5**), *E. faecalis* (128 µg/mL and 64 µg/mL respectively), *M. luteus* (128 µg/mL for both) and *B. brevis* (512 µg/mL and 64 µg/mL, respectively). The interesting results show that additional studies are necessary to explore the prospects of controlling pathogenic microorganisms with these natural compounds or other natural AKGs obtained by synthesis.

## CONFLICTS OF INTEREST

The authors declare that there is no conflict of interest.

## ACKNOWLEDGEMENTS

The authors thank the National university of Colombia and Colciencias for financial support. The Ministry of Environment, Housing and Territorial Development (MAVDT) granted permission for research on marine organisms [No. 4 of 10/02/2010, and No. 0306 of 22/02/2011]. We are grateful to the research group Studies and use of marine natural products and Colombian fruits, for their collaboration.

## AUTHORS' CONTRIBUTIONS

CBA performed the experimental work. ESG supervised the biological assays and contributed to the critical reading of the manuscript. BMM made contributions to the preparation of the paper as a leader of the research team. HMW supervised the research project and the drafting of the article. All authors approved the final version of the manuscript.

## REFERENCES

1. El Arbi M, Théolier J, Pigeon P, Jellali K, Trigui F, Top S, Aifa S, Fliess I, Jaouen G, Hammami R. Antibacterial properties and mode of action of new triaryl butane citrate compounds. *Eur J Med Chem.* 2014 April; 76: 408-13.
2. Benzekri R, Bouslama L, Papetti A, Snoussi M, Benslimene I, Hamami M, Limam F. Isolation and identification of an antibacterial compound from *Diplotaxis harra* (Forssk.) Boiss. *Ind Crops Prod.* 2016 February; 80: 228-34.
3. Habbu P, Warad V, Shastri R, Madagundi S, Kulkarni VH. Antimicrobial metabolites from marine microorganisms. *Chin J Nat Med.* 2016 February; 14 (2): 101-16.
4. Brango-Vanegas J, Costa GM, Ortman CF, Schenkel EP, Reginatto FH, Ramos FA, Arevalo-Ferro C, Castellanos L. Glycosylflavonoids from *Cecropia pachystachya* Trécul are quorum sensing inhibitors. *Phytomedicine.* 2014 April; 21 (5): 670-75.
5. Dobretsov S, Teplitski M, Bayer M, Gunasekera S, Proksch P, Paul VJ. Inhibition of marine biofouling by bacterial quorum sensing inhibitors. *Biofouling.* 2011 September; 27 (8): 893-05.
6. Maisarah NS, Yin WF, Chan KG. Caffeine as a potential quorum sensing inhibitor. *Sensors.* 2013 April; 13 (4): 5117-29.
7. Castillo-Juárez I, Maeda T, Mandujano-Tinoco E, Tomás M, Pérez-Eretza B, García-Contreras S, Wood TK, García-Contreras R. Role of quorum sensing in bacterial infections. *World J Clin Cases.* 2015 July; 3 (7): 575-98.
8. Rasmussen TB, Givskov M. Quorum-sensing inhibitors as anti-pathogenic drugs. *Int J Med Microbiol.* 2006 April; 296 (2-3): 149-61.
9. Castellanos LH, Mayorga HW, Duque CB. Estudio de la composición química y actividad antifouling del extracto de la esponja marina *Cliona delitrix*. *Vitae.* 2010 may-august; 17 (2): 209-24.
10. Mayorga H, Urrego NF, Castellanos L, Duque, C. Cembradienes from the Caribbean Sea whip *Eunicea* sp. *Tetrahedron Lett.* 2011 May; 52: 2515-18.
11. Tello E, Castellanos L, Duque C, Synthesis of cembranoid analogues and evaluation of their potential as quorum sensing inhibitors. *Bioorg Med Chem.* 2013 January; 21 (1): 242-56.
12. Martínez YD, Vanegas GL, Reina LG, Mayorga HW, Arévalo-Ferro C, Ramos FR, Duque CB, Castellanos LH. Biofilm inhibition activity of compounds isolated from two *Eunicea* species collected at the Caribbean Sea. *Rev Bras Farmacogn.* 2015 November-December; 25 (6): 605-11.
13. Deniau AL, Mosset P, Pédrone F, Mitre R, Le Bot D, Legrand AB. Multiple beneficial health effects of natural alkylglycerols from shark liver oil. *Mar Drugs.* 2010 July; 8 (7): 2175-84
14. Iannitti T, Palmieri B. An Update on the therapeutic role of alkylglycerols. *Mar Drugs.* 2010 August; 8 (8): 2267-300.
15. Magnusson CD, Gudmundsdottir AV, Haraldsson GG. Chemoenzymatic synthesis of a focused library of enantiopure structured 1-O-alkyl-2,3-diacyl-sn-glycerol type ether lipids. *Tetrahedron.* 2011 March; 67 (10): 1821-36.
16. Pemha R, Pegnyemb DE, Mosset P. Synthesis of (Z)-(20R)-1-O-(20-methoxynonadec-100-enyl)-sn-glycerol, a new analog of bioactive ether lipids. *Tetrahedron.* 2012 April; 68 (14): 2973-83
17. Ved HS, Gustov E, Mahadevans V, Pieringer RA. Dodecylglycerol a new type of antibacterial agent which stimulates autolysin activity in *Streptococcus faecium* ATCC 9790. *J Biol Chem.* 1984 July; 259 (13): 8115-21.
18. Lin YC, Schlievert PM, Anderson MJ, Fair CL, Schaeffers MM, Muthyala R, Peterson ML. Glycerol monolaurate and dodecylglycerol effects on *Staphylococcus aureus* and toxic shock syndrome toxin-1 *in vitro* and *in vivo*. *PLoS One* 2009 October, 4 (10), e7499.

19. Magnusson CD, Haraldsson GG. Synthesis of enantiomerically pure (Z)-(2'R)-1-O-(2'-methoxyhexadec-4'-enyl)-sn-glycerol present in the liver oil of cartilaginous fish Tetrahedron: Asymmetry. 2010 December; 21 (23): 2841-47.
20. Hanus L, Abu-Lafi S, Fride E, Breuer A, Vogel Z, Shalev DE, Kustanovich I, Mechoulam R. 2-Arachidonyl glyceryl ether, an endogenous agonist of the cannabinoid CB1 receptor. Proc Natl Acad Sci USA. 2001 March; 98 (7): 73662-65.
21. McLean RJ, Pierson LS, Fuqua C. A simple screening protocol for the identification of quorum signal antagonists. J Microbiol Methods. 2004 September; 58 (3): 351-60.
22. Zhu H, He CC, Chu QH. Inhibition of quorum sensing in *Chromobacterium violaceum* by pigments extracted from *Auricularia auricular*. Lett Appl Microbiol. 2011 March; 52 (3): 269-74.
23. Baker CN, Thornsberry C, Ronald W, Hawkinson RW. Inoculum standardization in antimicrobial susceptibility testing: Evaluation of overnight agar cultures and the rapid inoculum standardization system. J Clin Microbiol. 1983 March; 17 (3): 450-57.
24. Zgoda JR, Porter JR. A convenient microdilution method for screening natural products against bacteria and fungi. Pharm Biol. 2001 September; 39 (3): 221-25.
25. Balouiri M, Sadiki M, Ibensouda SK. Methods for in vitro evaluating antimicrobial activity: A review. J Pharm Anal. 2016 April; 6 (2): 71-79.
26. Montgomery, D.C. Design and Analysis of Experiments. 2nd edition. Mexico: Limusa; 2007. 750 p.
27. Wei X, Rodríguez AD, Baran P, Raptis RG, Sánchez JA, Ortega-Barria E, González J. Antiplasmodial cembradiene diterpenoids from a Southwestern Caribbean gorgonian octocoral of the genus *Eunicea*. Tetrahedron 2004 December; 60 (51): 11813-19.
28. Martínez DM, Pinzón AE, Laiton MF, Cárdenas JM, Duque CB, Castellanos LH, Ramos FR. Searching compounds with biological activity by bacteria from marine environments. Abstracts, V Congreso Iberoamericano de Productos Naturales. Bogotá. Colombia, 2016. 293 p.
29. Choo, JH, Rukayadi Y, Hwang JK. Inhibition of bacterial quorum sensing by vanilla extract. Lett Appl Microbiol. 2006 April; 42 (6): 637-41.
30. Blunt, JW, Copp BR, Keyzers RA, Munro MH, Prinsep MR. Marine natural products. Nat Prod Rep. 2017 March; 34 (3): 235-94.
31. LaSarre B, Federle MJ. Exploiting quorum sensing to confuse bacterial pathogens. Microbiol Mol Biol Rev. 2013 March; 77 (1): 73-11.
32. Qi SH, Zhang S, Yang LH, Qian PY. Antifouling and antibacterial compounds from the gorgonians *Subergorgia suberosa* and *Scripearia gracilis*. Nat Prod Rep. 2008 July; 22 (2): 154-66.
33. Liu H, Pei H, Han Z, Feng G, Li D. The antimicrobial effects and synergistic antibacterial mechanism of the combination of  $\epsilon$ -Polylysine and nisin against *Bacillus subtilis*. Food Control, 2015 January; 47: 444-50.
34. Favaro M, Sarmati L, Sancesario G, Fontana C. First case of *Listeria innocua* meningitis in a patient on steroids and etanercept. JMM Case Rep. 2014 June; 1: 1-5.
35. Carvalheira A, Eusébio C, Silva J, Gibbs P, Teixeira P. Influence of *Listeria innocua* on the growth of *Listeria monocytogenes*. Food Control 2010 November; 21 (11): 1492-96.
36. Oulahlal N, Brice W, Martial A, Degraeve P. Quantitative analysis of survival of *Staphylococcus aureus* or *Listeria innocua* on two types of surfaces: Polypropylene and stainless steel in contact with three different dairy products. Food Control. 2008 February; 19 (2): 178-85.
37. Painter KL, Krishna A, Wigneshweraraj S, Edwards A.M. What role does the quorum-sensing accessory gene regulator system play during *Staphylococcus aureus* bacteremia?. Trends Microbiol. 2014 December; 22 (12): 676-85.
38. Pereira UA, Barbosa LC, Maltha CR, Demuner AJ, Masood MA, Pimenta AL. Inhibition of *Enterococcus faecalis* biofilm formation by highly active lactones and lactams analogues of rubrolides. Eur J Med Chem. 2014 July; 82: 127-38.
39. Mauclair L, Egli M. Effect simulated microgravity on growth and production of exopolymeric substances of *Micrococcus luteus* space and earth isolates. FEMS Immunol Med Microbiol. 2010 August; 59 (3): 350-56.
40. Yilmaz M, Soran H, Beyatli Y. Antimicrobial activities of some *Bacillus* spp. strains isolated from the soil. Microbiol Res. 2006 August; 161 (2): 127-31.